University of Copenhagen
Faculty of Science
Department of Chemistry

Optimized Virtual Orbital Space for Quantum Computing

Master's Thesis in Chemistry / Quantum Chemistry

Author
Tobias ⟨*Surname*⟩
xxx@ku.dk

Supervisor
⟨Supervisor Name⟩
Department of Chemistry, UCPH

Co-supervisor (if any)
⟨Co-supervisor Name⟩

Copenhagen, March 6, 2026

Abstract

⟨*Write the abstract here after all chapters are complete.*⟩

Keywords: Optimized virtual orbital space, OVOS, MP2, orbital optimisation, virtual orbital reduction, quantum computing, PySCF, Newton-Raphson.

Resumé

⟨*Danish translation of the abstract.*⟩

Acknowledgements

⟨*Write acknowledgements here.*⟩

List of Abbreviations

HF	Hartree-Fock
RHF	Restricted Hartree-Fock
UHF	Unrestricted Hartree-Fock
MP2	Second-Order Møller-Plesset Perturbation Theory
MPn	Møller-Plesset Perturbation Theory of order n
CI	Configuration Interaction
CISD	Configuration Interaction Singles and Doubles
FCI	Full Configuration Interaction
CC	Coupled Cluster
CCSD	Coupled Cluster Singles and Doubles
MCSCF	Multi-Configurational Self-Consistent Field
CASSCF	Complete Active Space Self-Consistent Field
DFT	Density Functional Theory
OVOS	Optimized Virtual Orbital Space
FNO	Frozen Natural Orbitals
PNO	Pair Natural Orbitals
DLPNO	Domain-Based Local Pair Natural Orbitals
NO	Natural Orbitals
VQE	Variational Quantum Eigensolver
QPE	Quantum Phase Estimation
JW	Jordan-Wigner
BK	Bravyi-Kitaev
NISQ	Noisy Intermediate-Scale Quantum

QC	Quantum Computing
BO	Born-Oppenheimer
AO	Atomic Orbital
MO	Molecular Orbital
SCF	Self-Consistent Field
LCAO	Linear Combination of Atomic Orbitals
ERI	Electron Repulsion Integral
CBS	Complete Basis Set
GTO	Gaussian-Type Orbital
STO	Slater-Type Orbital
DIIS	Direct Inversion in the Iterative Subspace
NR	Newton-Raphson
RLE	Reduced Linear Equation
PySCF	Python-based Simulations of Chemistry Framework
UCPH	University of Copenhagen

Contents

Abstract iii

Resumé v

Acknowledgements vii

List of Abbreviations ix

1 Introduction 1
- 1.1 Motivation . . . 1
- 1.2 The Virtual Orbital Space Problem . . . 1
- 1.3 Quantum Computing Context . . . 1
- 1.4 Objectives . . . 1
- 1.5 Thesis Outline . . . 1

2 Background 3
- 2.1 The Electronic Structure Problem . . . 3
- 2.2 Hartree-Fock Theory . . . 3
  - 2.2.1 Restricted and Unrestricted Formalism . . . 3
- 2.3 Molecular Orbital Integrals . . . 3
  - 2.3.1 One-Electron Integrals . . . 3
  - 2.3.2 Two-Electron Integrals . . . 3
  - 2.3.3 Integral Notation: Chemists' vs. Physicists' Conventions . . . 3
- 2.4 Møller-Plesset Perturbation Theory . . . 3
  - 2.4.1 MP2 Correlation Energy . . . 3
  - 2.4.2 Spin-Orbital Formulation . . . 3
- 2.5 Spin-Orbital Formalism . . . 3
- 2.6 Orbital Space Partitioning and Index Notation . . . 3
- 2.7 Brueckner Orbitals . . . 3
  - 2.7.1 Definition and Physical Interpretation . . . 3
  - 2.7.2 Brueckner Orbitals in Coupled Cluster Theory . . . 3
  - 2.7.3 Connection to Orbital-Optimised MP2 . . . 3
- 2.8 Quantum Computing and Orbital Space . . . 4
  - 2.8.1 Second Quantisation and Qubit Mapping . . . 4
  - 2.8.2 The Variational Quantum Eigensolver . . . 4
  - 2.8.3 OVOS Orbitals as a VQE Reference State . . . 4

3 Theory **5**
3.1 The Second-Order Hylleraas Functional . . . 5
3.2 Orbital Rotation Parameterisation . . . 5
3.2.1 Active and Inactive Virtual Subspaces . . . 5
3.2.2 Unitary Transformation via Matrix Exponential . . . 5
3.3 Gradient of J_2 with Respect to Orbital Rotations . . . 5
3.3.1 MP2 Virtual-Virtual Density Matrix . . . 5
3.3.2 Gradient Expression . . . 5
3.4 Hessian of J_2 with Respect to Orbital Rotations . . . 5
3.4.1 Block Structure of the Hessian . . . 5
3.4.2 Negative Eigenvalues and Level-Shift Stabilisation . . . 5
3.5 Newton-Raphson Orbital Optimisation . . . 5
3.6 Generating the Unitary Rotation Matrix . . . 5
3.7 Orbital Canonicalisation . . . 5
3.8 Energy Recovery and the OVOS Error Metric . . . 5
3.9 OVOS Orbitals as Approximate Brueckner Orbitals . . . 6
3.9.1 Orbital Stationarity and the Brueckner Condition . . . 6
3.9.2 Relationship to Orbital-Optimised MP2 . . . 6
3.9.3 Expected Reduction of T1 Amplitudes . . . 6
3.10 OVOS Molecular Orbitals as a Trial State for VQE . . . 6
3.10.1 Reference-State Overlap with the FCI Ground State . . . 6
3.10.2 Reduction of T1 Amplitudes and Circuit Depth . . . 6
3.10.3 Qubit Resource Savings . . . 6

4 Method **7**
4.1 Algorithm Overview . . . 7
4.2 Integral Transformation . . . 7
4.2.1 AO-to-MO Transformation . . . 7
4.2.2 Spin-Orbital Construction . . . 7
4.2.3 Notation Convention: Chemists' vs. Physicists' Integrals . . . 7
4.3 Initial Active Space Selection . . . 7
4.4 MP1 Amplitudes and MP2 Energy . . . 7
4.5 Gradient and Hessian Assembly . . . 8
4.5.1 Block Structure and Efficient Inversion . . . 8
4.5.2 Level-Shift Stabilisation . . . 8
4.6 Orbital Rotation and Update . . . 8
4.7 Convergence and Iteration Control . . . 8
4.8 Initialisation Strategies . . . 8
4.8.1 Default SCF Initialisation . . . 8
4.8.2 Warm-Start from Previous Orbital Set . . . 8
4.8.3 Random Unitary Sampling . . . 8

5 Computational Details **9**
5.1 Software and Libraries . . . 9
5.2 Hardware . . . 9
5.3 Molecular Geometries . . . 9
5.4 Basis Sets . . . 9
5.5 Reference SCF Calculations . . . 9

5.6 OVOS Calculation Settings . . . 9
5.7 Data Storage and Reproducibility . . . 9

6 Results **11**
6.1 Validation of the Implementation . . . 11
6.2 Energy Recovery as a Function of Active Virtual Orbital Count . . . 11
6.3 Convergence Behaviour . . . 11
6.4 Effect of Initialisation Strategy . . . 12
6.5 Basis Set Dependence . . . 12
6.6 Comparison with Adamowicz and Bartlett (1987) . . . 12
6.7 Computational Cost . . . 12
6.8 T1 Amplitudes and the Brueckner Orbital Connection . . . 12
6.9 OVOS Orbitals as a VQE Reference State: Resource Estimates . . . 12

7 Conclusion **13**
7.1 Summary of Findings . . . 13
7.2 Critical Assessment . . . 13
7.3 Implications for Quantum Computing . . . 13

8 Outlook **15**
8.1 Algorithmic Improvements . . . 15
8.2 Method Extensions . . . 15
8.3 Quantum Computing Applications . . . 15
8.3.1 OVOS Orbitals as a VQE Reference State . . . 15
8.3.2 Brueckner Orbitals and Circuit Depth on NISQ Devices . . . 15
8.3.3 Qubit Count Reduction and Fault-Tolerant Perspectives . . . 15
8.4 Larger Molecular Systems . . . 15
8.5 Benchmarking and Validation . . . 15

Bibliography **17**

A Mathematical Derivations **19**
A.1 Spin-Orbital MP2 Energy from the Hylleraas Functional . . . 19
A.2 Derivation of the Orbital Rotation Gradient . . . 19
A.3 Derivation of the Orbital Rotation Hessian . . . 19
A.4 Matrix Exponential for the Rotation Matrix . . . 19
A.5 Notation Conventions for Two-Electron Integrals . . . 19

B Additional Data and Figures **21**
B.1 Full MP2 Reference Energies . . . 21
B.2 Complete OVOS Energy Recovery Tables . . . 21
B.3 Convergence Plots for All Systems . . . 21
B.4 Hessian Eigenvalue Analysis . . . 21

C Implementation Details and Source Code **23**
C.1 Repository and Version Information . . . 23
C.2 Spin-Orbital Integral Conversion . . . 23
C.3 MP2 Energy and Amplitude Computation . . . 23

C.4 Orbital Optimisation: Gradient and Hessian 23
C.5 Profiling Results 23

List of Figures

List of Tables

5.1 Molecular geometries and electronic structure parameters used in this work. 9

A.1 Summary of two-electron integral notation used in this thesis. 19

Chapter 1

Introduction

1.1 Motivation

1.2 The Virtual Orbital Space Problem

1.3 Quantum Computing Context

1.4 Objectives

1.5 Thesis Outline

Chapter 2

Background

2.1 The Electronic Structure Problem

2.2 Hartree-Fock Theory

2.2.1 Restricted and Unrestricted Formalism

2.3 Molecular Orbital Integrals

2.3.1 One-Electron Integrals

2.3.2 Two-Electron Integrals

2.3.3 Integral Notation: Chemists' vs. Physicists' Conventions

2.4 Møller-Plesset Perturbation Theory

2.4.1 MP2 Correlation Energy

2.4.2 Spin-Orbital Formulation

2.5 Spin-Orbital Formalism

2.6 Orbital Space Partitioning and Index Notation

2.7 Brueckner Orbitals

2.7.1 Definition and Physical Interpretation

2.7.2 Brueckner Orbitals in Coupled Cluster Theory

2.7.3 Connection to Orbital-Optimised MP2

2.8 Quantum Computing and Orbital Space

2.8.1 Second Quantisation and Qubit Mapping

2.8.2 The Variational Quantum Eigensolver

2.8.3 OVOS Orbitals as a VQE Reference State

Chapter 3

Theory

3.1 The Second-Order Hylleraas Functional

3.2 Orbital Rotation Parameterisation

3.2.1 Active and Inactive Virtual Subspaces

3.2.2 Unitary Transformation via Matrix Exponential

3.3 Gradient of J_2 with Respect to Orbital Rotations

3.3.1 MP2 Virtual-Virtual Density Matrix

3.3.2 Gradient Expression

3.4 Hessian of J_2 with Respect to Orbital Rotations

3.4.1 Block Structure of the Hessian

3.4.2 Negative Eigenvalues and Level-Shift Stabilisation

3.5 Newton-Raphson Orbital Optimisation

3.6 Generating the Unitary Rotation Matrix

3.7 Orbital Canonicalisation

3.8 Energy Recovery and the OVOS Error Metric

3.9 OVOS Orbitals as Approximate Brueckner Orbitals

3.9.1 Orbital Stationarity and the Brueckner Condition

3.9.2 Relationship to Orbital-Optimised MP2

3.9.3 Expected Reduction of T1 Amplitudes

3.10 OVOS Molecular Orbitals as a Trial State for VQE

3.10.1 Reference-State Overlap with the FCI Ground State

3.10.2 Reduction of T1 Amplitudes and Circuit Depth

3.10.3 Qubit Resource Savings

Chapter 4

Method

4.1 Algorithm Overview

Algorithm 1 OVOS Iterative Optimisation

```
Perform SCF calculation to obtain initial MO coefficients C
Transform AO integrals to MO basis: ⟨ij|ab⟩
Select initial active virtual space
repeat
    Compute MP1 amplitudes t_ij^ab and MP2 energy J_2
    Assemble gradient G and Hessian H
    Solve Newton-Raphson equation: ΔR = −H^{−1} G
    Generate unitary matrix U = exp(R)
    Update MO coefficients: C ← C U
    Canonicalise active Fock block
    Evaluate new J_2
until |J_2^(n) − J_2^(n−1)| < ε_conv
```

4.2 Integral Transformation

4.2.1 AO-to-MO Transformation

4.2.2 Spin-Orbital Construction

4.2.3 Notation Convention: Chemists' vs. Physicists' Integrals

4.3 Initial Active Space Selection

4.4 MP1 Amplitudes and MP2 Energy

4.5 Gradient and Hessian Assembly

4.5.1 Block Structure and Efficient Inversion

4.5.2 Level-Shift Stabilisation

4.6 Orbital Rotation and Update

4.7 Convergence and Iteration Control

4.8 Initialisation Strategies

4.8.1 Default SCF Initialisation

4.8.2 Warm-Start from Previous Orbital Set

4.8.3 Random Unitary Sampling

Chapter 5

Computational Details

5.1 Software and Libraries

5.2 Hardware

5.3 Molecular Geometries

Table 5.1: Molecular geometries and electronic structure parameters used in this work.

Molecule	Geometry (Å)	Multiplicity	Charge	Symmetry
H_2	$r(\mathrm{H}-\mathrm{H}) = 0.74144$	1	0	$D_{\infty h}$
LiH	$r(\mathrm{Li}-\mathrm{H}) = 1.595$	1	0	$C_{\infty v}$
H_2O	$r = 0.957$, $\angle = 104.5$ř	1	0	C_{2v}
CH_2	planar, see text	3	0	C_{2v}
BH_3	$r(\mathrm{B}-\mathrm{H}) = 1.19$, planar	1	0	D_{3h}
N_2	$r(\mathrm{N}-\mathrm{N}) = 1.10$	1	0	$D_{\infty h}$
CO	$r(\mathrm{C}-\mathrm{O}) = 1.128$	1	0	$C_{\infty v}$
HF	$r(\mathrm{H}-\mathrm{F}) = 0.917$	1	0	$C_{\infty v}$
NH_3	$r(\mathrm{N}-\mathrm{H}) = 1.012$	1	0	C_{3v}

5.4 Basis Sets

5.5 Reference SCF Calculations

5.6 OVOS Calculation Settings

5.7 Data Storage and Reproducibility

Chapter 6

Results

In this chapter, the results of the OVOS implementation are presented. The chapter is structured as follows:

- **Section 6.1** validates the implementation by comparing full-space MP2 correlation energies against PySCF and literature values.
- **Section 6.2** presents the main result: the fraction of MP2 correlation energy recovered as a function of the active virtual orbital count.
- **Section 6.3** analyses the convergence behaviour of the OVOS algorithm.
- **Section 6.4** compares different initialisation strategies and their impact on final energies and convergence.
- **Section 6.5** investigates how the choice of basis set affects energy recovery.
- **Section 6.6** provides a direct comparison with the results published by Adamowicz and Bartlett in their original OVOS paper.
- **Section 6.7** assesses the computational cost and scaling of the OVOS algorithm.
- **Section 6.8** examines the T1 amplitudes in the converged OVOS orbitals to test for Brueckner-like properties.
- **Section 6.9** estimates the practical benefits of using OVOS orbitals as a reference state for quantum computing applications, including overlap with FCI states and resource estimates for VQE.

6.1 Validation of the Implementation

6.2 Energy Recovery as a Function of Active Virtual Orbital Count

6.3 Convergence Behaviour

6.4 Effect of Initialisation Strategy

6.5 Basis Set Dependence

6.6 Comparison with Adamowicz and Bartlett (1987)

6.7 Computational Cost

6.8 T1 Amplitudes and the Brueckner Orbital Connection

6.9 OVOS Orbitals as a VQE Reference State: Resource Estimates

Chapter 7

Conclusion

7.1 Summary of Findings

7.2 Critical Assessment

7.3 Implications for Quantum Computing

Chapter 8

Outlook

8.1 Algorithmic Improvements

8.2 Method Extensions

8.3 Quantum Computing Applications

8.3.1 OVOS Orbitals as a VQE Reference State

8.3.2 Brueckner Orbitals and Circuit Depth on NISQ Devices

8.3.3 Qubit Count Reduction and Fault-Tolerant Perspectives

8.4 Larger Molecular Systems

8.5 Benchmarking and Validation

Bibliography

Appendix A

Mathematical Derivations

A.1 Spin-Orbital MP2 Energy from the Hylleraas Functional

A.2 Derivation of the Orbital Rotation Gradient

A.3 Derivation of the Orbital Rotation Hessian

A.4 Matrix Exponential for the Rotation Matrix

A.5 Notation Conventions for Two-Electron Integrals

Table A.1: Summary of two-electron integral notation used in this thesis.

Notation	Definition	Used where
$(pq\|rs)$	$\int\int \phi_p^*(1)\phi_q(1)r_{12}^{-1}\phi_r^*(2)\phi_s(2)\,d1\,d2$	PySCF output
$\langle pr\|qs\rangle$	same as $(pq\|rs)$	Physicists' (non-antisym.)
$\langle pq\|\|rs\rangle$	$(pr\|qs) - (ps\|qr)$	Antisymmetrised; used in all formulas

Appendix B

Additional Data and Figures

B.1 Full MP2 Reference Energies

B.2 Complete OVOS Energy Recovery Tables

B.3 Convergence Plots for All Systems

B.4 Hessian Eigenvalue Analysis

Appendix C

Implementation Details and Source Code

C.1 Repository and Version Information

C.2 Spin-Orbital Integral Conversion

Listing C.1: Spatial-to-spin ERI conversion (`spatial_2_spin_eri_optimized`).

```
# Convert spatial ERIs to spin-orbital basis (interleaved alpha
    /beta)
def spatial_2_spin_eri_optimized(self, eri_aaaa, eri_aabb,
    eri_bbbb):
    n_spin = self.tot_num_spin_orbs
    eri_spin = np.zeros((n_spin, n_spin, n_spin, n_spin))
    # alpha-alpha block (even indices)
    eri_spin[0::2, 0::2, 0::2, 0::2] = eri_aaaa
    # beta-beta block (odd indices)
    eri_spin[1::2, 1::2, 1::2, 1::2] = eri_bbbb
    # alpha-alpha / beta-beta cross block
    eri_spin[0::2, 0::2, 1::2, 1::2] = eri_aabb
    eri_spin[1::2, 1::2, 0::2, 0::2] = eri_aabb.transpose(2, 3,
        0, 1)
    return eri_spin
```

C.3 MP2 Energy and Amplitude Computation

C.4 Orbital Optimisation: Gradient and Hessian

C.5 Profiling Results