




$$\sin \theta = \frac{\text{OPP}}{\text{HYP}}$$

$$\csc \theta = \frac{\text{HYP}}{\text{OPP}}$$

$$\cos \theta = \frac{\text{ADJ}}{\text{HYP}}$$

$$\sec \theta = \frac{\text{HYP}}{\text{ADJ}}$$

$$\tan \theta = \frac{\text{OPP}}{\text{ADJ}}$$

$$\cot \theta = \frac{\text{ADJ}}{\text{OPP}}$$

Opposite Angle Identities

$$\begin{aligned} \sin(-\theta) &= -\sin \theta & \cos(-\theta) &= \cos \theta & \tan(-\theta) &= -\tan \theta \\ \csc(-\theta) &= -\csc \theta & \sec(-\theta) &= \sec \theta & \cot(-\theta) &= -\cot \theta \end{aligned}$$

Complete Rotation and Half Rotation Identities

$$\begin{aligned} \sin(\theta \pm 360^\circ) &= \sin \theta & \sin(\theta \pm 180^\circ) &= -\sin \theta \\ \cos(\theta \pm 360^\circ) &= \cos \theta & \cos(\theta \pm 180^\circ) &= -\cos \theta \\ \tan(\theta \pm 360^\circ) &= \tan \theta & \csc(\theta \pm 180^\circ) &= -\csc \theta \\ \csc(\theta \pm 360^\circ) &= \csc \theta & \sec(\theta \pm 180^\circ) &= -\sec \theta \\ \sec(\theta \pm 360^\circ) &= \sec \theta & \tan(\theta \pm 180^\circ) &= \tan \theta \\ \cot(\theta \pm 360^\circ) &= \cot \theta & \cot(\theta \pm 180^\circ) &= \cot \theta \end{aligned}$$

Complete Rotation Angle Pair Identities and Supplementary Angle Identities

$$\begin{aligned} \sin(360^\circ - \theta) &= -\sin \theta & \sin(180^\circ - \theta) &= \sin \theta \\ \cos(360^\circ - \theta) &= \cos \theta & \csc(180^\circ - \theta) &= \csc \theta \\ \tan(360^\circ - \theta) &= -\tan \theta & \cos(180^\circ - \theta) &= -\cos \theta \\ \csc(360^\circ - \theta) &= -\csc \theta & \sec(180^\circ - \theta) &= -\sec \theta \\ \sec(360^\circ - \theta) &= \sec \theta & \tan(180^\circ - \theta) &= -\tan \theta \\ \cot(360^\circ - \theta) &= -\cot \theta & \cot(180^\circ - \theta) &= -\cot \theta \end{aligned}$$

Sum to Product Formulas

$$\begin{aligned} \sin A + \sin B &= 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right) \\ \sin A - \sin B &= 2 \sin\left(\frac{A-B}{2}\right) \cos\left(\frac{A+B}{2}\right) \\ \cos A + \cos B &= 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right) \\ \cos A - \cos B &= -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right) \end{aligned}$$

Product to Sum Formulas

$$\begin{aligned} \sin A \cos B &= \frac{1}{2} [\sin(A+B) + \sin(A-B)] \\ \cos A \sin B &= \frac{1}{2} [\sin(A+B) - \sin(A-B)] \\ \cos A \cos B &= \frac{1}{2} [\cos(A+B) + \cos(A-B)] \\ \sin A \sin B &= \frac{1}{2} [\cos(A-B) - \cos(A+B)] \end{aligned}$$

The Area of an SAS Triangle

The area of a triangle is equal to the length of one side times the length of another side times the sine of the measure of the angle between the two sides divided by two.

$$\text{Area} = \frac{1}{2} ab \sin C \quad \text{Area} = \frac{1}{2} bc \sin A$$

$$\text{Area} = \frac{1}{2} ac \sin B$$

30°-60°-90° Triangle Theorem: If the inner angles of a triangle are 30°, 60° and 90°, and the length of the side opposite the 30° angle is w , then the length of the hypotenuse is $2w$ and the length of the side opposite the 60° angle is $w\sqrt{3}$.



45°-45°-90° Triangle Theorem: If the inner angles of a triangle are 45°, 45° and 90°, and the length of the two sides opposite the 45° angles are w , then the length of the hypotenuse is $w\sqrt{2}$.



Quarter Rotation Identities

$$\begin{aligned} \sin(\theta \pm 90^\circ) &= \pm \cos \theta & \cos(\theta \pm 90^\circ) &= \mp \sin \theta \\ \csc(\theta \pm 90^\circ) &= \pm \sec \theta & \sec(\theta \pm 90^\circ) &= \mp \csc \theta \\ \tan(\theta \pm 90^\circ) &= -\cot \theta & \cot(\theta \pm 90^\circ) &= -\tan \theta \end{aligned}$$

Complementary Angle Identities

$$\begin{aligned} \sin(90^\circ - \theta) &= \cos \theta & \csc(90^\circ - \theta) &= \sec \theta \\ \cos(90^\circ - \theta) &= \sin \theta & \sec(90^\circ - \theta) &= \csc \theta \\ \tan(90^\circ - \theta) &= \cot \theta & \cot(90^\circ - \theta) &= \tan \theta \end{aligned}$$

Scenarios when Solving Triangles

The Law of Sines

AAS ASA SSA

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

Ambiguity
i) No triangle
ii) One triangle
iii) Two triangles

The Law of Cosines

SSS SAS


$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$b^2 = c^2 + a^2 - 2ac \cos B$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Heron's Formula

If a , b , and c are the side lengths of a triangle, then the area of the triangle is equal to the expression below.



$$\text{Area} = \sqrt{s(s-a)(s-b)(s-c)}$$

$$\text{where } s = \frac{a+b+c}{2}$$

Reciprocal Identities

$$\begin{aligned} \sin \theta &= \frac{1}{\csc \theta} & \csc \theta &= \frac{1}{\sin \theta} \\ \cos \theta &= \frac{1}{\sec \theta} & \sec \theta &= \frac{1}{\cos \theta} \\ \tan \theta &= \frac{1}{\cot \theta} & \cot \theta &= \frac{1}{\tan \theta} \end{aligned}$$

Ratio Identities

$$\begin{aligned} \tan \theta &= \frac{\sin \theta}{\cos \theta} \\ \cot \theta &= \frac{\cos \theta}{\sin \theta} \end{aligned}$$

Pythagorean Identities

$$\begin{aligned} \sin^2 \theta + \cos^2 \theta &= 1 \\ \tan^2 \theta + 1 &= \sec^2 \theta & 1 + \cot^2 \theta &= \csc^2 \theta \end{aligned}$$

Sum and Difference Formulas

$$\sin(A \pm B) = \sin A \cos B \pm \sin B \cos A$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

Double Angle Formulas

$$\sin(2A) = 2 \sin A \cos A$$

$$\cos(2A) = \cos^2 A - \sin^2 A$$

$$\cos(2A) = 1 - 2 \sin^2 A$$

$$\cos(2A) = 2 \cos^2 A - 1$$

Half Angle Formulas

$$\sin\left(\frac{A}{2}\right) = \pm \sqrt{\frac{1 - \cos A}{2}}$$

$$\cos\left(\frac{A}{2}\right) = \pm \sqrt{\frac{1 + \cos A}{2}}$$

General Forms of Trigonometric Functions

$$\begin{aligned} y &= A \sin(\omega x - \phi) + B \\ y &= A \cos(\omega x - \phi) + B \end{aligned}$$


$$\begin{aligned} y &= A \csc(\omega x - \phi) + B & y &= A \tan(\omega x - \phi) + B \\ y &= A \sec(\omega x - \phi) + B & y &= A \cot(\omega x - \phi) + B \end{aligned}$$

Sinusoidal Function: A function whose graph has the same general form as the sine or cosine functions.


Midline: A horizontal line drawn through the middle of the graph of a trigonometric function to assist when graphing.

For the functions above, the equation of the midline is $y = B$.

Example:



Period: The number of degrees required for a trigonometric function to complete one whole cycle.



The period of the sine, cosine, cosecant, and secant functions is given by the following equation.

$$P = \frac{2\pi}{\omega}$$

The period of the tangent and cotangent functions is given by the following equation.

$$P = \frac{\pi}{\omega}$$

Phase Shift: The horizontal distance that a parent graph is translated (to the right or left).

The phase shift is given by the equation below.


$$PS = \frac{\phi}{\omega}$$

Amplitude: The vertical distance from the midline to the highest or lowest point on a sinusoidal graph.


The amplitude is the magnitude (i.e. the absolute value) of the coefficient of the trigonometric function.

Example: If $y = A \sin(\omega x - \phi) + B$, then Amplitude = $|A|$.


$y = \sin x$




$y = \cos x$




$y = \tan x$




$y = \csc x$



$y = \sec x$



$y = \cot x$




Arc Length

$$s = r\theta$$

The Area of a Sector

$$A = \frac{\theta}{2} r^2$$

Angle of Elevation and Angle of Depression



Converting from Degrees to Radians

$$\text{degree measure} \times \frac{\pi \text{ radians}}{180 \text{ degrees}} = \text{radian measure}$$

Converting from Radians to Degrees

$$\text{Radian measure} \times \frac{180 \text{ degrees}}{\pi \text{ radians}} = \text{degree measure}$$

If an object moves s units on a circular path in t units of time, then v is defined as the linear velocity according to the equation below.

$$v = \frac{s}{t}$$

If an object on a circle rotates θ angle units in t units of time, then ω is defined as the angular velocity according to the equation below.

$$\omega = \frac{\theta}{t}$$

$$v = r\omega$$

Revolution: A complete rotation.

Revolutions per minute = RPM

$$1 \text{ Revolution} = 2\pi \text{ radians} = 360^\circ$$

The Limit Laws

If both $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist, and " c " is a constant, then

$$1) \lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$$

$$2) \lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$$

$$3) \lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$$

$$4) \lim_{x \rightarrow a} \left[\frac{f(x)}{g(x)} \right] = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \quad \lim_{x \rightarrow a} g(x) \neq 0$$

$$5) \lim_{x \rightarrow a} [cf(x)] = c \lim_{x \rightarrow a} f(x)$$

$$6) \lim_{x \rightarrow a} c = c$$

$$7) \lim_{x \rightarrow a} f(g(x)) = f\left(\lim_{x \rightarrow a} g(x)\right) \text{ if "f" is continuous where its input is } \lim_{x \rightarrow a} g(x).$$

8) If " f " is continuous at $x=a$, then $\lim_{x \rightarrow a} f(x) = f(a)$.
*Remember, this is the definition of continuity at $x=a$.

Formulas for Calculating Derivatives (Part II)

Differentiation Formulas

Constant Rule i) $\frac{d}{dx}(c) = 0$ "c" is a constant.

Power Rule ii) $\frac{d}{dx}(x^n) = nx^{n-1}$ "n" is a constant.

Constant Multiple Rule iii) $\frac{d}{dx}[cf(x)] = cf'(x)$ "c" is a constant.

Sum Rule iv) $\frac{d}{dx}[f(x) + g(x)] = f'(x) + g'(x)$

Difference Rule v) $\frac{d}{dx}[f(x) - g(x)] = f'(x) - g'(x)$

Product Rule vi) $\frac{d}{dx}[f(x)g(x)] = f'(x)g(x) + f(x)g'(x)$

Quotient Rule vii) $\frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$

Trigonometric Differentiation Formulas

I) $\frac{d}{dx}(\sin x) = \cos x$

II) $\frac{d}{dx}(\cos x) = -\sin x$

III) $\frac{d}{dx}(\csc x) = -\csc x \cot x$

IV) $\frac{d}{dx}(\sec x) = \sec x \tan x$

V) $\frac{d}{dx}(\tan x) = \sec^2 x$

VI) $\frac{d}{dx}(\cot x) = -\csc^2 x$

A Proof of Formula I

If $f(x) = \sin x$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin x \cos(h) + \sin(h) \cos x - \sin x}{h}$$

$\sin(A \pm B) = \sin A \cos B \pm \sin B \cos A$

$$= \lim_{h \rightarrow 0} \frac{\sin x \cos(h) - \sin x + \sin(h) \cos x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin x (\cos(h) - 1) + \sin(h) \cos x}{h}$$

$$= \lim_{h \rightarrow 0} \left[\frac{\sin x (\cos(h) - 1)}{h} + \frac{\sin(h) \cos x}{h} \right]$$

$$= \lim_{h \rightarrow 0} \left[\sin x \frac{(\cos(h) - 1)}{h} \right] + \lim_{h \rightarrow 0} \left[\frac{\sin(h)}{h} \cos x \right]$$

$$= \lim_{h \rightarrow 0} \sin x \lim_{h \rightarrow 0} \frac{\cos(h) - 1}{h} + \lim_{h \rightarrow 0} \frac{\sin(h)}{h} \lim_{h \rightarrow 0} \cos x$$

$$= \lim_{h \rightarrow 0} \sin x \lim_{h \rightarrow 0} \frac{\cos(h) - 1}{h} + \lim_{h \rightarrow 0} \frac{\sin(h)}{h} \lim_{h \rightarrow 0} \cos x$$

$$= \sin x (0) + (1) \cos x$$

$$= 0 + \cos x$$

$$= \cos x$$

Practice

Find $f'(x)$.

① $f(x) = x^3 + \sin x + 8$

$$f'(x) = 3x^2 + \cos x + 0$$

$$f'(x) = \boxed{3x^2 + \cos x}$$

② $f(x) = 3 \tan x - 8x^5$

$$f'(x) = \boxed{3 \sec^2 x - 40x^4}$$

$$\textcircled{3} f(x) = \cos x - 2 - \frac{\sec x}{5} - 10x$$

$$f(x) = \cos x - 2 - \frac{1}{5}\sec x - 10x$$

$$f'(x) = -\sin x + 0 - \frac{1}{5}\sec x \tan x - 10$$

$$f'(x) = \boxed{-\sin x - \frac{1}{5}\sec x \tan x - 10}$$

Find $\frac{dy}{dx}$.

$$\textcircled{4} y = 3x^7 - 32.1 \csc x + 64.87 - \cot x$$

$$\frac{dy}{dx} = 21x^6 - 32.1(-\csc x \cot x) + 0 - (-\csc^2 x)$$

$$\frac{dy}{dx} = \boxed{21x^6 + 32.1 \csc x \cot x + \csc^2 x}$$

• $\textcircled{5} y = 4 \cos x + \sin x + 6$

$$\frac{dy}{dx} = 4(-\sin x) + \cos x + 0$$

$$\frac{dy}{dx} = \boxed{-4 \sin x + \cos x}$$

$$\textcircled{6} y = \tan x - x^6$$

$$\frac{dy}{dx} = \boxed{\sec^2 x - 6x^5}$$

Find y' .

$$\textcircled{7} y = \frac{\cot x}{2} - 10 \csc x - 15x^3 + 18$$

$$y = \frac{1}{2} \cot x - 10 \csc x - 15x^3 + 18$$

$$y' = \frac{1}{2}(-\csc^2 x) - 10(-\csc x \cot x) - 45x^2 + 0$$

$$y' = \boxed{-\frac{1}{2}\csc^2 x + 10 \csc x \cot x - 45x^2}$$

$$\textcircled{8} y = -9x + \sec x$$

$$y' = \boxed{-9 + \sec x \tan x}$$

A Proof of the Product Rule

If $r(x) = f(x)g(x)$, then

$$r'(x) = \lim_{h \rightarrow 0} \frac{r(x+h) - r(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - \cancel{f(x+h)g(x)} + \cancel{f(x+h)g(x)} - f(x)g(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f(x+h)[g(x+h) - g(x)] + g(x)[f(x+h) - f(x)]}{h}$$

$$= \lim_{h \rightarrow 0} \left\{ \frac{f(x+h)[g(x+h) - g(x)]}{h} + \frac{g(x)[f(x+h) - f(x)]}{h} \right\}$$

$$= \lim_{h \rightarrow 0} \left\{ f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right\}$$

$$\stackrel{1}{=} \lim_{h \rightarrow 0} \left[f(x+h) \frac{g(x+h) - g(x)}{h} \right] + \lim_{h \rightarrow 0} \left[g(x) \frac{f(x+h) - f(x)}{h} \right]$$

$$\stackrel{3}{=} \lim_{h \rightarrow 0} f(x+h) \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} + \lim_{h \rightarrow 0} g(x) \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$\stackrel{886}{=} f(x+0) g'(x) + g(x) f'(x)$$

$$= f(x)g'(x) + f'(x)g(x)$$

Find $\frac{dy}{dx}$.

• ⑨ $y = x^5 \sin x$

$$\frac{dy}{dx} = \left[\frac{d}{dx} (x^5) \right] \sin x + x^5 \frac{d}{dx} (\sin x)$$

$$\frac{dy}{dx} = \boxed{5x^4 \sin x + x^5 \cos x}$$

⑩ $y = \cos x \cot x$

$$\frac{dy}{dx} = \left[\frac{d}{dx}(\cos x) \right] \cot x + \cos x \frac{d}{dx}(\cot x)$$

$$\frac{dy}{dx} = -\sin x \cot x + \cos x (-\csc^2 x)$$

$$\frac{dy}{dx} = -\sin x \frac{\cos x}{\sin x} - \cos x \csc^2 x$$

$$\frac{dy}{dx} = -\cos x - \cos x \csc^2 x$$

⑪ $y = x^2 \csc x$

$$\frac{dy}{dx} = \left[\frac{d}{dx}(x^2) \right] \csc x + x^2 \frac{d}{dx}(\csc x)$$

$$\frac{dy}{dx} = 2x \csc x + x^2 (-\csc x \cot x)$$

$$\frac{dy}{dx} = 2x \csc x - x^2 \csc x \cot x$$

Find y'

⑫ $y = \sec x \tan x$

$$y' = \left[\frac{d}{dx}(\sec x) \right] \tan x + \sec x \frac{d}{dx}(\tan x)$$

$$y' = \sec x \tan x \tan x + \sec x \sec^2 x$$

$$y' = \sec x \tan^2 x + \sec^3 x$$

• ⑬ $y = x^4 \tan x$ $\frac{d}{dx}[f(x)g(x)] = f'(x)g(x) + f(x)g'(x)$

$$y' = \left[\frac{d}{dx}(x^4) \right] \tan x + x^4 \frac{d}{dx}(\tan x)$$

$$y' = 4x^3 \tan x + x^4 \sec^2 x$$

$$\textcircled{14} y = \sin x \cos x$$

$$y' = \left[\frac{d}{dx}(\sin x) \right] \cos x + \sin x \frac{d}{dx}(\cos x)$$

$$y' = \cos x \cos x + \sin x (-\sin x)$$

$$y' = \cos^2 x - \sin^2 x$$

$$y' = \boxed{\cos(2x)}$$

Find $\frac{dy}{dx}$.

$$\textcircled{15} y = x^3 \sec x$$

$$\frac{dy}{dx} = \left[\frac{d}{dx}(x^3) \right] \sec x + x^3 \frac{d}{dx}(\sec x)$$

$$\frac{dy}{dx} = \boxed{3x^2 \sec x + x^3 \sec x \tan x}$$

$$\textcircled{16} y = \cot x \csc x$$

$$\frac{dy}{dx} = \left[\frac{d}{dx}(\cot x) \right] \csc x + \cot x \frac{d}{dx}(\csc x)$$

$$\frac{dy}{dx} = (-\csc^2 x) \csc x + \cot x (-\csc x \cot x)$$

$$\frac{dy}{dx} = \boxed{-\csc^3 x - \csc x \cot^2 x}$$

• $\textcircled{17} y = x \cot x + x^6$

$$\frac{dy}{dx} = \left[\frac{d}{dx}(x) \right] \cot x + x \frac{d}{dx}(\cot x) + 6x^5$$

$$\frac{dy}{dx} = (1) \cot x + x (-\csc^2 x) + 6x^5$$

$$\frac{dy}{dx} = \boxed{\cot x - x \csc^2 x + 6x^5}$$

Find y' .

$$\textcircled{18} y = -3x^2 - x^2 \sec x$$

$$y' = -6x - \frac{d}{dx}(x^2 \sec x)$$

$$y' = -6x - \left(\frac{d}{dx}(x^2) \sec x + x^2 \frac{d}{dx}(\sec x) \right)$$

$$y' = -6x - (2x \sec x + x^2 \sec x \tan x)$$

$$y' = \boxed{-6x - 2x \sec x - x^2 \sec x \tan x}$$

$$\bullet \textcircled{19} y = x^{10} \csc x + 5x^8$$

$$y' = \left[\frac{d}{dx}(x^{10}) \right] \csc x + x^{10} \frac{d}{dx}(\csc x) + 40x^7$$

$$y' = 10x^9 \csc x + x^{10}(-\csc x \cot x) + 40x^7$$

$$y' = \boxed{10x^9 \csc x - x^{10} \csc x \cot x + 40x^7}$$

$$\textcircled{20} y = x^7 - x \tan x$$

$$y' = 7x^6 - \frac{d}{dx}(x \tan x)$$

$$y' = 7x^6 - \left(\left[\frac{d}{dx}(x) \right] \tan x + x \frac{d}{dx}(\tan x) \right)$$

$$y' = 7x^6 - (1 \tan x + x \sec^2 x)$$

$$y' = \boxed{7x^6 - \tan x - x \sec^2 x}$$

Find $\frac{dy}{dx}$.

$$\bullet \textcircled{21} y = 3 \sin x - x^2 \cos x$$

$$\frac{dy}{dx} = 3 \cos x - \frac{d}{dx}(x^2 \cos x)$$

$$\frac{dy}{dx} = 3 \cos x - \left(\left[\frac{d}{dx}(x^2) \right] \cos x + x^2 \frac{d}{dx}(\cos x) \right)$$

$$\frac{dy}{dx} = 3 \cos x - (2x \cos x + x^2(-\sin x))$$

$$\frac{dy}{dx} = 3 \cos x - (2x \cos x - x^2 \sin x)$$

$$\frac{dy}{dx} = \boxed{3 \cos x - 2x \cos x + x^2 \sin x}$$

• (22) $y = 2\tan x - x^4 \cot x$

$$\frac{dy}{dx} = 2\sec^2 x - \frac{d}{dx}(x^4 \cot x)$$

$$\frac{dy}{dx} = 2\sec^2 x - \left(\left[\frac{d}{dx} x^4 \right] \cot x + x^4 \frac{d}{dx} (\cot x) \right)$$

$$\frac{dy}{dx} = 2\sec^2 x - (4x^3 \cot x + x^4 (-\csc^2 x))$$

$$\frac{dy}{dx} = 2\sec^2 x - (4x^3 \cot x - x^4 \csc^2 x)$$

$$\frac{dy}{dx} = \boxed{2\sec^2 x - 4x^3 \cot x + x^4 \csc^2 x}$$

The Quotient Rule

Find y' .

• (23) $y = \frac{\sin x}{x^3}$

$$y' = \frac{\left[\frac{d}{dx} (\sin x) \right] x^3 - (\sin x) \frac{d}{dx} (x^3)}{(x^3)^2}$$

$$y' = \frac{(\cos x) x^3 - (\sin x) (3x^2)}{x^6}$$

$$y' = \frac{x^2 (x \cos x - 3 \sin x)}{x^2 x^4}$$

$$y' = \boxed{\frac{x \cos x - 3 \sin x}{x^4}}$$

$$\textcircled{24} y = \frac{9x}{5x^2+2}$$

$$y' = \frac{\left[\frac{d}{dx}(9x) \right] (5x^2+2) - (9x) \frac{d}{dx}(5x^2+2)}{(5x^2+2)^2}$$

$$y' = \frac{9(5x^2+2) - 9x(10x+0)}{(5x^2+2)^2}$$

$$y' = \frac{45x^2+18-90x^2}{(5x^2+2)^2}$$

$$y' = \boxed{\frac{-45x^2+18}{(5x^2+2)^2}}$$

$$\bullet \textcircled{25} y = \frac{\sec x}{2x^2}$$

$$y' = \frac{\left[\frac{d}{dx}(\sec x) \right] (2x^2) - (\sec x) \frac{d}{dx}(2x^2)}{(2x^2)^2}$$

$$y' = \frac{(\sec x \tan x)(2x^2) - (\sec x)(4x)}{4x^4}$$

$$y' = \frac{2x(x \sec x \tan x - 2 \sec x)}{2x \cdot 2x^3}$$

$$y' = \boxed{\frac{x \sec x \tan x - 2 \sec x}{2x^3}}$$

$$\textcircled{26} \ y = \frac{x^3}{x^4-6}$$

$$y' = \frac{\left[\frac{d}{dx}(x^3) \right] (x^4-6) - x^3 \frac{d}{dx}(x^4-6)}{(x^4-6)^2}$$

$$y' = \frac{3x^2(x^4-6) - x^3(4x^3+0)}{(x^4-6)^2}$$

$$y' = \frac{3x^6 - 18x^2 - 4x^6}{(x^4-6)^2}$$

$$y' = \boxed{\frac{-x^6 - 18x^2}{(x^4-6)^2}}$$

Find $\frac{dy}{dx}$.

$$\bullet \textcircled{27} \ y = \frac{3x-2}{4x^3+5}$$

$$\frac{dy}{dx} = \frac{\left[\frac{d}{dx}(3x-2) \right] (4x^3+5) - (3x-2) \frac{d}{dx}(4x^3+5)}{(4x^3+5)^2}$$

$$\frac{dy}{dx} = \frac{(3+0)(4x^3+5) - (3x-2)(12x^2+0)}{(4x^3+5)^2}$$

$$\frac{dy}{dx} = \frac{12x^3+15 - (36x^3-24x^2)}{(4x^3+5)^2}$$

$$\frac{dy}{dx} = \frac{12x^3+15 - 36x^3+24x^2}{(4x^3+5)^2}$$

$$\frac{dy}{dx} = \boxed{\frac{-24x^3+24x^2+15}{(4x^3+5)^2}}$$

• ②⑧ $y = \frac{x+1}{9x^2-5}$

$$\frac{dy}{dx} = \frac{\left[\frac{d}{dx}(x+1) \right] (9x^2-5) - (x+1) \frac{d}{dx}(9x^2-5)}{(9x^2-5)^2}$$

$$\frac{dy}{dx} = \frac{(1+0)(9x^2-5) - (x+1)(18x+0)}{(9x^2-5)^2}$$

$$\frac{dy}{dx} = \frac{9x^2-5 - (18x^2+18x)}{(9x^2-5)^2}$$

$$\frac{dy}{dx} = \frac{9x^2-5-18x^2-18x}{(9x^2-5)^2}$$

$$\frac{dy}{dx} = \frac{-9x^2-18x-5}{(9x^2-5)^2}$$

• ②⑨ $y = \frac{2x^5+6}{-4x^6+2}$

$$\frac{dy}{dx} = \frac{\left[\frac{d}{dx}(2x^5+6) \right] (-4x^6+2) - (2x^5+6) \frac{d}{dx}(-4x^6+2)}{(-4x^6+2)^2}$$

$$\frac{dy}{dx} = \frac{(10x^4+0)(-4x^6+2) - (2x^5+6)(-24x^5+0)}{(-4x^6+2)^2}$$

$$\frac{dy}{dx} = \frac{-40x^{10}+20x^4 - (-48x^{10}-144x^5)}{(-4x^6+2)^2}$$

$$\frac{dy}{dx} = \frac{-40x^{10}+20x^4+48x^{10}+144x^5}{(-4x^6+2)^2}$$

$$\frac{dy}{dx} = \frac{8x^{10}+144x^5+20x^4}{(-4x^6+2)^2}$$

• ③⑩ $y = \frac{4x^2-1}{-6x^3-10}$

$$\frac{dy}{dx} = \frac{\left[\frac{d}{dx}(4x^2-1)\right](-6x^3-10) - (4x^2-1)\frac{d}{dx}(-6x^3-10)}{(-6x^3-10)^2}$$

$$\frac{dy}{dx} = \frac{(8x+0)(-6x^3-10) - (4x^2-1)(-18x^2+0)}{(-6x^3-10)^2}$$

$$\frac{dy}{dx} = \frac{-48x^4-80x - (-72x^4+18x^2)}{(-6x^3-10)^2}$$

$$\frac{dy}{dx} = \frac{-48x^4-80x+72x^4-18x^2}{(-6x^3-10)^2}$$

$$\frac{dy}{dx} = \boxed{\frac{24x^4-18x^2-80x}{(-6x^3-10)^2}}$$

Find y' .

• ③⑪ $y = \frac{2}{3\cot x+1}$

$$y' = \frac{\left[\frac{d}{dx}(2)\right](3\cot x+1) - 2\frac{d}{dx}(3\cot x+1)}{(3\cot x+1)^2}$$

$$y' = \frac{0(3\cot x+1) - 2(3(-\csc^2 x)+0)}{(3\cot x+1)^2}$$

$$y' = \frac{0 - 2(-3\csc^2 x)}{(3\cot x+1)^2}$$

$$y' = \boxed{\frac{6\csc^2 x}{(3\cot x+1)^2}}$$

• ③② $y = \frac{-1}{8\cos x - 10}$

$$y' = \frac{\left[\frac{d}{dx}(-1) \right] (8\cos x - 10) - (-1) \frac{d}{dx}(8\cos x - 10)}{(8\cos x - 10)^2}$$

$$y' = \frac{0(8\cos x - 10) - (-1)(8(-\sin x) + 0)}{(8\cos x - 10)^2}$$

$$y' = \frac{0 - (-1)(-8\sin x)}{(8\cos x - 10)^2}$$

$$y' = \boxed{\frac{-8\sin x}{(8\cos x - 10)^2}}$$

Examples where We Do Not Need the Quotient Rule

• ③③ $f(x) = \frac{x^3}{4}$

$$f(x) = \frac{1}{4}x^3$$

$$f'(x) = \boxed{\frac{3}{4}x^2}$$

• ③④ $f(x) = \frac{\csc x}{\sqrt{10}}$

$$f(x) = \frac{1}{\sqrt{10}} \csc x$$

$$f'(x) = \frac{1}{\sqrt{10}} (-\csc x \cot x) = \boxed{-\frac{1}{\sqrt{10}} \csc x \cot x}$$

③⑤ $f(x) = \frac{9x^2 + 4x - 2}{3}$

$$f(x) = \frac{1}{3}(9x^2 + 4x - 2)$$

$$f'(x) = \frac{1}{3}(18x + 4) = \boxed{6x + \frac{4}{3}}$$

$$f(x) = 3x^2 + \frac{4}{3}x - \frac{2}{3}$$

$$f'(x) = \boxed{6x + \frac{4}{3}}$$