



### Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Age * Follow_MLS_or_not	132	100.0%	0	0.0%	132	100.0%
Gender * Follow_MLS_or_not	131	99.2%	1	0.8%	132	100.0%
Annual_Household_Income * Follow_MLS_or_not	132	100.0%	0	0.0%	132	100.0%

### Age \* Follow\_MLS\_or\_not

#### Crosstab



Count

		Follow_MLS_or_not		Total
		0	1	
Age	1	1	4	5
	2	19	28	47
	3	24	18	42
	4	14	14	28
	5	4	2	6
	6	4	0	4
Total		66	66	132

#### Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	9.047 <sup>a</sup>	5	.107
Likelihood Ratio	10.746	5	.057
Linear-by-Linear Association	6.161	1	.013
N of Valid Cases	132		

a. 6 cells (50.0%) have expected count less than 5. The minimum expected count is 2.00.



### Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	.079 <sup>a</sup>	1	.778		
Continuity Correction <sup>b</sup>	.011	1	.917		
Likelihood Ratio	.079	1	.778		
Fisher's Exact Test				.860	.459
Linear-by-Linear Association	.079	1	.779		
N of Valid Cases	131				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 26.79.

b. Computed only for a 2x2 table



### Symmetric Measures

	Value	Asymptotic Standard Error <sup>a</sup>	Approximate T <sup>b</sup>	Approximate Significance
Interval by Interval Pearson's R	-.025	.087	-.280	.780 <sup>c</sup>
Ordinal by Ordinal Spearman Correlation	-.025	.087	-.280	.780 <sup>c</sup>
N of Valid Cases	131			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.



### Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)
Pearson Chi-Square	4.930 <sup>a</sup>	5	.425
Likelihood Ratio	4.999	5	.416
Linear-by-Linear Association	1.277	1	.258
N of Valid Cases	132		

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 7.50.

### Symmetric Measures

	Value	Asymptotic Standard Error <sup>a</sup>	Approximate T <sup>b</sup>	Approximate Significance
Interval by Interval Pearson's R	.099	.086	1.131	.260 <sup>c</sup>
Ordinal by Ordinal Spearman Correlation	.093	.087	1.065	.289 <sup>c</sup>
N of Valid Cases	132			

- a. Not assuming the null hypothesis.
- b. Using the asymptotic standard error assuming the null hypothesis.
- c. Based on normal approximation.

