

THEOREM 1

Let A , B , and C be matrices of the same size, and let r and s be scalars.

a. $A + B = B + A$

b. $(A + B) + C = A + (B + C)$

c. $A + 0 = A$

d. $r(A + B) = rA + rB$

e. $(r + s)A = rA + sA$

f. $r(sA) = (rs)A$

Matrix Multiplication

When a matrix B multiplies a vector \mathbf{x} , it transforms \mathbf{x} into the vector $B\mathbf{x}$. If this vector is then multiplied in turn by a matrix A , the resulting vector is $A(B\mathbf{x})$. See Fig. 2.





FIGURE 2 Multiplication by B and then A .

Thus $A(B\mathbf{x})$ is produced from \mathbf{x} by a *composition* of mappings—the linear transformations studied in Section 1.8. Our goal is to represent this composite mapping as multiplication by a single matrix, denoted by AB , so that

$$A(B\mathbf{x}) = (AB)\mathbf{x} \quad (1)$$

See Fig. 3.



If A is an $m \times n$ matrix, and if B is an $n \times p$ matrix with columns $\mathbf{b}_1, \dots, \mathbf{b}_p$, then the product AB is the $m \times p$ matrix whose columns are $A\mathbf{b}_1, \dots, A\mathbf{b}_p$. That is,

$$AB = A[\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_p] = [A\mathbf{b}_1 \quad A\mathbf{b}_2 \quad \cdots \quad A\mathbf{b}_p]$$

multiplication of matrices corresponds to composition of linear transformations

each column of AB is a linear combination of the columns of A using weights from the corresponding column of B .

Properties of Matrix Multiplication

The following theorem lists the standard properties of matrix multiplication. Recall that I_m represents the $m \times m$ identity matrix and $I_m \mathbf{x} = \mathbf{x}$ for all \mathbf{x} in \mathbb{R}^m .

THEOREM 2

Let A be an $m \times n$ matrix, and let B and C have sizes for which the indicated sums and products are defined.

- a. $A(BC) = (AB)C$ (associative law of multiplication)
 - b. $A(B + C) = AB + AC$ (left distributive law)
 - c. $(B + C)A = BA + CA$ (right distributive law)
 - d. $r(AB) = (rA)B = A(rB)$
for any scalar r
 - e. $I_m A = A = A I_n$ (identity for matrix multiplication)
-

WARNINGS:

1. In general, $AB \neq BA$.
2. The cancellation laws do *not* hold for matrix multiplication. That is, if $AB = AC$, then it is *not* true in general that $B = C$. (See Exercise 10.)
3. If a product AB is the zero matrix, you *cannot* conclude in general that either $A = 0$ or $B = 0$. (See Exercise 12.)

THEOREM 3

Let A and B denote matrices whose sizes are appropriate for the following sums and products.

- a. $(A^T)^T = A$
 - b. $(A + B)^T = A^T + B^T$
 - c. For any scalar r , $(rA)^T = rA^T$
 - d. $(AB)^T = B^T A^T$
-