

1. a) Given,

$$H = - \left(\frac{w_{12} + w_{21}}{2} \right) S_1 S_2$$

$$\therefore w_{12} = 2 \text{ and } w_{21} = -1$$

$$\Rightarrow H = -0.5 S_1 S_2$$

When we update S_2 asynchronously,

$$H' = -0.5 S_1 S_2'$$

$$\begin{aligned} \therefore \Delta H &= H' - H = -0.5 S_1 (S_2' - S_2) \\ &= -0.5 S_1 (\operatorname{sgn}(-S_1) - S_2) \\ &= -0.5 S_1 (-S_1 - S_2) \\ &= 0.5 S_1 (S_1 + S_2) = 0.5 S_1^2 + 0.5 S_1 S_2 \\ &= \cancel{0.5} \cancel{S_1^2} = 0.5 (1 + S_1 S_2) \because (S_1^2 = 1) \\ &\geq 0 \quad \checkmark \quad (\because S_1 S_2 = \pm 1) \end{aligned}$$

\Rightarrow Energy can increase. \checkmark

When we update S_1 asynchronously,

$$H' = -0.5 S_1' S_2$$

$$\begin{aligned} \therefore \Delta H &= -0.5 S_2 (S_1' - S_1) \\ &= -0.5 S_2 (\operatorname{sgn}(2S_2) - S_1) \\ &= -0.5 S_2 (S_2 - S_1) \\ &= 0.5 (S_1 S_2 - S_2^2) \\ &= 0.5 (S_1 S_2 - 1) \leq 0 \quad \checkmark \end{aligned}$$

$$\therefore \Delta H \leq 0$$

Q, S

\Rightarrow Energy cannot increase. \checkmark

(b)

$$H = - \left(\frac{w_{12} + w_{21}}{2} \right) S_1 S_2$$

$$= -0.5 S_1 S_2$$

After update, $H' = -0.5 S_1' S_2'$

$$\therefore \Delta H = -0.5 (S_1' S_2' - S_1 S_2) \quad \checkmark$$

$$\begin{aligned}\Delta H &= -0.5 \left[\operatorname{sgn}(2s_2) \operatorname{sgn}(-s_1) - s_1 s_2 \right] \\ &= -0.5 \left[-s_2 s_1 - s_1 s_2 \right] \\ &= s_1 s_2 \quad \checkmark\end{aligned}$$

$\therefore s_1$ and s_2 can take values ± 1 only.

So, $s_1 s_2$ can be ± 1 only.

So, $\Delta H \neq 0$

0.5

\Rightarrow Energy can't stay constant after synchronous update. \checkmark

(c)

$$H = -\frac{1}{2} \sum_{i,j}^N w_{ij} n_i n_j + \sum_{i=1}^N \mu_i n_i$$

Update m^{th} neuron:

$$n'_m = \Theta_H(b_m), \quad b_m = \sum_{j=1}^N w_{mj} n_j - \mu_m$$

$$\begin{aligned}\therefore \hat{H} &= -\frac{1}{2} \sum_{i=1}^N \sum_{j=1}^N w_{ij} n_i n_j - \frac{1}{2} \sum_{j=1}^N w_{mj} n'_m n_j - \frac{1}{2} \sum_{i=1}^N w_{im} n_i n'_m \\ &\quad - \frac{1}{2} w_{mm} n'_m n'_m + \sum_{i=1, i \neq m}^N \mu_i n_i + \mu_m n'_m\end{aligned}$$

$$\therefore \Delta H = \hat{H} - H$$

$$= -\frac{1}{2} \sum_{i=1, i \neq m}^N (w_{mi} + w_{im}) (n_i n'_m - n_i n_m) - \frac{1}{2} w_{mm} (n'_m{}^2 - n_m{}^2) + \mu_m (n'_m - n_m)$$

$\left[\because \text{First term } \hat{H} \text{ gets cancelled as it remains same as in } H \right]$

0.5

Now $\because w_{ij} = w_{ji}$

$$\Rightarrow \Delta H = -\sum_{i=1, i \neq m}^N w_{im} n_i n'_m - \frac{1}{2} w_{mm} (n'_m{}^2 - n_m{}^2) + \mu_m (n'_m - n_m)$$

$$\begin{aligned}&= -n'_m \left[\sum_{i=1}^N w_{im} n_i - \mu_m \right] + w_{mm} n_m n'_m - \frac{1}{2} w_{mm} (n'_m{}^2 - n_m{}^2) \\ &= -b_m n'_m - \underline{w_{mm} (n'_m{}^2 - n_m n'_m + 2 n_m n'_m)} - \mu_m n'_m \end{aligned}$$

$\rightarrow \mu_m n'_m$

$$\text{If } n_m' = n_m \Rightarrow \Delta H = 0 \quad \checkmark$$

Now if $n_m = 0$ and $n_m' = 1 \Rightarrow b_m > 0$

$$\begin{aligned} \Rightarrow \Delta H &= -b_m - \frac{w_{mm}}{2} (1-0+0) - \mu_m(0) \\ &= -b_m - \frac{w_{mm}}{2} \quad \checkmark \end{aligned}$$

$\therefore b_m > 0$ and $w_{mm} > 0 \quad \checkmark$

$$\Rightarrow \Delta H < 0 \quad \checkmark$$

If $n_m = 1$ and $n_m' = 0 \Rightarrow b_m < 0$

$$\begin{aligned} \Rightarrow \Delta H &= -\frac{w_{mm}}{2} (-1) - \mu_m \\ &= \frac{w_{mm}}{2} - \mu_m < 0 \quad (\because b_m < 0) \end{aligned}$$

$$\therefore \Delta H \leq 0$$

$\Rightarrow H$ cannot increase under this dynamics. $\checkmark \quad 0.5$

2/2

2. (a) Let $H_{\mu\nu}$ denote Hamming distance between pattern μ and ν .

$$\text{Let } Q_{\mu\nu} = \sum_{j=1}^N x_j^\mu x_j^\nu$$

$$\text{So, } Q_{\mu\nu} = N - 2H_{\mu\nu} = 35 - 2H_{\mu\nu}$$

$$\therefore \text{for } \mu=1, H_{11} = 0 \Rightarrow Q_{11} = 35 \checkmark$$

$$H_{12} = 10 \Rightarrow Q_{12} = 15 \checkmark$$

$$H_{13} = 8 \Rightarrow Q_{13} = 19 \checkmark$$

$$H_{14} = 11 \Rightarrow Q_{14} = 13 \checkmark$$

$$H_{15} = 35 \Rightarrow Q_{15} = -35 \checkmark$$

$$\text{For } \mu=2, H_{21} = 10 \Rightarrow Q_{21} = 15 \checkmark$$

$$H_{22} = 0 \Rightarrow Q_{22} = 35 \checkmark$$

$$H_{23} = 14 \Rightarrow Q_{23} = 7 \checkmark$$

$$H_{24} = 3 \Rightarrow Q_{24} = 29 \checkmark$$

$$H_{25} = 25 \Rightarrow Q_{25} = -15 \checkmark$$

$$\begin{aligned}
 (b) b_i^{(v)} &= \sum_{j=1}^N w_{ij} x_j^{(v)} \\
 &= \sum_{j=1}^N \left(\frac{1}{35} \sum_{\mu=1}^2 x_i^{(\mu)} x_j^{(\mu)} \right) x_j^{(v)} \\
 &= \frac{1}{35} \left[\sum_{j=1}^N x_i^{(1)} x_j^{(1)} x_j^{(v)} + \sum_{j=1}^N x_i^{(2)} x_j^{(2)} x_j^{(v)} \right] \\
 &= \frac{1}{35} \left[x_i^{(1)} Q_{1v} + x_i^{(2)} Q_{2v} \right]
 \end{aligned}$$

$$\therefore b_i^{(1)} = \frac{1}{35} \left[35 x_i^{(1)} + 15 x_i^{(2)} \right] = x_i^{(1)} + \frac{3}{7} x_i^{(2)} \checkmark$$

$$b_i^{(2)} = \frac{1}{35} \left[15 x_i^{(1)} + 35 x_i^{(2)} \right] = \frac{3}{7} x_i^{(1)} + x_i^{(2)} \checkmark$$

$$b_i^{(3)} = \frac{1}{35} (19x_i^{(1)} + 7x_i^{(2)}) = \frac{19}{35} x_i^{(1)} + \frac{1}{5} x_i^{(2)} \quad \checkmark$$

$$b_i^{(4)} = \frac{1}{35} (13x_i^{(1)} + 29x_i^{(2)}) = \frac{13}{35} x_i^{(1)} + \frac{29}{35} x_i^{(2)} \quad \checkmark$$

$$b_i^{(5)} = \frac{1}{35} (-35x_i^{(1)} - 15x_i^{(2)}) = -x_i^{(1)} - \frac{3}{7} x_i^{(2)} \quad \checkmark$$

(c)

If $\operatorname{sgn}(b_i^{(v)}) = x_i^{(v)}$ then pattern remains same.

$$\Rightarrow v=1, \operatorname{sgn}(b_i^{(1)}) = \operatorname{sgn}(x_i^{(1)} + \frac{3}{7}x_i^{(2)}) = x_i^{(1)} \quad \checkmark$$

$$v=2, \operatorname{sgn}(b_i^{(2)}) = \operatorname{sgn}(\frac{3}{7}x_i^{(1)} + x_i^{(2)}) = x_i^{(2)} \quad \checkmark$$

$$v=3, \operatorname{sgn}(b_i^{(3)}) = \operatorname{sgn}(\frac{19}{35}x_i^{(1)} + \frac{1}{5}x_i^{(2)}) = x_i^{(1)} \quad \checkmark$$

$$v=4, \operatorname{sgn}(b_i^{(4)}) = \operatorname{sgn}(\frac{13}{35}x_i^{(1)} + \frac{29}{35}x_i^{(2)}) = x_i^{(2)} \quad \checkmark$$

$$v=5, \operatorname{sgn}(b_i^{(5)}) = \operatorname{sgn}(-x_i^{(1)} - \frac{3}{7}x_i^{(2)}) = -x_i^{(1)} = x_i^{(5)} \quad \checkmark$$

$$\therefore x_i^{(5)} = -x^{(1)}$$

\Rightarrow Only pattern 1, 2 and 5 remain same.

 \checkmark

2/2

3. (a) Take points $(-2, 1), (0, 1), (-1, -2), (-1, 3)$

$$\bullet(-1, 3) \rightarrow -1$$

$$+1 \leftarrow (-2, 1) \bullet \bullet (0, 1) \rightarrow +1$$

$$\circ(-1, -2) \rightarrow -1$$

We cannot construct a line (decision boundary) that separates the points mentioned above. Thus problem's not linearly separable. OK

(b) On decision boundary the argument of signum function is zero.

$$\text{Thus, } w_{11}x_1 + w_{12}x_2 - \theta_1 = 0$$

$(-1, -1)$ and $(-3, 2)$ lie on this line \Rightarrow

~~$w_{11} + w_{12} - \theta_1 = 0$~~ $-w_{11} - w_{12} = \theta_1$

~~$-3w_{11} + 2w_{12} - \theta_1 = 0$~~ $-3w_{11} + 2w_{12} = \theta_1$

$\text{Take } w_{11} = 1 \Rightarrow 2w_{11} - 3w_{12} = 0$

$\Rightarrow w_{12} = 2/3 \quad \& \quad \theta_1 = -5/3 \quad \text{OK}$

$\text{And, } w_{21}x_1 + w_{22}x_2 = \theta_2$

$(-3, 2)$ & $(2, 3)$ lie on this line \Rightarrow

$-3w_{21} + 2w_{22} = \theta_2$

$2w_{21} + 3w_{22} = \theta_2$

$\Rightarrow 5w_{21} + w_{22} = 0$

$\text{Take } w_{22} = 1 \Rightarrow w_{21} = -1/5 \quad \& \quad \theta_2 = 13/5$

OK

$w_{31}x_1 + w_{32}x_2 = \theta_3$

$(2, 3)$ & $(-1, -1)$ lie on this line \Rightarrow




$2w_{31} + 3w_{32} = \theta_3 \Rightarrow 3w_{31} + 4w_{32} = 0$

$-w_{31} - w_{32} = \theta_3$

OK

$\text{Take } w_{31} = +1 \Rightarrow w_{32} = -3/4 \quad \& \quad \theta_3 = -1/4$

3(c)



We can draw a plane passing through the (x) marks above.

$$\text{i.e. } P_1(0, -1, -1), P_2(1, 0, -1), P_3(1, -1, 0)$$

$$\Rightarrow W_1V_1 + W_2V_2 + W_3V_3 = \theta$$

$$P_1 \Rightarrow -W_2 - W_3 = \theta \quad \text{--- (1)}$$

$$P_2 \Rightarrow W_1 - W_3 = \theta \quad \text{--- (2)}$$

$$P_3 \Rightarrow W_1 - W_2 = \theta \quad \text{--- (3)}$$

V_j	V_1	V_2	V_3	θ
	-1	-1	-1	-1
	-1	-1	+1	-1
	-1	+1	-1	-1
	-1	+1	+1	(-)
	+1	-1	-1	+1
	+1	-1	+1	-1
	+1	+1	-1	-1
	+1	+1	+1	-1

$$(2) - (3) \Rightarrow W_2 - W_3 = 0 \Rightarrow W_2 = W_3 = -\theta/2 \text{ or } \theta = -2W_3$$

$$\text{Take } W_1 = 1$$

$$\therefore (2) \Rightarrow W_1 - W_3 = -2W_3$$

$$\Rightarrow W_3 = -1 = W_2$$

$$\therefore \theta = \underline{\underline{2}}$$

$$\text{So, } W = \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix} \text{ and } \theta = 2$$


OK (Output layer)

$$\& \quad W = \begin{bmatrix} 1 & 2/3 \\ -1/5 & 1 \\ 1 & -3/4 \end{bmatrix} \quad \& \quad \theta = \begin{bmatrix} -5/3 \\ 13/5 \\ -1/4 \end{bmatrix}$$

(Hidden layer)

OK

4(a)



$$\bullet = +1$$

$$\circ = -1$$

Simple perception cannot solve this problem as the problem is linearly not separable and simple perceptions can only solve linearly separable problems. OK

(b)

μ	x_1^μ	x_2^μ	x_3^μ	t^μ	g_1^μ	g_2^μ	
1.	1	0	1	0	-1	e^{-1}	e^{-2}
2	1	0	1	-1	e^{-2}	e^{-1}	
3	0	0	0	+1	1	e^{-3}	
4	1	0	0	-1	e^{-1}	e^{-2}	
5	1	1	0	-1	e^{-2}	e^{-1}	
6	0	0	1	-1	e^{-1}	e^{-2}	
7	0	1	1	-1	e^{-2}	e^{-1}	
8	1	1	1	+1	e^{-3}	1	

$$w_1 = (0, 0, 0)$$

$$w_2 = (1, 1, 1)$$

OK

$$\mu=1, g_1^{(1)} = \exp(-|((0, 1, 0))^T|^2) = \exp(-1)$$

$$g_2^{(1)} = \exp(-|((1, 0, 1))^T|^2) = \exp(-2)$$

$$\mu=2, g_1^{(2)} = \exp(-|((1, 0, 1))^T|^2) = \exp(-2) \quad \text{OK}$$

$$g_2^{(2)} = \exp(-|((0, 1, 0))^T|^2) = \exp(-1)$$

$$\mu=3, g_1^{(3)} = \exp(-|((0, 0, 0))^T|^2) = 1$$

$$g_2^{(3)} = \exp(-|((1, 1, 1))^T|^2) = \exp(-3)$$

$$\mu = 4, \quad g_1^{(4)} = \exp(-\|(1, 0, 0)\|^2) = \exp(-1)$$

~~g₂~~

$$g_2^{(4)} = \exp(-\|(0, 1, 1)\|^2) = \exp(-2)$$

$$\mu = 5, \quad g_1^{(5)} = \exp(-\|(1, 1, 0)\|^2) = \exp(-2)$$

$$g_2^{(5)} = \exp(-\|(0, 0, 1)\|^2) = \exp(-1)$$

$$\mu = 6, \quad g_1^{(6)} = \exp(-\|(0, 0, 1)\|^2) = \exp(-1)$$


$$g_2^{(6)} = \exp(-\|(1, 1, 0)\|^2) = \exp(-2)$$

$$\mu = 7, \quad g_1^{(7)} = \exp(-\|(0, 1, 1)\|^2) = \exp(-2)$$

$$g_2^{(7)} = \exp(-\|(1, 0, 0)\|^2) = \exp(-1)$$

$$\mu = 8, \quad g_1^{(8)} = \exp(-\|(1, 1, 1)\|^2) = \exp(-3)$$

$$g_2^{(8)} = \exp(-\|(0, 0, 0)\|^2) = 1$$



Here
 $\bullet \rightarrow +1$
 $\circ \rightarrow -1$

OK

Let weight vector be $(w_1, w_2)^T$


and threshold θ .

∴ On decision boundary argument of signum function should be zero.

$$\therefore \theta = \text{sgn}(w_1 g_1 + w_2 g_2 - \theta)$$

$$\Rightarrow w_1 g_1 + w_2 g_2 = \theta \quad \text{on decision boundary!}$$

Points $(1, 0)$ and $(0, 1)$ lie on this line.



$$(g_1, g_2) = (1, 0) \Rightarrow w_1 = \theta$$

$$(g_1, g_2) = (0, 1) \Rightarrow w_2 = \theta$$

Choose $w_1 = 1 \Rightarrow w_2 = 1 = \theta$

Thus $w = (1, 1)^T$ and $\theta = 1$. OK

5.)

$$v_j^{(M)} = g\left(\sum_{k=1}^3 w_{jk} x_k^{(M)} - \theta_j\right) \quad \& \quad b_j^M = \sum_{k=1}^3 w_{jk} x_k^{(M)} - \theta_j$$

$$o_1^{(M)} = g\left(\sum_{j=1}^5 w_{1j} v_j^{(M)} - \theta_1\right) \quad \& \quad B_1^M = \sum_{j=1}^5 w_{1j} v_j^{(M)} - \theta_1$$

$$H = \frac{1}{2} \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)})^2 \quad \text{OK}$$

Now,

$$\delta w_{1jm} = -\eta \frac{\partial H}{\partial w_{1jm}} \quad \text{OK}$$

$$= +\eta \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)}) \frac{\partial o_i^{(\mu)}}{\partial w_{1jm}}$$

$$= \eta \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)}) \frac{\partial o_i^{(\mu)}}{\partial B_1^M} \cdot \frac{\partial B_1^M}{\partial w_{1jm}}$$

$$= \eta \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)}) g'(B_1^M) \left[\sum_{j=1}^5 \delta_{mj} v_j^{(M)} \right] \quad \text{OK}$$

$$\delta w_{1jm} = \eta \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)}) g'(B_1^M) v_m^{(M)}$$

We get update rule for θ_1 if we replace $v_j^{(M)}$ by -1.

$$\Rightarrow \delta \theta_1 = -\eta \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)}) g'(B_1^M) \quad \text{OK}$$

Now for the hidden layer:

$$\delta w_{mn} = -\eta \frac{\partial H}{\partial w_{mn}} = -\cancel{\eta} \cancel{\partial}$$

$$= \eta \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)}) \frac{\partial o_i^{(\mu)}}{\partial w_{mn}}$$

$$= \eta \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)}) \frac{\partial o_i^{(\mu)}}{\partial B_1^M} \cdot \frac{\partial B_1^M}{\partial w_{mn}}$$

$$= \eta \sum_{\mu=1}^p (t_i^{(\mu)} - o_i^{(\mu)}) g'(B_1^M) \left[\sum_{j=1}^5 w_{1j} \frac{\partial v_j^{(M)}}{\partial w_{mn}} \right] \quad \text{OK}$$

CHALMERS	Anonymous code	FFR135-0152-PXZ	Points for question (to be filled in by teacher)	Consecutive page no. Löpande sid nr
	Anonym kod		Poäng på uppgiften (fylltes av lärare)	12

Poäng på uppgiften
(fylltes av lärare)Question no.
Uppgift nr

5

2-

$$\begin{aligned}
 \text{Now, } \frac{\partial V_j^M}{\partial w_{mn}} &= \frac{\partial V_j^M}{\partial b_j^M} \cdot \frac{\partial b_j^M}{\partial w_{mn}} \\
 &= g'(b_j^M) \cdot \left(\sum_{k=1}^3 \delta_{jm} \delta_{kn} x_k^M \right) \quad \text{OK} \\
 &= g'(b_j^M) x_n^M \delta_{jm} \quad \text{OK}
 \end{aligned}$$

$$\begin{aligned}
 \text{So, } \delta w_{mn} &= \eta \sum_{i=1}^P (t_i^{(M)} - o_i^{(M)}) g'(B_i^{(M)}) w_{im} g'(b_m^{(M)}) x_n^{(M)} \\
 &= \eta \sum_{i=1}^P (t_i^{(M)} - o_i^{(M)}) g'(B_i^{(M)}) g'(b_m^{(M)}) w_{im} x_n^{(M)} \quad \text{OK}
 \end{aligned}$$

For thresholds update rule simply replace $x_n^M \rightarrow -1$

$$\Rightarrow \delta \theta_m = -\eta \sum_{i=1}^P (t_i^{(M)} - o_i^{(M)}) g'(B_i^{(M)}) g'(b_m^{(M)}) w_{im} \quad \text{OK}$$

Sequential
training?

CHALMERS	Anonymous code	FFR135-0152-PXZ	Points for question (to be filled in by teacher)	Consecutive page no. Löpande sid nr
	Anonym kod		Poäng på uppgiften (fyller av lärare)	Question no. Uppgift nr

1

6. Input layer. - This layer feeds the input image in all channels into the neural network.

Convolution layer - This layer performs convolution operation with kernels on the input layer to get feature maps which capture features of images. ✓

Max-pooling layer - This layer takes maximum value in a receptive area thus summarizing the feature maps or the output of previous layer. ✓

Fully-connected: This layer takes the output from max-pooling layer and acts as buffer layer before feeding its own output to the output layer. ✓

Output layer: This layer produces the final output of the network. (classes or real values). ✓

$$\rightarrow x_1 = 31, y_1 = 31, z_1 = 3 \quad \checkmark$$

\rightarrow if j is the number of strides, ~~then's~~'s step size then

$$1 + js = x_1 - (3-1) \quad \because 3 \text{ is field size.}$$

$$\Rightarrow j = \frac{x_1 - 3}{s}$$

$$\Rightarrow x_2 = j+1 \quad (\because \text{one more than the no. of strides.})$$

$$= \frac{x_1 - 3}{s} + 1$$

$$= \frac{31-3}{2} + 1 = 15$$

$$\Rightarrow x_2 = 15, y_2 = 15 \quad (\because \text{similar calculation}), z_2 = 10$$

$$\rightarrow x_3 = x_2/5 \quad (\because \text{max-pooling layer})$$

$$x_3 = 3 \quad \checkmark$$

$$\text{Thus, } y_3 = 3 \quad \checkmark \text{ and } z_3 = 10 \quad \checkmark \text{ (same as } z_2).$$

CHALMERS	Anonymous code	FFR135-0152-PXZ	Points for question (to be filled in by teacher)	Consecutive page no. Löpande sid nr
	Anonym kod		Poäng på uppgiften (fyller av lärare)	Question no. Uppgift nr
				14 6

Also, $y_4 = 10$ ✓ and $y_5 = 5$ ✓

→ Now for convolution layer, no. of weights = $(3 \times 3 \times 3) = 27$
& each feature map has one threshold
⇒ total parameters = $(27 + 1) \times 10$
= 280 ✓

→ No trainable parameters between convolution layer & max-pooling layer. ✓

→ Between max-pooling & fully connected layer:

No. of weights = ~~$(3 \times 3) \times 2 \times 10$~~ $(z_3 \times y_3) \times z_3 \times y_4$
= $3 \times 3 \times 10 \times 10$
= 900 ✓

⇒ ~~No. of thresholds~~, No. of thresholds = 10 (= y_4)
⇒ total = 910 ✓

→ In between output layer & fully connected layer

No. of weights = $y_5 \times y_4 = 50$ ✓
No. of thresholds = $y_5 = 5$ ✓
∴ total = 55 ✓

2/2

Rough Work

4 → ⑥

4.60
6
40

$$\textcircled{6} \quad n'n' - nn + 2nn'$$

$$n'n' + nn' + nn' - nn$$

$$n(n+n) + n(n'-n)$$

35-16
35-22

$$b_m = \sum_{i+m} w_{min} n_i + w_{mm} n_m - \mu_m n_m$$

n=0 n=1 4, 15

n=i n=0

→ -11

8, 5

$$\sum_{i+m} w_{min} n_i + w_{mm} < \mu_m$$

$$w_{mm} - \mu_m =$$

$$b_m \leq w_{mm} - \mu_m$$

$$b_m n_m \cdot b_m \theta_H(b_m)$$

F

$$b_m n_m \cdot -b_m - \frac{w}{2} < 0$$

$$1 \rightarrow 0 \Rightarrow \frac{w}{2} - \mu_m$$

$$b_m < 0$$

35-28

35-6

$$\sum \phi < \mu_m \quad 35-50$$


-15

$$9 \times 3 = 27 + \textcircled{1} \times 10$$

$$x_1 = 31, y_1 = 31, z_1 = 3$$

$$x_2 = 15, y_2 = 15, z_2 = 10$$

$$x_3 = 3, y_3 = 3, z_3 = 10$$



$$y_4 = 10 \quad -\frac{2}{5} + 3.$$

$$8 \cdot 5 - \frac{(3-1)}{2}$$

$$7 - \quad 1 + s_x$$

$$\frac{13}{5}$$

$$\frac{4.60}{6} = \textcircled{10}$$

120 ✓

$$\frac{5-3-1}{2}$$

$$1 \rightarrow 1+s \rightarrow 1+2s$$

$$1+js = N - (n-1) \quad \textcircled{6} \quad js = \frac{(n-1)}{s}$$

$$1+s_j = N - \frac{n-1}{s}$$

$$\frac{31-3}{2} \quad \textcircled{14+1} \quad 15$$

$$\frac{N-n}{s} + 1$$