
Bulk · normalization — complementary evidence


A

Library-size distributions by condition




B

Paired donor RLE medians before/after normalization



C

Reference-feature normalization sensitivity



Estimate [interval] · q
Independent n = 24

STAT1 · Primary	-0.70 [-0.88, -0.52] 0.002
STAT1 · Alternative	-0.62 [-0.80, -0.44] 0.005
IRF7 · Primary	-0.50 [-0.68, -0.32] 0.02
IRF7 · Alternative	-0.42 [-0.60, -0.24] 0.023
ISG15 · Primary	-0.30 [-0.48, -0.12] 0.038
ISG15 · Alternative	-0.22 [-0.40, -0.04] 0.041
COL1A1 · Primary	-0.10 [-0.28, 0.08] 0.65
COL1A1 · Alternative	-0.02 [-0.20, 0.16] 0.65
NDUFA1 · Primary	0.10 [-0.08, 0.28] 0.65
NDUFA1 · Alternative	0.18 [0.00, 0.36] 0.077
SRSF1 · Primary	0.30 [0.12, 0.48] 0.092
SRSF1 · Alternative	0.38 [0.20, 0.56] 0.095
MKI67 · Primary	0.50 [0.32, 0.68] 0.11
MKI67 · Alternative	0.58 [0.40, 0.76] 0.11
KLF2 · Primary	0.70 [0.52, 0.88] 0.13
KLF2 · Alternative	0.78 [0.60, 0.96] 0.13
JUN · Primary	0.90 [0.72, 1.08] 0.15
JUN · Alternative	0.98 [0.80, 1.16] 0.15

Software test fixture; not biological evidence