
Single cell · multi_contrast — complementary evidence


A

Independent support across contrast families




B

Complete signed effect atlas



C

Cross-contrast replication effects



Estimate [interval] · q
Independent n = 12

T cells · Cohort A	-0.70 [-0.88, -0.52] 0.002
T cells · Cohort B	-0.62 [-0.80, -0.44] 0.005
B cells · Cohort A	-0.50 [-0.68, -0.32] 0.02
B cells · Cohort B	-0.42 [-0.60, -0.24] 0.023
Myeloid · Cohort A	-0.30 [-0.48, -0.12] 0.038
Myeloid · Cohort B	-0.22 [-0.40, -0.04] 0.041
Stromal · Cohort A	-0.10 [-0.28, 0.08] 0.65
Stromal · Cohort B	-0.02 [-0.20, 0.16] 0.65
Endothelial · Cohort A	0.10 [-0.08, 0.28] 0.65
Endothelial · Cohort B	0.18 [0.00, 0.36] 0.077
NK cells · Cohort A	0.30 [0.12, 0.48] 0.092
NK cells · Cohort B	0.38 [0.20, 0.56] 0.095
Pericytes · Cohort A	0.50 [0.32, 0.68] 0.11
Pericytes · Cohort B	0.58 [0.40, 0.76] 0.11
Mast cells · Cohort A	0.70 [0.52, 0.88] 0.13
Mast cells · Cohort B	0.78 [0.60, 0.96] 0.13
Epithelial · Cohort A	0.90 [0.72, 1.08] 0.15
Epithelial · Cohort B	0.98 [0.80, 1.16] 0.15