
Leave-one-cohort stability
Estimate [interval] · q
T cells · Full cohorts
T cells · Omit cohort A
B cells · Full cohorts
B cells · Omit cohort A
Myeloid · Full cohorts
Myeloid · Omit cohort A
Stromal · Full cohorts
Stromal · Omit cohort A
Endothelial · Full cohorts
Endothelial · Omit cohort A
NK cells · Full cohorts
NK cells · Omit cohort A
-0.70 [-0.88, -0.52] 0.002 n=12
-0.62 [-0.80, -0.44] 0.005 n=8
-0.38 [-0.56, -0.20] 0.02 n=12
-0.30 [-0.48, -0.12] 0.023 n=8
-0.06 [-0.24, 0.12] 0.65 n=12
0.02 [-0.16, 0.20] 0.65 n=8
0.26 [0.08, 0.44] 0.056 n=12
0.34 [0.16, 0.52] 0.059 n=8
0.58 [0.40, 0.76] 0.074 n=12
0.66 [0.48, 0.84] 0.077 n=8
0.90 [0.72, 1.08] 0.092 n=12
0.98 [0.80, 1.16] 0.095 n=8
−1.0
−0.5
0.0
0.5
1.0
STAT1 donor log2 fold change · supplied 95% interval