Single cell · resolution — complementary evidence

A

Donor support at declared resolutions



B

Donor mixture within selected clusters



C

Independent annotation consistency



Estimate [interval] · q
Independent n = 12

T cells · Cohort A	-0.22 [-0.30, -0.14] 0.002
T cells · Cohort B	-0.14 [-0.22, -0.06] 0.005
B cells · Cohort A	-0.16 [-0.24, -0.08] 0.02
B cells · Cohort B	-0.08 [-0.16, 0.00] 0.65
Myeloid · Cohort A	-0.10 [-0.18, -0.02] 0.038
Myeloid · Cohort B	-0.02 [-0.10, 0.06] 0.65
Stromal · Cohort A	-0.03 [-0.11, 0.05] 0.65
Stromal · Cohort B	0.05 [-0.03, 0.13] 0.65
Endothelial · Cohort A	0.03 [-0.05, 0.11] 0.65
Endothelial · Cohort B	0.11 [0.03, 0.19] 0.077
NK cells · Cohort A	0.10 [0.02, 0.18] 0.092
NK cells · Cohort B	0.18 [0.10, 0.26] 0.095
Pericytes · Cohort A	0.16 [0.08, 0.24] 0.11
Pericytes · Cohort B	0.24 [0.16, 0.32] 0.11
Mast cells · Cohort A	0.22 [0.14, 0.30] 0.13
Mast cells · Cohort B	0.30 [0.22, 0.38] 0.13
Epithelial · Cohort A	0.29 [0.21, 0.37] 0.15
Epithelial · Cohort B	0.37 [0.29, 0.45] 0.15