❖ MID RNS (35)

Histology

1- Which is correct about Macula densa:

A) has no basement membrane B) has thick basement membrane

C) located in wall of afferent arteriole D) located in PCT

2- About Podocyte:

A) specialized epithelial cell B) Neuroepithelium

C) connective tissue cell D) non-branched cell

3- All of the following are parts of nephron except:

A) PCT B) DCT

C) Loop of Henle D) collecting tubule

Anatomy

4- Length of ureter:

A) 10 inches B) 5 inches C) 15 inches D) 25 inches

5- The shortest part of urethra is:

A) membranous B) prostatic C) internal urethral meatus D) penile

6- Abnormal fusion of lower poles of both kidneys leads to:

A) Horseshoe kidney B) Rosette shaped kidney

C) Congenital polycystic kidney D) Pelvic kidney

7- All of the following are anterior relations of right kidney except:

A) spleen B) right Supra renal gland

C) right colic flexure D) 2nd part of duodenum

Biochemistry

8- Stone-inhibitor substances include:

A) Citrate B) Vitamin D

C) Vitamin C D) Phosphate

Parasitology

9- All of the following could be found in urine EXCEPT:

A) Echinococcus granulosus eggs

B) Trichomonas vaginalis trophozoite

C) Strongyloides stercoralis larva

D) Wuchereria bancrofti microfilaria

Pharmacology

10- Drug causes tubular cell toxicity:

A) Aminoglycoside B) Acetaminophen

C) Sulfonamides D) Clopidogrel

11- Thrombotic microangiopathy caused by which drug:

A) Clopidogrel B) Amphotericin B C) Sulfonamides

12- Statement shows correct relation between disease and drug caused:

A) Crystal nephropathy - Sulfonamides

B) Thrombotic microangiopathy - Aminoglycoside

C) tubular cell toxicity - Acetaminophen

Pathology

13- The outer surface of kidney shows tiny abscesses in the cortex and medulla in:

A) toxic tubular necrosis B) ischemic tubular necrosis

C) acute pyelonephritis D) interstitial nephritis

14- Which of the following show irregular depressions due to cortical scarring:

A) acute glomerulonephritis B) acute pyelonephritis

C) chronic pyelonephritis D) chronic glomerulonephritis

15- Acute diffuse proliferative glomerulonephritis is due to:

A) Blood-borne infection B) Ascending infection

C) Deposition of immune complexes in the glomeruli D) Autoimmune reaction

---

16- Edema of the nephritic syndrome appears in:

A) Face B) Lower limbs

C) Back D) All-over the body

17- The following is NOT a cause of nephrotic syndrome:

A) Acute diffuse proliferative glomerulonephritis

B) Membranous glomerulonephritis

C) Membrano- proliferative glomerulonephritis

D) Minimal change glomerulonephritis

18- Epithelial crescents are characteristic of:

A) Rapidly progressive glomerulonephritis

B) Membranous glomerulonephritis

C) Membrano-proliferative glomerulonephritis

D) Minimal change glomerulonephritis

Microbiology

19- Which of the following cause non gonococcal UTI ?

A) Neisseria gonorrhoea B) Mycoplasma hominis

C) E. Coli D) proteus mirabilis

20- Child with recurrent UTI mainly has:

A) ureteric and renal calculi B) vesico ureteric reflux

C) neurogenic bladder D) urethral valve

Physiology

21- H secretion decreases in case of:

A) increased plasma Co2 concentration B) carbonic anhydrase inhibitor

C) decreased K secretion D) increased Na reabsorption

22- Glucose transport through luminal border by:

A) Na cotransport B) passive diffusion C) H cotransport

23- it's correct to say:

A) renin secretion is inversely proportional to sodium concentration in DCT

24- it's correct to say :

A) below renal threshold non substance appears in urine

B) sodium has Tm

25- counter current multiplier :

A) medullary collecting duct B) vasa recta in juxtamedullary nephron

C) loop of Henle in juxtamedullary nephron D) loop of Henle in cortical nephron

---

26- Aldosterone increases Na reabsorption at:

A) proximal convoluted tubule B) thick ascending limb

C) descending loop of Henle D) DCT and collecting duct

27- Tubular fluid is more concentrated in:

A) descending Loop of Henle B) early ascending loop of Henle C) proximal convoluted tubule

---

28- Which of the following lead to increase of GFR :

A) contraction of afferent arterioles B) efferent arterioles mildly contracted

C) efferent arterioles severely contracted D) contraction of mesangial cell

---

ANSWERS									
1	**2**	**3**	**4**	**5**	**6**	**7**	**8**	**9**	**10**
A	A	D	A	A	A	A	A	A	A
11	**12**	**13**	**14**	**15**	**16**	**17**	**18**	**19**	**20**
A	A	C	C	C	A	A	A	B	B
21	**22**	**23**	**24**	**25**	**26**	**27**	**28**	**29**	**30**
B	A	A	A	C	D	A	B		