

1. Most excitatory neurotransmission in the brain is believed to be mediated by:

A- Glutamate
GnRH.

B. glycine

C. GABA

D.

2. Synaptic conduction is mostly orthodromic (one way conduction) because:

A. dendrites cannot be depolarized.

B. once repolarized, an area cannot be depolarized.

C. the strength of antidromic impulses is less.

D- the neurotransmitter is in the presynaptic terminal.

3- At chemical synapses, excitatory postsynaptic potentials (EPSPs) can be produced by:

A- Opening of Na⁺ channels

B- Opening of K⁺ channels

C- Opening of Cl⁻ channels

D- Closure of Ca⁺ channels.

4- Opening of ligand-gated Cl⁻ channels causes :

A. inhibition of the postsynaptic neuron

B. depolarization of the postsynaptic neuron

C. initiation of an action potential

D. block of ligand-gated cation channels.

5- Temporal summation

A - Occurs when several presynaptic fibers are stimulated simultaneously.

B- Occurs when presynaptic fiber is repeatedly stimulated.

C- Is due to overlap of the discharging zone.

d- None of the above is correct.

6- Which reflexes property of the following describe the spreading of the impulses to wider areas?

A. Occlusion

B. lateral inhibition.

C. Divergence

D. Convergence.