



# Random Variables

Chris Piech

CS109, Stanford University

# Ultimate Probability



Ultimate Probability

2,734 views

11 0 SHARE SAVE ...



Maika Isogawa

Published on 1 Dec 2018

SUBSCRIBE 11

<https://www.youtube.com/watch?v=H2IfTwGisOg>





# Let's Play a Game

- Game set-up
  - We have a fair coin (come up “heads” with  $p = 0.5$ )
  - Let  $n$  = number of coin flips (“heads”) before first “tails”
  - You win  $\$2^n$
- How much would you pay to play?



# *Mutual exclusion And Independence*

Are two properties of events that make it easy to calculate probabilities.



# Conditional Probability



$$P(E|F) = \frac{P(EF)}{P(F)}$$

What is your new belief that E will occur, given that you have observed F occurred





In the conditional  
paradigm, the formulas of  
probability are preserved.



# BAE's Theorem?

$$P(A \mid B \wedge E) = \frac{P(B \mid A \wedge E) P(A \mid E)}{P(B \mid E)}$$





# Learning Goals

1. Be able to use conditional independence definition
2. Be able to define a random variable (R.V.)
3. Be able to use and produce a PMF of a R.V.
4. Be able to calculate the expectation of the R.V.





$G_1$

$G_2$





$G_3$

$G_4$

$G_5$

**T**





```
dna.txt — dna
dna.txt
1 False,True,False,False,True,False
2 True,True,False,True,True,False
3 True,True,False,True,True,True
4 False,True,False,True,True,False
5 False,True,False,False,True,False
6 True,True,False,True,True,True
7 False,False,True,False,False,False
8 False,False,True,False,True,False
9 True,False,False,True,False,False
10 False,True,False,True,True,False
11 True,False,False,True,False,False
12 True,False,True,True,False,False
13 False,True,False,False,True,False
14 False,False,True,True,False,False
15 True,True,False,False,True,True
16 True,False,True,True,False,False
17 True,True,True,True,True,True |
18 True,False,True,False,False,True
19 False,True,False,True,True,True
20 False,False,True,False,False,False
21 False,False,False,True,True,False
22 False,True,False,False,True,False
23 True,True,False,True,True,True
24 False,True,False,True,True,False
25 True,False,False,False,False,True
26 False,False,True,True,False,True
27 False,False,False,True,False,False
28 False,True,True,False,False,True
29 False,True,False,False,True,True
30 False,False,False,False,False,True
31 False,True,False,True,True,False
32 True,False,False,True,False,False
33 True,True,False,True,True,True
34 True,True,False,False,True,True
35 True,True,False,True,True,True
36 False,False,True,True,False,False
--
```

100,000  
samples

6 observations per sample



# Discovered Pattern

```
[Piech-2:dna piech$ python findStructure.py  
size data = 100000  
p(G1) = 0.500  
p(G2) = 0.545  
p(G3) = 0.299  
p(G4) = 0.701  
p(G5) = 0.600  
p(T) = 0.390  
p(T and G1) = 0.291 , P(T)p(G1) = 0.195  
p(T and G2) = 0.300 , P(T)p(G2) = 0.213  
p(T and G3) = 0.116 , P(T)p(G3) = 0.117  
p(T and G4) = 0.273 , P(T)p(G4) = 0.273  
p(T and G5) = 0.309 , P(T)p(G5) = 0.234
```

• • •

$$\begin{aligned} p(T \text{ and } G5 \mid G2) &= 0.450 \\ p(T \mid G2)p(G5 \mid G2) &= 0.450 \end{aligned}$$





Independence  
relationships can change  
with conditioning.

If  $E$  and  $F$  are independent, that does not mean they will still  
be independent given another event  $G$ .

*There is additional reading about this in the course reader. You will explore this more in depth in CS228*





# Two Great Tastes

Conditional Probability

Independence



# Conditional Independence

- Two events  $E$  and  $F$  are called **conditionally independent given  $G$** , if

$$P(EF|G) = P(E|G)P(F|G)$$

- Or, equivalently if:

$$P(E|FG) = P(E|G)$$



# Conditional Paradigm

- For any events A, B, and E, you can condition consistently on E, and all formulas still hold:

$$P(A \text{ B} \mid E) = P(B \text{ A} \mid E)$$

$$P(A \text{ B} \mid E) = P(A \mid B \text{ E}) P(B \mid E)$$

- Can think of E as “everything you already know”
- Formally,  $P(\bullet \mid E)$  satisfies 3 axioms of probability



# NETFLIX

**And Learn**



# Netflix and Learn

What is the probability  
that a user will watch  
Life is Beautiful?

$$P(E)$$



$$P(E) = \lim_{n \rightarrow \infty} \frac{n(E)}{n} \approx \frac{\text{\#people who watched movie}}{\text{\#people on Netflix}}$$


$$P(E) = 10,234,231 / 50,923,123 = 0.20$$



# Netflix and Learn

What is the probability  
that a user will watch  
Life is Beautiful, given  
they watched Amelie?

$$P(E|F)$$



$$P(E|F) = \frac{P(EF)}{P(F)} = \frac{\text{\#people who watched both}}{\text{\#people who watched } F}$$


$$P(E|F) = 0.42$$




Conditioned on liking a set of movies?

# Netflix and Learn


Each event corresponds to liking a particular movie




$E_1$



$E_2$



$E_3$



$E_4$

$$P(E_4|E_1, E_2, E_3)?$$




Is  $E_4$  independent of  $E_1, E_2, E_3$ ?

# Netflix and Learn


Is  $E_4$  independent of  $E_1, E_2, E_3$ ?




$E_1$



$E_2$



$E_3$



$E_4$


$$P(E_4|E_1, E_2, E_3) \stackrel{?}{=} P(E_4)$$

# Netflix and Learn


Is  $E_4$  independent of  $E_1, E_2, E_3$ ?




$E_1$



$E_2$



$E_3$



$E_4$

$$P(E_4|E_1, E_2, E_3) = \frac{P(E_1 E_2 E_3 E_4)}{P(E_1 E_2 E_3)}$$

# Netflix and Learn

- What is the probability that a user watched four particular movies?
  - There are 13,000 titles on Netflix
  - The user watches 30 random titles.
  - $E$  = movies watched include the given four.

- Solution:

$$P(E) = \frac{\overset{\text{Watch those four}}{\binom{4}{4}} \overset{\text{Choose 24 movies not in the set}}{\binom{12996}{24}}}{\underset{\text{Choose 30 movies from netflix}}{\binom{13000}{30}}} = 10^{-11}$$






# Netflix and Learn




$E_1$



$E_2$



$E_3$




$E_4$




# Netflix and Learn

$K_1$


*Like foreign emotional comedies*




$E_1$



$E_2$



$E_3$



$E_4$

Assume  $E_1$ ,  $E_2$ ,  $E_3$  and  $E_4$  are conditionally independent given  $K_1$



# Netflix and Learn

$K_1$


*Like foreign emotional comedies*



Assume  $E_1$ ,  $E_2$ ,  $E_3$  and  $E_4$  are conditionally independent given  $K_1$




# Netflix and Learn



Assume  $E_1$ ,  $E_2$ ,  $E_3$  and  $E_4$  are conditionally independent given  $K_1$



# Netflix and Learn



Assume  $E_1, E_2, E_3$  and  $E_4$  are conditionally independent given  $K_1$





Conditional independence is a practical, real world way of decomposing hard probability questions.

# Conditional Independence



If  $E$  and  $F$  are  
dependent,  
that does not mean  $E$  and  
 $F$  will be dependent  
when another event is  
observed.



# Conditional Dependence



If  $E$  and  $F$  are  
independent,

that does not mean  $E$  and  
 $F$  will be independent  
when another event is  
observed.



# Big Deal

“Exploiting conditional independence to generate fast probabilistic computations is one of the main contributions CS has made to probability theory”

-Judea Pearl wins 2011 Turing Award, *“For fundamental contributions to artificial intelligence through the development of a calculus for probabilistic and causal reasoning”*



Ready for the next (cs109) episode



# Random Variables

# Remember Learning to Code?

*type*

*name*

*value*

```
int a = 5;  
double b = 4.2;  
bit c = 1;  
choice d = medium;
```

$z \in \{\text{high, medium, low}\}$

Random variables are like programming variables, with uncertainty

# Pirates of the Random Variables

**int** a = 5;

$A$  is the number of pirate ships in our *future* armada.

$$A \in \{1, 2, \dots, 10\}$$



**double** b = 4.2;

$B$  is the amount of money we get *after* we defeat Blackbeard.

$$B \in \mathbb{R}^+$$



**bit** c = 1;

$C$  is 1 *if* we successfully raid Isla de Muerta. 0 otherwise.

$$C \in \{0, 1\}$$



# Random Variable

- A **Random Variable** is a variable will have a value. But there is uncertainty as to what value.
- Example:
  - 3 fair coins are flipped.
  - $Y$  = number of “heads” on 3 coins
  - **$Y$  is a random variable**
  - $P(Y = 0) = 1/8$  (T, T, T)
  - $P(Y = 1) = 3/8$  (H, T, T), (T, H, T), (T, T, H)
  - $P(Y = 2) = 3/8$  (H, H, T), (H, T, H), (T, H, H)
  - $P(Y = 3) = 1/8$  (H, H, H)
  - $P(Y \geq 4) = 0$

It is confusing that both random variables  
and events use the same notation





Random variables and  
events are two *different*  
things





We can define an event to  
be a particular assignment  
to a random variables



# Example Random Variable

- Consider 5 coin flips, each which independently come up heads with probability  $p$

- Recall:

$$P(2 \text{ heads}) = \binom{5}{2} p^2 (1 - p)^3$$

$$P(3 \text{ heads}) = \binom{5}{3} p^3 (1 - p)^2$$

- $Y$  = number of “heads” on 5 flips

$$Y \in \{1, 2, \dots, 5\}$$

$$P(Y = k) = \binom{5}{k} p^k (1 - p)^{5-k}$$

\* Pro tip: no coin works like this... but many real world binary events do

# Fun with Random Variables

- Probability Mass Function:

$$P(X = a)$$

- Expectation:

$$E[X]$$

- Variance:

$$\text{Var}(X)$$



Learning  
goals for  
today

# 1. Probability Mass Function



All the different assignments to a random variable make a function

Let  $Y$  be a random variable



$Y$



For example  $Y$  is the number of heads in 5 coin flips

$$Y = 2$$

It is an *event* when  
Y takes on a value

For example Y is the number of heads in 5 coin flips



If this is a number


$$P(Y = 2)$$


Then this is a number  
(between 0 and 1)

For example  $Y$  is the number of heads in 5 coin flips


If this is a variable


$$P(Y = k)$$


Then this is a function

For example  $Y$  is the number of heads in 5 coin flips

# Random Variables -> Functions

$$P(Y = k)$$


A blue arrow points from the expression  $k = 5$  to the variable  $k$  in the probability expression  $P(Y = k)$ . Another blue arrow points from the probability expression  $P(Y = k)$  down to the numerical value  $0.03125$ .

$$k = 5$$
$$0.03125$$

For example  $Y$  is the number of heads in 5 coin flips

# Random Variables -> Functions

$$P(Y = k)$$

```
private double eventProbability(int k) {  
    int ways = choose(N, k);  
    double a = Math.pow(P, k);  
    double b = Math.pow(P, N-k);  
    return ways * a * b;  
}
```

```
private static final int N = 5;  
private static final double P = 0.5;
```

For example Y is the number of heads in 5 coin flips





If a random variable is  
discrete we call this function  
the **Probability Mass**  
*Function*



# Probability Mass Function

Let  $X$  be a random variable that represents the result of a **single dice roll**.  $X$  can take on the values  $\{1, 2, 3, 4, 5, 6\}$

$$P(X = x)$$


$$p(x)$$

This is shorthand  
notation for the PMF


$$p_X(x)$$

This is also shorthand  
notation for the PMF

# PMF For a Single 6 Sided Dice




# PMF for the sum of two dice



# PMF as an Equation

$$p(X = x) = \begin{cases} \frac{x-1}{36} & \text{if } x \in \mathbb{Z}, 1 \leq x \leq 6 \\ \frac{13-x}{36} & \text{if } x \in \mathbb{Z}, 7 \leq x \leq 12 \\ 0 & \text{else} \end{cases}$$

Again, this is the probability for the sum of two dice



\*errata: in lecture this formula had some small mistakes ☺


# Sanity Check

$$\sum_{\text{all } k} P(Y = k) \stackrel{?}{=}$$

# Sanity Check

$$\sum_{\text{all } k} P(Y = k) \stackrel{?}{=}$$

---






# Sanity Check

$$\sum_k P(Y = k) = 1$$

---



## 2. Expectation

# Expected Value

- The Expected Values for a discrete random variable  $X$  is defined as:

$$E[X] = \sum_{x:p(x)>0} x \cdot p(x)$$

- Note: sum over all values of  $x$  that have  $p(x) > 0$ .
- Expected value also called: **Mean**, *Expectation*, **Weighted Average**, **Center of Mass**, *1<sup>st</sup> Moment*

# Expected Value

- Roll a 6-Sided Die.  $X$  is outcome of roll

- $p(1) = p(2) = p(3) = p(4) = p(5) = p(6) = 1/6$

- $E[X] = 1\left(\frac{1}{6}\right) + 2\left(\frac{1}{6}\right) + 3\left(\frac{1}{6}\right) + 4\left(\frac{1}{6}\right) + 5\left(\frac{1}{6}\right) + 6\left(\frac{1}{6}\right) = \frac{7}{2}$

- $Y$  is random variable

- $P(Y = 1) = 1/3, \quad P(Y = 2) = 1/6, \quad P(Y = 3) = 1/2$

- $E[Y] = 1 (1/3) + 2 (1/6) + 3 (1/2) = 13/6$

# Lying with Statistics

“There are three kinds of lies:  
lies, damned lies, and statistics”

– *Mark Twain*

- School has 3 classes with 5, 10 and 150 students
- Randomly choose a class with equal probability
- $X$  = size of chosen class
- What is  $E[X]$ ?
  - $E[X] = 5 (1/3) + 10 (1/3) + 150 (1/3)$   
 $= 165/3 = 55$

# Lying with Statistics

“There are three kinds of lies:  
lies, damned lies, and statistics”

– *Mark Twain*

- School has 3 classes with 5, 10 and 150 students
- Randomly choose a student with equal probability
- $Y$  = size of class that student is in
- What is  $E[Y]$ ?
  - $E[Y] = 5 (5/165) + 10 (10/165) + 150 (150/165)$   
 $= 22635/165 \approx 137$
- Note:  $E[Y]$  is students' perception of class size
  - But  $E[X]$  is what is usually reported by schools!

# Properties of Expectation

- **Linearity:**

$$E[aX + b] = aE[X] + b$$

- Consider  $X = 6$ -sided die roll,  $Y = 2X - 1$ .
- $E[X] = 3.5$                        $E[Y] = 6$

- **Expectation of a sum** is the sum of expectations

$$E[X + Y] = E[X] + E[Y]$$

- **Unconscious statistician:**

$$E[g(x)] = \sum_x g(x)p(x)$$




Wonderful

# St Petersburg

- Game set-up

- We have a fair coin (come up “heads” with  $p = 0.5$ )
- Let  $n$  = number of coin flips (“heads”) before first “tails”
- You win  $\$2^n$

- How much would you pay to play?



# St Petersburg

- Game set-up
  - We have a fair coin (come up “heads” with  $p = 0.5$ )
  - Let  $n$  = number of coin flips (“heads”) before first “tails”
  - You win  $\$2^n$
- How much would you pay to play?
- Solution
  - Let  $X$  = your winnings
  - $$E[X] = \left(\frac{1}{2}\right)^1 2^0 + \left(\frac{1}{2}\right)^2 2^1 + \left(\frac{1}{2}\right)^3 2^2 + \left(\frac{1}{2}\right)^4 2^3 + \dots = \sum_{i=0}^{\infty} \left(\frac{1}{2}\right)^{i+1} 2^i$$
$$= \sum_{i=0}^{\infty} \frac{1}{2} = \infty$$
  - I'll let you play for \$1 thousand... but just once! Takers?

# St Petersburg + Reality

- What if Chris has only \$65,536?
  - Same game
  - If you win over \$65,536 I leave the country.
- Solution
  - Let  $X$  = your winnings
  - $$\begin{aligned} E[X] &= \left(\frac{1}{2}\right)^1 2^0 + \left(\frac{1}{2}\right)^2 2^1 + \left(\frac{1}{2}\right)^3 2^2 + \left(\frac{1}{2}\right)^4 2^3 + \dots \\ &= \sum_{i=0}^k \left(\frac{1}{2}\right)^{i+1} 2^i \text{ s.t. } k = \log_2(65,536) \\ &= \sum_{i=0}^{16} \frac{1}{2} = 8.5 \end{aligned}$$

# Learning Goals

1. Be able to use conditional independence
2. Be able to define a random variable (R.V.)
3. Be able to use + produce a PMF of a R.V.
4. Be able to calculate the expectation of the R.V.

