

**ST745, Spring 2006**

**Midterm, 1:30-2:45, Thursday, 3/2/2006**

**This is an in-class, open-note and open-book midterm exam.**

1. (20 pts) Suppose the hazard function of a random survival time (in years)  $T$  takes the form  $\lambda(t) = (\lambda/2)t^{-1/2}$ , where  $\lambda > 0$ . Do the following:

- (a) Find the cumulative hazard function, survival function and density function of  $T$ . (8 pts)
- (b) Given a random sample  $(\tilde{t}_1, \delta_1), (\tilde{t}_2, \delta_2), \dots, (\tilde{t}_n, \delta_n)$ , where  $\tilde{t}_i$  is the observed failure time or right censoring time;  $\delta_i = 1$  if  $\tilde{t}_i$  is a failure time and  $\delta_i = 0$  if  $\tilde{t}_i$  is a censoring time. Find the MLE of  $\lambda$  and its asymptotic variance. (8 pts)
- (c) Suppose we have a sample 1, 4+, 4, 9, where + indicates a right censored observation. Find the MLE of  $\lambda$  and its variance. (4 pts)

2. (20 pts) The following table gives grouped survival data in years for patients with certain disease:

time interval $i$	$n_i$	$d_i$	$w_i$
$i = 1: [0, 2)$	20	2	2
$i = 2: [2, 4)$	16	2	4
$i = 3: [4, 6)$	10	2	2

where  $n_i$  is the number of patients at the beginning of time interval  $i$ ,  $d_i$  and  $w_i$  are the number of deaths and censorings observed in that interval. **Assume** all censorings in an interval occurred at the **middle** of that interval. Do the following:

- (a) Find the life-table estimate of  $S(6) = P(T \geq 6)$ . (10 pts)
  - (b) Find the variance estimate for the estimate you got in (a). (10 pts)
3. (40 pts) The survival times of 10 patients with certain heart disease are given as follows:

2, 3, 4+, 5, 5+, 6+, 8, 10, 12, 15+,

where “+” means a right censored survival time, do the following:

- (a) Find the Nelson-Aalen estimator of the cumulative hazard  $\Lambda(8)$  and the Kaplan-Meier estimator of  $S(8) = P(T > 8)$ . (15 pts)

- (b) Find the variance estimates for the Nelson-Aalen and Kaplan-Meier estimators you got in (a). (15 pts)
- (c) Construct a 95% confidence interval for  $S(8)$  based on the Nelson-Aalen and Kaplan-Meier estimators you got in (a). (10 pts)
4. (20 pts) In a clinical trial, we want to compare a new treatment to the standard one on time (in months) to tumor recurrence for patients with certain type of cancer. Assume that enough patients will be available at the **beginning** of the trial and half of them will be randomly assigned to each treatment. It is known that the average tumor recurrence time is 2 month for the standard treatment; while for the new treatment, it is expected that it will extend the average tumor recurrence time to 4 month. The trial will last 10 months. Suppose that the tumor recurrence time for each treatment group follows an exponential distribution and the censoring time is uniformly distributed on the interval  $[0, 10]$ . Do the following:
- (a) Find the probability for each treatment that a patient will experience the tumor recurrence before the end of the trial. (10 pts)
- (b) Suppose we will use the log-rank test to test the difference in time to tumor recurrence for these two treatments with significance level  $\alpha = 0.05$  and 99% power to detect the expected difference. How many patients should be included in the study? (10 pts)