

SUPPORTING INFORMATION

Additional supporting information may be found online in the supporting information tab for this article.

List of information available

Table 1-SI: Perfusate composition.

Figure 1-SI: study recruitment flowchart

Figure 2-SI: vascular flows during NMP: (A) hepatic artery and (B) portal vein.

Figure 3-SI: IL-6 (A) and IL-10 (B) perfusate concentration during NMP.

Figure 4-SI: electron micrographs of liver samples after cold ischemia storage; (A) swollen mitochondria in hepatocytes from group CS; (B) magnification of the previous panel; (C, D) swollen mitochondria in hepatocytes from group NMP.

Figure 5-SI: electron micrographs of CS liver samples after cold ischemia storage; (A, B) cytoplasmic lipid droplets; (C, D) biliary poles with dense bodies.

Figure 6-SI: electron micrographs of liver samples after cold ischemia storage showing reduced Disse spaces (A: CS liver; B: NMP liver) and non-activated Kupffer cells (C: CS liver; D: NMP liver).

Figure 7-SI: electron micrographs of CS liver samples 2 hours after implantation; (A, B) mitochondrial swelling; (C, D) lipid droplets.

Figure 8-SI: electron micrographs of CS liver samples 2 hours after implantation: (A) numerous dense bodies in the biliary pole; (B) damaged vessel; (C) activated Kupffer cell; (D) infiltrating neutrophil.

Figure 9-SI: electron micrographs of NMP liver samples 2 hours after implantation: (A, B) mitochondria of normal size; (C, D) reduced lipid droplets

Table 1-SI. Perfusate composition.

COMPONENTS	VOLUME
ABO-compatible red blood cells concentrate	750 mL
Gelofusine® (Braun, Melsungen, Germany)	1000 mL
Human Albumin 200 g/L (Albital®, Kedrion, Barga (LU), Italy)	100 mL
Parenteral nutrition (Plusflex®, Braun, Melsungen, Germany)	10 mL
Multivitamins intravenous (IV) solution (Cernevit®, Italy)	10 mL
Metronidazole IV solution 5mg/mL (Baxter, Rome, Italy)	40 mL
Cefazolin 1000 mg (Teva Italia, Milan, Italy)	10 mL
Fluconazole 100 mg (Fresenius Kabi Italia, Verona, Italy)	50 mL
Fast acting Insulin 100 UI/ml (Actrapid®, Denmark)	20 mL
Calcium Gluconate IV solution 10% (Galenica Senese)	40 mL
Sterile water	50 mL
Sodium chlorate 0.9% solution (Fresenius Kabi Italia)	150 mL
Sodium bicarbonate 8.4% solution (Monico, Venice, Italy)	30 mL
Heparin 25,000 units (Epsoclar®, Hospira Italy, Naples, Italy)	5 mL
Total volume	2265 mL

Figure 1-SI. Study flowchart

CEFEMA Flowchart Sep. 2016




Figure 2-SI. Vascular flows during NMP: (A) hepatic artery and (B) portal vein.




Figure 3-SI. IL-6 (A) and IL-10 (B) perfusate concentration during NMP.




Figure 4-SI. Electron micrographs of liver samples after cold ischemia storage. **A)** Swollen mitochondria in hepatocytes from group CS; **B)** Magnification of the previous panel; **C, D)** Swollen mitochondria in hepatocytes from group NMP.




Figure 5-SI. Electron micrographs of CS liver samples after cold ischemia storage; (A, B) Cytoplasmic lipid droplets; (C, D) biliary poles with dense bodies.




Figure 6-SI. Electron micrographs of liver samples after cold ischemia storage showing reduced Disse spaces (A: CS liver; B: NMP liver) and non-activated Kupffer cells (C: CS liver; D: NMP liver).




Figure 7-SI. Electron micrographs of CS liver samples 2 hours after implantation; (A, B) mitochondrial swelling; (C, D) lipid droplets.







Figure 8-SI. Electron micrographs of CS liver samples 2 hours after implantation: (A) numerous dense bodies in the biliary pole; (B) damaged vessel; (C) activated Kupffer cell; (D) infiltrating neutrophil.




Figure 9-SI. Electron micrographs of NMP liver samples 2 hours after implantation.

A,B) mitochondria of normal size; **C,D)** reduced lipid droplets.

