

Supplementary Table 1. Clinical Outcomes for the Modified Intent-to-Treat Analysis Cohort*

	HMP-O ₂ N=65	SCS N=74	p-value
PRIMARY AND SECONDARY END-POINTS			
EAD, N (%)	8 (5.8)	12 (16.2)	0.51
Patient Survival (%)			0.11
3 months	96.9	98.7	
6 months	96.9	97.3	
12 months	90.8	97.3	
Graft Survival (%)			0.61
3 months	95.4	96.0	
6 months	95.4	93.2	
12 months	89.2	90.5	
Death Censored Graft Failure			0.34
3 months	3.1	4.1	
6 months	3.1	5.4	
12 months	3.1	8.2	
PNF, N (%)	1 (1.5)	3 (4.1)	0.38
Liver-Related Graft Failure, N (%)	2 (3.1)	5 (6.8)	0.32
Re-transplanted, N (%)	1 (1.5)	4 (5.4)	
Liver-Related Death, N (%)	1 (1.5)	1 (1.4)	
Biliary Complication, N (%)	8 (12.3)	18 (24.3)	0.15
Leak [§]	1 (1.5)	3 (4.1)	
Anastomotic Stricture ^{§§}	7 (10.8)	11 (14.9)	
Non-Anastomotic Stricture ^{§§§}	0 (0.0)	4 (5.4)	
Vascular Complications,[§] N (%)	1 (1.5)	2 (2.7)	0.89
Hepatic Artery Thrombosis	1 (1.5)	1 (1.4)	
Portal Vein Thrombosis	0 (0.0)	1 (1.4)	
Acute Kidney Injury within 7 days, N (%)	14 (21.5)	16 (21.6)	0.99
EXPLORATORY END-POINTS			
Hospital LOS[†], median (IQR)	9.0 (6.0, 14.0)	10.0 (6.0, 17.0)	0.38
ICU LOS[†], days, median (IQR)	3.0 (2.0, 4.0)	3.0 (2.0, 5.0)	0.53
POST-HOC ANALYSIS END-POINTS			
Severe Post-Operative Bleeding[*]	3 (4.6)	9 (12.2)	0.12
Moderate to Severe Ischemic Preservation Injury[*]	5 (7.7)	11 (14.9)	0.19
Acute Cellular Rejection, N (%)[*]	6 (9.2)	6 (8.1)	0.81
Incisional Hernia, N (%)[*]	0 (0.0)	4 (5.4)	0.12

*Includes patients planned for HMP-O₂ but receiving SCS, Excludes patients with major protocol deviations but includes patients with exclusion due to clinical parameters

§Defined as CT or drainage catheter evidence of bile leak in absence of hepatic artery thrombosis.

§§Defined as anastomotic bile duct narrowing with ERCP evidence of biliary ductal dilation requiring endoscopic or surgical intervention

§§§Defined as non-anastomotic/ intrahepatic narrowing with ERCP evidence of biliary ductal dilation or biliary casts

§Defined as portal vein thrombosis and/or hepatic artery thrombosis

†Defined as time from transplant to hospital discharge

EAD, early allograft dysfunction; HMP-O₂, hypothermic oxygenated machine perfusion; ICU, intensive care unit; IQR, interquartile range; LOS, length of stay; PNF, primary non-function; SCS, static cold storage.