

Supplementary Table 2. Biliary Complications in ECD Cohort†

	HMP-O ₂ N=15*	SCS N=21#	p-value
All Biliary Complications	2 (13.3)	8 (38.1)	0.14
DCD	2 (13.3)	3 (14.3)	
Age >65	0 (0.0)	5 (23.8)	
Steatosis >30%	0 (0.0)	0 (0.0)	
Biliary Leak	1 (6.7)	2 (9.5)	1.00
DCD	1 (6.7)	0 (0.0)	
Age >65	0 (0.0)	2 (9.5)	
Steatosis >30%	0 (0.0)	0 (0.0)	
Anastomotic Biliary Stricture	1 (6.7)	2 (9.5)	1.00
DCD	1 (6.7)	2 (9.5)	
Age >65	0 (0.0)	0 (0.0)	
Steatosis >30%	0 (0.0)	0 (0.0)	
Non-anastomotic Biliary Stricture	0 (0.0)	4 (19.1)	0.12
DCD	0 (0.0)	1 (4.8)	
Age >65	0 (0.0)	3 (14.3)	
Steatosis >30%	0 (0.0)	0 (0.0)	

†Results are presented as N (% of ECD recipients in treatment arm)

*HMP-O₂ ECD recipients are inclusive of 5 DCD recipients, 8 Advanced Age Recipients (>age 65), and 2 recipients with macrovesicular steatosis >30%.

#SCS ECD recipients are inclusive of 10 DCD recipients, 9 Advanced Age Recipients (>age 65), and 2 recipients with macrovesicular steatosis >30%.

DCD, donation after cardiac death; ECD, extended criteria donor; HMP-O₂, hypothermic oxygenated machine perfusion; SCS, static cold storage;