

Supplementary Table 3. Adverse events reported in per-protocol analysis cohort

	HMP-O₂ N=119 AEs	SCS N=176 AEs
POST-TRANSPLANT AEs AND SAEs		
All Adverse Events[#]	119 (100)	176 (100)
Mild AE	42 (35.3)	55 (31.3)
Moderate AE	62 (52.1)	84 (47.7)
Severe AE	15 (12.6)	37 (27.2)
Adverse Events ≤30 days	74 (62.2)	114 (64.8)
Adverse Events >30 days	45 (37.8)	62 (35.2)
Serious Adverse Event (SAE)	46 (38.7)	78 (44.3)
Infection-Related AEs[#]	54 (45.4)	57 (32.4)
Mild Infection-Related AE	22 (18.5)	30 (17.0)
Moderate Infection-Related AE	25 (21.0)	25 (14.2)
Severe Infection-Related AE	7 (5.9)	2 (1.1)
Non-Infection Related AEs[#]	65 (54.6)	122 (69.3)
Mild Noninfectious AE	20 (16.8)	29 (16.4)
Moderate Noninfectious AE	37 (31.1)	59 (33.5)
Severe Noninfectious AE	8 (6.7)	35 (19.9)
Non-Infection Related SAEs	28 (23.5)	64 (36.4)
POST-TRANSPLANT AEs BY CATEGORY		
Vascular - HAT[§]	1 (0.8)	1 (0.6)
Vascular - PVT[§]	0 (0.0)	1 (0.6)
Infection	54 (45.4)	57 (32.4)
≤30 Days	20 (16.8)	27 (15.3)
>30 Days	34 (28.6)	30 (17.0)
Ischemic Preservation Injury	3 (2.5)	10 (5.7)
Moderate	3 (2.5)	6 (3.4)
Severe	0 (0.0)	4 (2.3)
Early Allograft Dysfunction	7 (5.9)	12 (6.8)
Moderate-Severe Bleeding	10 (8.4)	19 (10.8)
Post-Op Return to OR [§]	6 (5.0)	8 (4.5)
Post-Op Transfusion	3 (2.5)	8 (4.5)
GI or Oropharyngeal Bleed	1 (0.8)	3 (1.7)
Biliary Complications	8 (6.7)	19 (10.8)
Bile Leak	1 (0.8)	3 (1.7)
Anastomotic Stricture	7 (5.9)	12 (6.8)
Non-Anastomotic Stricture	0 (0.0)	4 (2.3)
Acute Kidney Injury <7 days	14 (11.8)	17 (9.7)
Primary Nonfunction	0 (0.0)	3 (1.7)
Graft Failure	1 (0.8)	5 (2.8)
Acute Cellular Rejection	6 (5.0)	7 (4.0)
Incisional Hernia	0 (0.0)	4 (2.3)
Other	15 (12.6)	21 (11.9)
Neurologic	3 (2.5)	3 (1.7)
Cardiac	2 (1.7)	3 (1.7)
Pulmonary	2 (1.7)	3 (1.7)
Gastrointestinal	1 (0.8)	4 (2.3)
Wound Complication	2 (1.7)	0 (0.0)
Endocrine	0 (0.0)	2 (1.1)
Skin Cancer	1 (0.8)	0 (0.0)
Thromboembolic	4 (3.4)	4 (2.3)
Other	0 (0.0)	2 (1.1)

*All results reported as n/N(%Total Adverse Events for the treatment group)

[†]Defined as acute kidney injury by RIFLE criteria within 7 days of transplant

[#]Mild, Moderate and Severe complications, as defined by study site and sponsor definition; incidence calculated as discrete/total events

HAT, hepatic artery thrombosis; HMP-O₂, hypothermic oxygenated machine perfusion; OR, operating room; PVT, portal vein thrombosis; SCS, Static Cold Storage