

Supplementary Table 2 Demographics, Intraoperative, and Preservation Characteristics

Characteristics	All patients (n = 46)	SCS (n = 23)	HOPE (n = 23)	<i>P</i>
Donor characteristics				
Age, y, median (IQR)	72 (59–78)	71 (59–78)	73 (60–78)	0.875
BMI, kg/m ² , median (IQR)	28 (25–31)	28 (24–31)	28 (26–31)	0.700
Sex, female: male, n (%)	21 (46%): 25 (54%)	10 (44%): 13 (56%)	11 (48%): 12 (52%)	0.767
ET-DRI Score*	2.050 (1.878–2.218)	1.960 (1.830–2.240)	2.080 (1.910–2.210)	0.462
Cause of death, n (%)				
Cerebrovascular accident	30 (65%)	15 (65%)	15 (65%)	>0.999
Anoxia	10 (22%)	3 (13%)	7 (31%)	0.284
Trauma	6 (13%)	5 (22%)	1 (4%)	0.187
Donor Resuscitation, yes, n (%)	14 (31%)	6 (26%)	8 (35%)	0.749
Graft selection ECD criteria, n (%)†				
Donor age ≥65	31 (67%)	14 (61%)	17 (74%)	0.530
Donor ICU ≥7 days	13 (28%)	5 (22%)	8 (35%)	0.514
Donor BMI >30 kg/m ²	13 (28%)	7 (30%)	6 (26%)	>0.999
Macro/mixed steatosis >40%	10 (22%)	5 (22%)	5 (22%)	>0.999
Serum sodium >165 mmol/L	3 (7%)	2 (9%)	1 (4%)	>0.999
Serum AST/ALT >3× of normal	14 (30%)	8 (35%)	6 (26%)	0.749
Serum bilirubin >2 mg/dL	3 (7%)	1 (4%)	2 (9%)	>0.999
No. of ECD criteria reached, n (%)				
1 Of 7	13 (28%)	7 (30%)	6 (26%)	>0.999
2–3 Of 7	26 (57%)	14 (61%)	12 (52%)	0.767
>3 Of 7	7 (15%)	2 (9%)	5 (22%)	0.414
Microsteatosis, n (%)‡				
None to mild (0–29%)	32 (70%)	16 (70%)	16 (70%)	0.723
Moderate to	10 (22%)	6 (26%)	4 (17%)	0.723

severe ($\geq 30\%$)				
Macrosteatosis, n (%) [‡]				
None to mild (0–29%)	39 (85%)	21 (91%)	18 (78%)	0.598
Moderate to severe ($\geq 30\%$)	3 (7%)	1 (4%)	2 (9%)	0.598
Graft weight, g, median (IQR)	1643 (1379–1980)	1619 (1346–1984)	1643 (1342–1878)	0.943
Recipient characteristics				
Age, y, median (IQR)	62 (55–65)	63 (56–67)	60 (52–64)	0.262
BMI, kg/m ² , median (IQR)	28 (25–31)	28 (26–34)	28 (23–31)	0.443
Sex, female: male, n (%)	8 (17%): 38 (83%)	3 (13%): 20 (87%)	5 (22%): 18 (78%)	0.699
labMELD, median (IQR)	15 (9–25)	17 (8–25)	13 (9–18)	0.361
BAR Score, median (IQR) [§]	6 (4–10)	7 (4–11)	5 (3–8)	0.178
Listing indication, n (%)				
Alcoholic cirrhosis	11 (24%)	7 (31%)	4 (17%)	0.491
Viral hepatitis	3 (7%)	0 (0%)	3 (13%)	0.233
HCC in cirrhosis	20 (43%)	9 (39%)	11 (48%)	0.767
PSC/PBC	2 (4%)	1 (4%)	1 (4%)	>0.999
Other	10 (22%)	6 (26%)	4 (17%)	0.722
Pre-transplant abdominal surgery, n (%)	9 (20%)	2 (9%)	7 (30%)	0.135
eGFR at admission mL/min/1.73m ² , median (IQR)	89 (56–98)	79 (54–97)	91 (60–100)	0.473
Platelet counts at admission, nL, median (IQR)	98 (61–138)	98 (58–113)	97 (66–139)	0.866
Intraoperative and perfusion characteristics				
Total cold preservation time, min, median (IQR)	495 (465–593)	502 (467–584)	495 (447–612)	0.823
Cold preservation time before	375 (311–465)	NA	375 (311–465)	NA

HOPE, min, median (IQR)				
Machine perfusion time, min, median (IQR)	145 (101–203)	NA	145 (101–203)	NA
Warm ischemic, min, median (IQR)	43 (37–52)	45 (39–52)	39 (35–54)	0.275
Duration of surgery, min, median (IQR)	403 (324–474)	420 (308–495)	395 (329–452)	0.489
Intraoperative RBC, U, median (IQR)	4 (2–7)	4 (2–7)	3 (2–6)	0.327
Intraoperative FFP, U, median (IQR)	16 (10–24)	16 (10–18)	18 (9–26)	0.823

*Based on Braat et al.²⁴.

†Based on Czigany et al and the definitions of ECD criteria liver allografts in DBD donors as recommended by the Germany Medical Chamber (Bundesärztekammer)¹¹.

‡Based on McCormack et al.²²

§Based on Dutkowski et al.²⁵

||Based on the CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration) equation¹⁶.

BAR indicates balance of risk; BMI, body mass index; CKD-EPI, Chronic Kidney Disease Epidemiology Collaboration; FFP, fresh frozen plasma; HCC, hepatocellular carcinoma; ICU, intensive care unit; NA, not applicable; PBC, primary biliary cholangitis; PSC, primary sclerosing cholangitis; RBC, packed red blood cells.