

Lab Book

Brad Day

2025-05-19

Table of contents

Preface	3
1 Reading	4
1.1 Summary of “Broad Iron Lines in Active Galactic Nuclei” [1]	4
2 Initial Gradus Testing	8
3 Parameter Space Investigations	14
3.1 Ruling out α_{22} and α_{52}	14
3.2 Exploring the 5-dimensional parameter space	14
3.2.1 Line profiles	15
3.2.2 Redshift Image Renders	16
4 Fitting Spectral Lines	19
References	20

Preface

This book details my research during Summer 2026. Many thanks to Dr. Andrew Young and Mr Darius Michienzi for supervising this project.

This is a Quarto book.

To learn more about Quarto books visit <https://quarto.org/docs/books>.

1 Reading

1.1 Summary of “Broad Iron Lines in Active Galactic Nuclei” [1]

Fluorescent iron line, $\text{Fe-K}\alpha$, in the X-ray band at $6.4 - 6.9 \text{ keV}$ is the strongest narrow line emitted by an accretion disk around a black hole. Inverse Compton scattering of the soft photons (lower energy than the reactant particles) in a hot corona is the most likely candidate for the production of these X-rays. The emitted line is broadened and skewed due to the Doppler effect and gravitational redshift, and is often called the broad iron line due to being broadened to a FWHM of $5 \times 10^4 \text{ km s}^{-1}$.




Figure 1.1: The profile of the broad iron line, image credit to Fabian (2000) [1].

Soft Comptonisation (multiple inverse Compton scattering by hot thermal electrons) occurs giving a power law X-ray spectrum. Flares from active or flaring regions irradiate the accretion disk forming a “reflection” component in the spectrum.




Figure 1.2: Reflection spectrum from a Monte Carlo simulation of an illuminated slab. The dashed line is the incident continuum and the solid line shows the reflected spectrum. Image credit Reynolds (1996) [1]

The iron line is produced when one of the two K-shell electrons is ejected following absorption of an X-ray, the threshold for neutral iron is 7.1 keV. The resultant state can decay through the dropping of an L-shell electron into the K-shell, releasing 6.4 keV as either an emission line photon (34%) or an Auger electron (66%). Ionised iron has a photoelectric threshold and the $K\alpha$ of higher energy. The strength of the iron line is measured in terms of its equivalent width.

i Equivalent Width

The equivalent width of a spectral line is the width of the rectangle which has the same area as that between the line and the continuum level.




Figure 1.3: A diagram indicating the equivalent width. The width of the rectangle is the equivalent width.

The equivalent width is a function of the geometry of the accretion disk, the elemental abundances, the inclination angle, and the ionisation state.

! Key Points

- **X-ray Continuum:** The hard X-ray continuum in AGNs and GBHCs is thought to be from active/flaring regions in the corona. Thermal electrons multiply inverse Compton scatter optical and UV photons from the disk to X-ray energies. A hard X-ray power law results and irradiates the accretion disk to produce a reflection component.
- **Reflection Component:** The reflection component causes the spectrum to flatten above 10 keV as Compton recoil reduces the backscattered flux and results in a strong fluorescent iron line at ~ 6.4 keV. The energy and strength of this line depends on iron abundance, inclination of the disk, and its ionisation state. A strong iron absorption edge may also be present from an ionised disk.
- **Iron Line Profile:** Doppler shifts, relativistic broadening and gravitational redshifting give the broad and skewed line profile. As the line comes from the innermost regions, these effects are very pronounced and may be used to determine the black hole spin and inclination angle of the accretion disk. In most Seyfert 1 galaxies, the disk is inclined at around 30° , consistent with the standard model.
- **Observations:** The iron line was first clearly detected by Ginga and a line profile subsequently resolved by ASCA confirmed the broad and skewed shape expected from an accretion disk around a Schwarzschild black hole. RXTE has been used to study the line and continuum variability on shorter timescales, with reduced energy resolution.
- **Black Hole Spin:** The radius of the ISCO decreases with spin. Since the line profile is sensitive to the innermost radius of fluorescent emission this may be used to estimate the black hole, with some astrophysical assumptions. With present time-averaged observations such measurements may be ambiguous as alternative models with different values for the spin may produce identical line profiles.
- **Alternative Models for the Production of the Broad Iron Line:** Alternative models that do not require an accretion disk appear to fail, in particular the line width cannot be entirely due to Comptonisation. Hybrid models in which the Comptonisation and Doppler/gravitational effects produce the profile are heavily constrained.
- **Variability:** Rapid X-ray continuum variability is observed in most AGNs, and the iron line is expected to vary in response to this with short time lag. Whilst these timescales are too short to be probed presently, significant and complex iron line variability has been observed. Often the line flux is seen to remain constant whilst the continuum changes, and there appears to be an anticorrelation between the reflected fraction and the equivalent width. In another study the reflected fraction and the photon index of the power law are correlated for individual objects and between different ones. These need to be explained as they appear contrary to our simple model of reflection. Flux correlated changes in the ionisation state of the disk may explain some of these facts.
- **Reverberation Mapping:** The rapid variability is associated with activation of new flares in the corona above the accretion disk. X-ray reverberation mapping uses observations of the iron line response to sudden changes in the continuum to study the accretion disk and black hole. This may be used to determine the geometry of the X-ray emission and the black hole spin and mass.
- **Other Classes of Object** Iron lines are observed in quasars, LLAGNs, radio-loud AGNs, and GBHCs. Quasars have the iron line decrease with increasing luminosity which may be due to the more luminous sources accreting closer to the Eddington limit being more highly ionised. Observations in LLAGNs may determine whether the accretion with low rates is an advection-dominated accretion flow or a thin disk. Weak lines have been observed suggesting that their low accretion rate flows are thin disks rather than optically thick ADAFs. The broad iron lines and reflection humps are weak or absent perhaps because the reflection signature is swamped by a beamed continuum. GBHCs show smeared absorption and emission features in the reflection spectra, and there is a debate as to the precise nature of the accretion flow within a few tens of Schwarzschild radii.

2 Initial Gradus Testing

The below details initial experimentation with `Gradus.jl`. The code block below renders an image of a black hole with a thick accretion disk. Colour denotes the redshift of a given region on the disk.

```
using Gradus
using Plots

# Function defining the cross section of the disk
function discCrossSection(x)
    centre = 8
    radius = 3

    if (x < centre - radius) || (radius + centre < x)
        zero(x)
    else
        r = x - centre
        0.5(sqrt(radius^2 - r^2) + (0.5sin(3x)))
    end
end

# Metric Parameters
M = 1.0      # Mass
a = 0.7      # Black hole spin
13 = 0.0
22 = 0.0
52 = 0.0
3 = 0.0
i = deg2rad(70) # Inclination

# Instantiating the metric
m = JohannsenMetric(M, a, 13, 22, 52, 3)

# Position of the observer
# (x[1], Distance, Inclination, x[4])
x = SVector(0.0, 1000.0, i, 0.0)

# Maximum affine time, parameterises the solution
# _max ~ 2*x[2]
_max = 2000.0

# Setting up impact parameters
= range(-10.0, 10.0, 100)
= range(-10.0, 10.0, 100)
```



```

# Getting an array of velocities from the impact parameters
vs = vec([map_impact_parameters(m, x, a, b) for a in , b in ])
xs = fill(x, size(vs))

# Generating solutions
sols = tracegeodesics(m, xs, vs, _max, save_on = false)

# Generating points and reshaping to be the same shape as the image
points = unpack_solution.(sols.u)
points = reshape(points, (100, 100))

# Redshift point function
redshift = ConstPointFunctions.redshift(m, x)
redshiftGeometry = redshift ConstPointFunctions.filter_intersected()

# Disc
d = ThickDisc(r -> discCrossSection(r))

# this function returns the impact parameter axes
, , image = rendergeodesics(
    m, x, d, _max, pf = redshiftGeometry,
    # image parameters
    image_width = 200, image_height = 150,
    # the "zoom" -- use the impact parameter axes
    lims = (-20, 20), lims = (-15, 15), verbose = false
)

heatmap(, , image, aspect_ratio = 1)

```




Figure 2.1: The first render produced.

Once familiarity was gained with the basics of `Gradus.jl` the line profile functionality was utilised through the following function.

```
function computeLineProfile(m, i, height, bins = range(0.0, 1.5, 180))
    # Position of the observer
    # (x[1], Distance, Inclination, x[4])
    x = SVector(0.0, 10000.0, i, 0.0)

    # Disk
    d = ThinDisc(0.0, Inf)

    # Setting up the model and profile
    model = LampPostModel(h = height)
    profile = emissivity_profile(m, d, model)

    # Computing the line profile
    bins, flux = lineprofile(m, x, d, profile; bins=bins, verbose=true)

    return bins, flux
end
```

This will allow for the investigation of the free parameters for the Johannsen metric; α_{13} , α_{22} , α_{52} , and ϵ_3 . A lamppost corona model is used with variable height h . An example of the effect of varying the height of this corona is shown below.





Figure 2.2: Initial variations of the height of the lamppost corona.

Following initial experimentation the two functions above were integrated together for efficient investigation. The two free parameters that will be the focus of this investigation are α_{13} and ϵ_3 . Examples of the effects of these parameters on the emission lines are shown below






Figure 2.3: Initial variations of α_{13} and ϵ_3

It can be seen that α_{13} has a very smooth, almost linear effect on the shape of the line, while ϵ_3 is much more chaotic, particularly for $\epsilon_3 > 25$.

3 Parameter Space Investigations

Code was set up to easily run several variations to the model which will allow for the investigation of the parameter space. A thin disk accretion model and a lamppost corona were used. It was set up such that for a given configuration of h , a , i , ϵ_3 , and α_{13} the line profile would be calculated and an image of the accretion disk produced and plotted together.

3.1 Ruling out α_{22} and α_{52}

Before exploring the 5D parameter space, tests were ran on both α_{22} and α_{52} such that their effects could be noted, should they be useful in the future.




Figure 3.1: The resultant line profile from varying α_{52} between 0 and 40. A value of 50 did not successfully compute suggesting an upper limit on this parameter.

Varying α_{52} showed near random, and near zero effect on the line and will thus be disregarded for this investigation. Varying α_{22} often failed to compute and will thus similarly be disregarded for this investigation.

3.2 Exploring the 5-dimensional parameter space

Each of the 5 parameters were varied individual, with the remaining four being held at default values of

Parameter	Value
α_{13}	0.0
ϵ_3	0.0
a	0.998
h	$10 R_G$
i	60°

3.2.1 Line profiles




Figure 3.2: The line profiles resultant from varying α_{13} between 0 and 50.




Figure 3.3: The line profiles resultant from varying ϵ_3 between 0 and 30. Values of 40 and 50 were tested however caused an error, there must be an upper bound on the allowed values of this parameter.




Figure 3.4: The line profiles resultant from varying the inclination between 5 and 85 degrees.




Figure 3.5: The line profiles resultant from varying h between 5 and 30 gravitational radii.

- Figure 3.7 shows that increasing the free parameter α_{13} sharpens the two peaks of the line, increases the maximum flux, and makes it more narrow.
- Figure 3.3 shows that increasing the free parameter ϵ_3 similarly increases the maximum flux, however the resultant profile is much more noisy. While α_{13} provided sharp peaks, ϵ_3 provided broader peaks,




Figure 3.6: The line profiles resultant from varying a between 5 and 95% of $M(= 1)$.

particularly in the shorter one.

- Figure 3.4 shows that increasing the inclination smoothes out the line profile and shifts the peak to higher energies. The line is broadened significantly.
- Figure 3.5 shows that increasing the corona height makes the line profile sharper, increases the maximum flux, and shifts the maximum to slightly lower energies.
- Figure 3.6 shows that the spin had the least prominent impact on the profile of the line, seeming to only slightly decrease the height of the peaks with increasing spin. This may be an effect of the small range of spins available. It may therefore be informative to investigate this further by increasing the mass, using different corona heights, or varying the inclination.

3.2.2 Redshift Image Renders

Alongside the line profiles, images of the accretion disk were rendered with a colourmap denoting the redshift of light from a given part of the disk. The corona height was not varied here as it did not impact the render.




Figure 3.7: The effect of α_{13} on the accretion disk render. The event horizon appears to grow with α_{13}




Figure 3.8: The effect of ϵ_3 on the redshift image. The shape of the disk does not change but the redshift profile does, particularly in the inner regions where the redshift tends to increase.




Figure 3.9: The effect of inclination on the redshift image. The redshift becomes more apparent as the inclination increases and the disk becomes edge on. This agrees with the observed line profile trend of energy increasing with inclination as the effects of doppler shifting become more apparent with increasing energy.




Figure 3.10: The effect of black hole spin on the redshift image. The event horizon appears to shrink.

4 Fitting Spectral Lines

With the parameter space explored, it was useful to setup the architecture for fitting the model to spectral data. This was done using `SpectralFitting.jl` and defining a custom model using Gradus which fits the following parameters:

Parameter	Meaning
K	Normalisation
h	Corona Height
E	Line Energy
R_{in}	Inner Radius
R_{out}	Outer Radius
θ	Inclination
a	Spin
13	α_{13} Parameter
3	ϵ_3 Parameter

a

References

- [1] A. C. Fabian *et al.*, Publications of the Astronomical Society of the Pacific **112**, 1145 (2000).