

Lab Book

Bradley Day

2025-05-19

Table of contents

Preface	3
I Summer Reading	4
1 Work Done Over the Summer Break	5
II Teaching Block 1	6
2 Week 1	7
2.1 2026/09/21	7
2.2 2026/09/22	7
2.2.1 Meeting Minutes	7
2.3 2026/09/23	8
2.3.1 Reading - K. Nandra (2007) [1]	8
2.4 2026/09/26	9
2.4.1 Reading - Fabian et al. (2002) [2]	9
III Teaching Block 2	22
References	33

Preface

This book details my research during the final year of my degree at the University of Bristol. Thank you to Professor Andrew Young for supervising and my project partner Jacob Taylor for his work.

This is a Quarto book.

To learn more about Quarto books visit <https://quarto.org/docs/books>.

Part I

Summer Reading

1 Work Done Over the Summer Break

Some reading and meetings occurred over the summer break prior to the start of the project, here I detail notes from meetings and will note key points from reading.

This project will be based on emission line fitting to the Iron $K\alpha$ emission line seen when looking at black holes. The focus will be on Active Galactic Nuclei (AGN).

Part II

Teaching Block 1

2 Week 1

2.1 2026/09/21

Lab book set up and initialised as a Github page. Meeting with supervisor arranged for 2026/09/22.

2.2 2026/09/22

Plans/questions for meeting:

- Confirm that Quarto is allowed for writing lab book
 - Yes
- Get connected to solva
 - This will need to be after Rhys has set things up
 - Note 2026-09-23 have connected to solva, however will wait until Rhys has given an induction to use
- Make sure GitHub set up properly so we can commit/push to it
- Confirm scope of project and form initial plan with Jacob
 - Fitting models?
 - * XSPEC?
 - * Julia (SpectralFitting.jl)?
 - * Other?
 - Data Reduction
 - * Using XMM-Newton data so will need to become familiar with [XMM SAS](#)
 - * Are we using this on solva?
 - * Note to self: see project plan on Github for more resources

2.2.1 Meeting Minutes

- XMM is not a survey telescope so even random selections will usually be interesting objects
- Andy will add a link to Nandra's paper on the Github repo for bright sources
 - Read this paper
- Browse paper(s) to find interesting sources
 - NGC4151?
 - Make sure to note when and how long observations were taken
 - HEASARC archive should let us search for particular objects
 - Download data and go through the ABC guide
- Before next meeting:
 - Look into candiadate AGN

- * Look at Tartarus database?
- Read papers provided by Andy
- Look through XMM SAS ABC guide

2.3 2026/09/23

2.3.1 Reading - K. Nandra (2007) [1]

- Covers 37 observations of 26 Seyfert galaxies from XMM-Newton
 - All show Iron K emission, which makes this paper useful for candidate selection for our research
 - Around 45% of the data was best fit with a relativistically blurred reflection model
 - * Some objects may not show the broadening?
- General background information
 - Iron line emission broadening predicted in Fabian et al (1989)
 - This was alongside the discovery that the emission likely comes from the accretion disk
 - Useful reads?
 - * Fabian (1989)
 - * George, Nandra & Fabian (1990)
 - * Tanaka et al. (1995)
- Data Reduction
 - They give a lot of detail on their data reduction which may be useful further down the line
 - Source taken from a circle with radius 700 detector pixels
 - There are several windowing modes on the CCD which will be important to note
- Model
 - They found that the minimum set of models was:
 - * Power law continuum (modelling the corona)
 - * Reflection model (e.g. XILLVER)
 - * Can use relativistic blurring on this however not necessarily needed
 - * Ionized absorber model
 - The models used are covered in depth, including how they selected if and how to apply the absorber models
 - * This will be useful later down the line to give some direction when fitting
 - They modelled the emission using a broad and narrow component, this may be useful to consider?
- Potential candidate list (from Figure 3 of Nandra et al.)
 - NGC3516(2)
 - * Very broad emission centred around the Fe K α line
 - NGC3783(2)
 - * Quite a clean spectrum with low noise/small errors
 - NGC4151(2) or NGC4151(3)
 - * Clean spectrum with broad emission

2.4 2026/09/26

2.4.1 Reading - Fabian et al. (2002) [2]

2.4.1.1 Background

- Presents preliminary results from a 325 ks observation of MCG-6-30-15
 - Data taken from both the EPIC MOS cameras and the EPIC pn camera in small window mode with a medium filter
- Very clear detection of a strong, broad, and skewed iron line
- Spectrum characterised well by a steep emissivity profile within $6r_g$ and a flatter profile beyond
- Inner radius of emission appears to lie around $2r_g$
 - This was measured from the fit, where the inner radius of the disc was a fitting parameter
 - The measurements are dependent on an assumed power-law continuum and lack of complex absorption above 2.5 keV
- ASCA observation showed emission from an accretion disc extending from $6 - 40r_g$ with an inclination of around 30°
 - Measured from Fe- $k\alpha$ line
- The red wing of the line extends below 4 keV which can be explained by the disc extending within $6 r_g$
 - Why is this??
 - This implies rapidly spinning black hole (as ISCO shrinks with a_*)

2.4.1.2 Fitting

- Several models were used to fit the data, including a phenomenological model (parameterising the iron line with power law, Compton reflection, a narrow Gaussian at 6.4 keV and a broad Gaussian which was free)
 - This provided a good overall fit with $\chi^2 = 984.1$ for 1034 degrees of freedom
 - * How are the number of dof determined?
 - This was improved by allowing the narrow Gaussian to have some width, giving an “excellent” fit
- Second model used the Laor (1991) emission line profile
 - Fits were insensitive to the outer radius of the disc, which was thus fixed at $400r_g$
 - Free parameters were R_{in} , the inclination, and the emissivity as a function of radius (parametrised as R^{-q})
 - Almost comparable χ^2 to the first model with two broad Gaussians
 - * This falls short as with a simple power law emissivity, the red wing and narrow core of the line cannot be modelled simultaneously
- Further improvements were made including
 - Addition of an iron K-edge
 - * The K-edge is a sudden increase in x-ray absorption [FIND CITATION FOR THIS, [wiki for now](#)]
 - * This significantly improved the fit
 - * The K-edge however must be explained physically, several attempts were made
 1. Neutral iron along the line of sight
 - This would require strong iron-L absorption, which is not present

2. Neutral iron outside the line of sight
 - Reflection off of distant material which would require strong reflection
 - This is inconsistent with the weak, narrow 6.4 keV iron line
 3. Thick material partially covering the line of sight
 - This gave a comparable fit to the edge
 4. Sako et al. (2002) suggested significant absorption due to inner-shell transitions of ionised iron
 - Explored using an absorption line model, giving a good fit
 5. The complexity around 7.0 keV is emission
 - Recombination emission by H-like iron
- * It was not possible to determine whether the complexity was due to absorption or emission and both were thus considered
 - Use of a broken power law
 - * This is not physically self-consistent
 - Ignores Fe $K\beta$ at 7.05 keV
 - The high EW is not consistent with $R_{\text{refl}} = 1$, $i = 30^\circ$, solar abundances and neutral
 - High abundance was investigated using $3\times$ the solar abundances which improved the fit significantly
 - Spectral variability was investigated by taking the ratio of the spectra resulting from two 10 ks intervals from the maximum and minimum flux periods
 - Taking the ratio of the two spectra showed clear signs
 - The iron line core is stronger in the low-flux spectrum
 - The models assume that the effects of ionized absorption are negligible above 2.5 keV, correct for absorption rising from low-Z elements, but may not be true if e.g. Si, S are present

Part III

Teaching Block 2

References

- [1] K. Nandra *et al.*, Monthly Notices of the Royal Astronomical Society **382**, 194 (2007).
- [2] A. C. Fabian *et al.*, (2002).