



Self-Organization and Brain Function

Project URL: <http://brainhack.org/self-organization-and-brain-function>

J. P. Pfannmöller^{1*}, R. Mesquita², L.C.T. Herrera² and Daniela Dentico³

1 Introduction

Self-organization is a fundamental property of complex systems, describing the order spontaneously arising by the local interactions of the system components not mediated by top-down inputs. Though, self-organizing systems typically possess a large number of components and exhibit complex dynamics, their evolution is deterministic and governed by a small number of order parameters. This property is used here to model the self-organization of the ocular dominance columns of the striate cortex in patterns of neighboring stripes, which respond preferentially to inputs from the left or the right eye. Several models for self-organization with various degrees of complexity exist. We chose the most simple model due to its robustness and limited number of parameter.

2 Approach

The Swift-Hohenberg equation [1] was used to model the self-organization of the ocular dominance columns. There are two order parameters in this equation, the first one determines the spatial wavelength λ of the stripes and the second one the branchiness ϵ of the pattern. The algorithm used to generate the results has been modified from an open source script. The Swift-Hohenberg was solved by applying periodic boundary conditions after a Fourier transform to k space, which simplifies the computation of the solution to matrix operations:

for k = 1:N

```
q(k) = (2*(k-1)/N-1)*pi*N/L;
end
```

for k = 1:N

for l = 1:N

```
Q(k,l) = 1/(1-dt*(epsilon-(k0^2-(q(k)^2+q(l)^2))^2));
end
end.
```

3 Results

Figures (a), (b) and (c) show the temporal evolution of the solution to the Swift-Hohenberg equation for random initial conditions (a), constant ϵ and time increasing from (a) to (c). In (c), (d) and (e) three solutions with different ϵ are shown. The branchiness increases with ϵ from (c) to (e). The wavelength λ was set to the same value in all figures and the pattern in (d) is similar to the ocular dominance layers found in the visual cortex.

4 Conclusions

A simple model suffices to study basic properties of ocular dominance self-organization. Possibly, a combination with models for self-organization in neighboring cortical layers would allow to investigate higher organizational principles of the cortex [2], e.g. the coordination between ocular dominance, orientation, and cytochrome oxidase.

Availability of Supporting Data

More information about this project can be found at:

<http://brainhack.org/self-organization-and-brain-function>.

Further data and files supporting this project are hosted in the *GigaScience* repository

https://github.com/Brainhack-Proceedings-2015/Pfan_HBM_SOBF.

Competing interests

None

Author's contributions

JPP, RM, LCTH, and DD performed the project and wrote the report.

*Correspondence: pfannmoelj@uni-greifswald.de

¹Functional Imaging Unit, Center for Diagnostic Radiology, University Medicine Greifswald, Greifswald, Fleischmann Strasse 8, 17475, Germany
Full list of author information is available at the end of the article

Acknowledgements

The authors would like to thank the organizers and attendees of the 2015 OHBM Hackathon. This work was supported by generous donations from individuals to the Center for Investigating Healthy Minds, founded and led by Dr. Richard J. Davidson.

Author details

¹Functional Imaging Unit, Center for Diagnostic Radiology, University Medicine Greifswald, Greifswald, Fleischmann Strasse 8, 17475, Germany.

²Institute of Physics, University of Campinas, Campinas, Rua Sérgio Buarque de Holanda 777, 13083-859, Brazil. ³Waisman Center, University of Wisconsin, Madison, 1500 Highland Ave, WI 53705, USA.

References

1. Hohenberg, P.C., Swift, J.B.: Effects of additive noise at the onset of rayleigh-bénard convection. *Phys. Rev. A* **46**, 4773–4785 (1992).
doi:[10.1103/PhysRevA.46.4773](https://doi.org/10.1103/PhysRevA.46.4773)
2. Reichl, L., Heide, D., Löwel, S., Crowley, J.C., Kaschube, M., Wolf, F.: Coordinated optimization of visual cortical maps (i) symmetry-based analysis. *PLoS Comput Biol* **8**(11), 1002466 (2012).
doi:[10.1371/journal.pcbi.1002466](https://doi.org/10.1371/journal.pcbi.1002466)

