MEDICAL DISEASE KNOWLEDGE BASE

Respiratory Diseases

RAG-ready educational reference | Version date: 06 October 2026

Table of Contents

1. Asthma
2. Chronic obstructive pulmonary disease (COPD)
3. Pneumonia
4. Pulmonary tuberculosis
5. Obstructive sleep apnea
6. Pulmonary fibrosis

Respiratory Diseases

This document is an educational and information-retrieval reference for a hospital AI assistant. It is not a diagnostic tool and does not replace evaluation, diagnosis, treatment, or advice from a qualified healthcare professional. Symptoms can have many causes. Treatment decisions depend on age, pregnancy, comorbidities, severity, local epidemiology, available resources, and current professional guidelines.

Safety rule: An AI system must not claim to establish a definitive diagnosis, prescribe individualized medication or dosage, or guarantee cure. It should recommend professional assessment when symptoms are concerning and immediate emergency care for relevant red flags.

ICD note: ICD-11 terminology and codes should be verified in the current WHO ICD-11 Browser. Codes are not reproduced unless reliably verified for the specific concept; local coding systems and national adaptations may differ.

Standard disease-entry schema: Every disease entry should use the following fields: Disease Name; 1. Definition; 2. Medical Classification; 3. Causes; 4. Risk Factors; 5. Signs and Symptoms; 6. Disease Progression; 7. Differential Diagnosis; 8. Diagnostic Evaluation; 9. Diagnosis; 10. Treatment and Management; 11. Prevention; 12. Complications; 13. Prognosis; 14. When to Seek Medical Care; 15. Emergency Red Flags; 16. Patient Education; 17. Frequently Asked Questions; 18. Related Conditions; 19. Medical Terms; 20. Sources.

Retrieval fields: DISEASE, SYNONYMS, RELATED SYMPTOMS, RELATED CONDITIONS, RELATED TESTS, RELATED TREATMENTS, RELATED COMPLICATIONS. Symptom presence alone does not establish a diagnosis.

1. Asthma

DISEASE: Asthma

SYNONYMS: Asthma

RELATED SYMPTOMS: Variable wheeze, cough, chest tightness, and breathlessness; symptoms may fluctuate and be triggered by allergens, infection, exercise, or irritants.

RELATED TESTS: History, examination, and objective assessment of variable airflow limitation when feasible.

1. Definition

Asthma is a medical condition in the respiratory diseases category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Respiratory Diseases. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Variable wheeze, cough, chest tightness, and breathlessness; symptoms may fluctuate and be triggered by allergens, infection, exercise, or irritants. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History, examination, and objective assessment of variable airflow limitation when feasible. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Asthma should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Asthma? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Asthma; clinical terminology; patient terminology; suspected Asthma; symptoms associated with Asthma; diagnosis and management of Asthma.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

2. Chronic obstructive pulmonary disease (COPD)

DISEASE: Chronic obstructive pulmonary disease (COPD)

SYNONYMS: Chronic obstructive pulmonary disease (COPD)

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Chronic obstructive pulmonary disease (COPD) is a medical condition in the respiratory diseases category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Respiratory Diseases. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Chronic obstructive pulmonary disease (COPD) should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Chronic obstructive pulmonary disease (COPD)? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Chronic obstructive pulmonary disease (COPD); clinical terminology; patient terminology; suspected Chronic obstructive pulmonary disease (COPD); symptoms associated with Chronic obstructive pulmonary disease (COPD); diagnosis and management of Chronic obstructive pulmonary disease (COPD).

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

3. Pneumonia

DISEASE: Pneumonia

SYNONYMS: Pneumonia

RELATED SYMPTOMS: Cough, fever, breathlessness, chest discomfort, and fatigue; older adults may present with confusion or weakness.

RELATED TESTS: Clinical examination, oxygen assessment, and chest imaging or microbiology when indicated.

1. Definition

Pneumonia is a medical condition in the respiratory diseases category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Respiratory Diseases. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Cough, fever, breathlessness, chest discomfort, and fatigue; older adults may present with confusion or weakness. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include Clinical examination, oxygen assessment, and chest imaging or microbiology when indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Pneumonia should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Pneumonia? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Pneumonia; clinical terminology; patient terminology; suspected Pneumonia; symptoms associated with Pneumonia; diagnosis and management of Pneumonia.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

4. Pulmonary tuberculosis

DISEASE: Pulmonary tuberculosis

SYNONYMS: Pulmonary tuberculosis

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Pulmonary tuberculosis is a medical condition in the respiratory diseases category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Respiratory Diseases. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Pulmonary tuberculosis should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Pulmonary tuberculosis? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Pulmonary tuberculosis; clinical terminology; patient terminology; suspected Pulmonary tuberculosis; symptoms associated with Pulmonary tuberculosis; diagnosis and management of Pulmonary tuberculosis.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

5. Obstructive sleep apnea

DISEASE: Obstructive sleep apnea

SYNONYMS: Obstructive sleep apnea

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Obstructive sleep apnea is a medical condition in the respiratory diseases category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Respiratory Diseases. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Obstructive sleep apnea should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Obstructive sleep apnea? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Obstructive sleep apnea; clinical terminology; patient terminology; suspected Obstructive sleep apnea; symptoms associated with Obstructive sleep apnea; diagnosis and management of Obstructive sleep apnea.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

6. Pulmonary fibrosis

DISEASE: Pulmonary fibrosis

SYNONYMS: Pulmonary fibrosis

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Pulmonary fibrosis is a medical condition in the respiratory diseases category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Respiratory Diseases. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Pulmonary fibrosis should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Pulmonary fibrosis? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Pulmonary fibrosis; clinical terminology; patient terminology; suspected Pulmonary fibrosis; symptoms associated with Pulmonary fibrosis; diagnosis and management of Pulmonary fibrosis.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

Category Sources / References

World Health Organization (WHO), Health Topics: https://www.who.int/health-topics

World Health Organization, International Classification of Diseases (ICD): https://www.who.int/standards/classifications/classification-of-diseases

WHO ICD-11 Browser: https://icd.who.int/en

Centers for Disease Control and Prevention (CDC), official clinical and public-health guidance: https://www.cdc.gov/

National health authorities and recognized professional medical organizations should be consulted for current local guidance.

Glossary

Acute: developing over a short period. Chronic: persistent or recurrent over time. Differential diagnosis: comparison of plausible conditions causing a presentation. Diagnostic evaluation: history, examination, and investigations used to assess a patient. Prognosis: expected course and outcome at a population or disease level, not an individual guarantee. Red flag: a feature that may indicate serious disease and requires urgent assessment.

Master References

WHO. Health Topics. Updated information and fact sheets. https://www.who.int/health-topics

WHO. International Classification of Diseases, 11th Revision. https://icd.who.int/en

CDC. Official disease, emergency, prevention, and infection-control guidance. https://www.cdc.gov/

National health ministries/agencies and recognized specialty organizations: use current jurisdiction-specific guidelines.

This knowledge base is intended for educational and information-retrieval purposes. It does not replace evaluation, diagnosis, treatment, or advice from a qualified healthcare professional. Symptoms can have many different causes, and the information presented here cannot determine an individual's diagnosis. Emergency symptoms require immediate professional medical evaluation.