MEDICAL DISEASE KNOWLEDGE BASE

Pain, Symptoms, and Clinical Presentations

RAG-ready educational reference | Version date: 06 October 2026

Table of Contents

1. Headache
2. Migraine
3. Fever
4. Cough
5. Shortness of breath
6. Chest pain
7. Abdominal pain
8. Back pain
9. Neck pain
10. Joint pain
11. Muscle pain
12. Sore throat
13. Nausea
14. Vomiting
15. Diarrhea
16. Constipation
17. Dizziness
18. Fainting
19. Fatigue
20. Weakness
21. Swelling
22. Rash
23. Itching
24. Weight loss
25. Weight gain
26. Loss of appetite
27. Difficulty swallowing
28. Blood in stool
29. Blood in urine
30. Painful urination
31. Frequent urination
32. Excessive thirst
33. Excessive hunger
34. Vision changes
35. Hearing changes
36. Numbness
37. Tingling
38. Confusion
39. Seizures
40. Sleep problems
41. Anxiety
42. Depression symptoms

43. Nerve pain
44. Neuropathic pain
45. Chronic pain
46. Acute pain
47. Cancer-related pain
48. Post-surgical pain

Pain, Symptoms, and Clinical Presentations

This document is an educational and information-retrieval reference for a hospital AI assistant. It is not a diagnostic tool and does not replace evaluation, diagnosis, treatment, or advice from a qualified healthcare professional. Symptoms can have many causes. Treatment decisions depend on age, pregnancy, comorbidities, severity, local epidemiology, available resources, and current professional guidelines.

Safety rule: An AI system must not claim to establish a definitive diagnosis, prescribe individualized medication or dosage, or guarantee cure. It should recommend professional assessment when symptoms are concerning and immediate emergency care for relevant red flags.

ICD note: ICD-11 terminology and codes should be verified in the current WHO ICD-11 Browser. Codes are not reproduced unless reliably verified for the specific concept; local coding systems and national adaptations may differ.

Standard disease-entry schema: Every disease entry should use the following fields: Disease Name; 1. Definition; 2. Medical Classification; 3. Causes; 4. Risk Factors; 5. Signs and Symptoms; 6. Disease Progression; 7. Differential Diagnosis; 8. Diagnostic Evaluation; 9. Diagnosis; 10. Treatment and Management; 11. Prevention; 12. Complications; 13. Prognosis; 14. When to Seek Medical Care; 15. Emergency Red Flags; 16. Patient Education; 17. Frequently Asked Questions; 18. Related Conditions; 19. Medical Terms; 20. Sources.

Retrieval fields: DISEASE, SYNONYMS, RELATED SYMPTOMS, RELATED CONDITIONS, RELATED TESTS, RELATED TREATMENTS, RELATED COMPLICATIONS. Symptom presence alone does not establish a diagnosis.

1. Headache

DISEASE: Headache

SYNONYMS: Headache

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Headache is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Headache should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Headache? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Headache; clinical terminology; patient terminology; suspected Headache; symptoms associated with Headache; diagnosis and management of Headache.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

2. Migraine

DISEASE: Migraine

SYNONYMS: Migraine

RELATED SYMPTOMS: Recurrent headache, often throbbing, sometimes with nausea, light or sound sensitivity, and aura.

RELATED TESTS: Clinical history and examination; additional testing is guided by red flags or atypical features.

1. Definition

Migraine is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Recurrent headache, often throbbing, sometimes with nausea, light or sound sensitivity, and aura. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include Clinical history and examination; additional testing is guided by red flags or atypical features. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Migraine should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Migraine? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Migraine; clinical terminology; patient terminology; suspected Migraine; symptoms associated with Migraine; diagnosis and management of Migraine.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

3. Fever

DISEASE: Fever

SYNONYMS: Fever

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Fever is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Fever should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Fever? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Fever; clinical terminology; patient terminology; suspected Fever; symptoms associated with Fever; diagnosis and management of Fever.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

4. Cough

DISEASE: Cough

SYNONYMS: Cough

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Cough is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Cough should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Cough? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Cough; clinical terminology; patient terminology; suspected Cough; symptoms associated with Cough; diagnosis and management of Cough.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

5. Shortness of breath

DISEASE: Shortness of breath

SYNONYMS: Shortness of breath

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Shortness of breath is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Shortness of breath should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Shortness of breath? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Shortness of breath; clinical terminology; patient terminology; suspected Shortness of breath; symptoms associated with Shortness of breath; diagnosis and management of Shortness of breath.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

6. Chest pain

DISEASE: Chest pain

SYNONYMS: Chest pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Chest pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Chest pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Chest pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Chest pain; clinical terminology; patient terminology; suspected Chest pain; symptoms associated with Chest pain; diagnosis and management of Chest pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

7. Abdominal pain

DISEASE: Abdominal pain

SYNONYMS: Abdominal pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Abdominal pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Abdominal pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Abdominal pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Abdominal pain; clinical terminology; patient terminology; suspected Abdominal pain; symptoms associated with Abdominal pain; diagnosis and management of Abdominal pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

8. Back pain

DISEASE: Back pain

SYNONYMS: Back pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Back pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Back pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Back pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Back pain; clinical terminology; patient terminology; suspected Back pain; symptoms associated with Back pain; diagnosis and management of Back pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

9. Neck pain

DISEASE: Neck pain

SYNONYMS: Neck pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Neck pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Neck pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Neck pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Neck pain; clinical terminology; patient terminology; suspected Neck pain; symptoms associated with Neck pain; diagnosis and management of Neck pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

10. Joint pain

DISEASE: Joint pain

SYNONYMS: Joint pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Joint pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Joint pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Joint pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Joint pain; clinical terminology; patient terminology; suspected Joint pain; symptoms associated with Joint pain; diagnosis and management of Joint pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

11. Muscle pain

DISEASE: Muscle pain

SYNONYMS: Muscle pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Muscle pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Muscle pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Muscle pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Muscle pain; clinical terminology; patient terminology; suspected Muscle pain; symptoms associated with Muscle pain; diagnosis and management of Muscle pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

12. Sore throat

DISEASE: Sore throat

SYNONYMS: Sore throat

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Sore throat is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Sore throat should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Sore throat? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Sore throat; clinical terminology; patient terminology; suspected Sore throat; symptoms associated with Sore throat; diagnosis and management of Sore throat.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

13. Nausea

DISEASE: Nausea

SYNONYMS: Nausea

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Nausea is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Nausea should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Nausea? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Nausea; clinical terminology; patient terminology; suspected Nausea; symptoms associated with Nausea; diagnosis and management of Nausea.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

14. Vomiting

DISEASE: Vomiting

SYNONYMS: Vomiting

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Vomiting is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Vomiting should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Vomiting? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Vomiting; clinical terminology; patient terminology; suspected Vomiting; symptoms associated with Vomiting; diagnosis and management of Vomiting.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

15. Diarrhea

DISEASE: Diarrhea

SYNONYMS: Diarrhea

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Diarrhea is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Diarrhea should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Diarrhea? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Diarrhea; clinical terminology; patient terminology; suspected Diarrhea; symptoms associated with Diarrhea; diagnosis and management of Diarrhea.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

16. Constipation

DISEASE: Constipation

SYNONYMS: Constipation

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Constipation is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Constipation should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Constipation? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Constipation; clinical terminology; patient terminology; suspected Constipation; symptoms associated with Constipation; diagnosis and management of Constipation.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

17. Dizziness

DISEASE: Dizziness

SYNONYMS: Dizziness

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Dizziness is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Dizziness should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Dizziness? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Dizziness; clinical terminology; patient terminology; suspected Dizziness; symptoms associated with Dizziness; diagnosis and management of Dizziness.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

18. Fainting

DISEASE: Fainting

SYNONYMS: Fainting

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Fainting is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Fainting should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Fainting? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Fainting; clinical terminology; patient terminology; suspected Fainting; symptoms associated with Fainting; diagnosis and management of Fainting.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

19. Fatigue

DISEASE: Fatigue

SYNONYMS: Fatigue

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Fatigue is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Fatigue should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Fatigue? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Fatigue; clinical terminology; patient terminology; suspected Fatigue; symptoms associated with Fatigue; diagnosis and management of Fatigue.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

20. Weakness

DISEASE: Weakness

SYNONYMS: Weakness

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Weakness is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Weakness should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Weakness? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Weakness; clinical terminology; patient terminology; suspected Weakness; symptoms associated with Weakness; diagnosis and management of Weakness.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

21. Swelling

DISEASE: Swelling

SYNONYMS: Swelling

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Swelling is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Swelling should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Swelling? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Swelling; clinical terminology; patient terminology; suspected Swelling; symptoms associated with Swelling; diagnosis and management of Swelling.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

22. Rash

DISEASE: Rash

SYNONYMS: Rash

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Rash is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Rash should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Rash? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Rash; clinical terminology; patient terminology; suspected Rash; symptoms associated with Rash; diagnosis and management of Rash.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

23. Itching

DISEASE: Itching

SYNONYMS: Itching

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Itching is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Itching should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Itching? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Itching; clinical terminology; patient terminology; suspected Itching; symptoms associated with Itching; diagnosis and management of Itching.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

24. Weight loss

DISEASE: Weight loss

SYNONYMS: Weight loss

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Weight loss is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Weight loss should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Weight loss? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Weight loss; clinical terminology; patient terminology; suspected Weight loss; symptoms associated with Weight loss; diagnosis and management of Weight loss.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

25. Weight gain

DISEASE: Weight gain

SYNONYMS: Weight gain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Weight gain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Weight gain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Weight gain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Weight gain; clinical terminology; patient terminology; suspected Weight gain; symptoms associated with Weight gain; diagnosis and management of Weight gain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

26. Loss of appetite

DISEASE: Loss of appetite

SYNONYMS: Loss of appetite

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Loss of appetite is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Loss of appetite should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Loss of appetite? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Loss of appetite; clinical terminology; patient terminology; suspected Loss of appetite; symptoms associated with Loss of appetite; diagnosis and management of Loss of appetite.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

27. Difficulty swallowing

DISEASE: Difficulty swallowing

SYNONYMS: Difficulty swallowing

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Difficulty swallowing is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Difficulty swallowing should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Difficulty swallowing? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Difficulty swallowing; clinical terminology; patient terminology; suspected Difficulty swallowing; symptoms associated with Difficulty swallowing; diagnosis and management of Difficulty swallowing.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

28. Blood in stool

DISEASE: Blood in stool

SYNONYMS: Blood in stool

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Blood in stool is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Blood in stool should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Blood in stool? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Blood in stool; clinical terminology; patient terminology; suspected Blood in stool; symptoms associated with Blood in stool; diagnosis and management of Blood in stool.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

29. Blood in urine

DISEASE: Blood in urine

SYNONYMS: Blood in urine

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Blood in urine is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Blood in urine should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Blood in urine? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Blood in urine; clinical terminology; patient terminology; suspected Blood in urine; symptoms associated with Blood in urine; diagnosis and management of Blood in urine.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

30. Painful urination

DISEASE: Painful urination

SYNONYMS: Painful urination

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Painful urination is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Painful urination should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Painful urination? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Painful urination; clinical terminology; patient terminology; suspected Painful urination; symptoms associated with Painful urination; diagnosis and management of Painful urination.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

31. Frequent urination

DISEASE: Frequent urination

SYNONYMS: Frequent urination

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Frequent urination is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Frequent urination should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Frequent urination? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Frequent urination; clinical terminology; patient terminology; suspected Frequent urination; symptoms associated with Frequent urination; diagnosis and management of Frequent urination.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

32. Excessive thirst

DISEASE: Excessive thirst

SYNONYMS: Excessive thirst

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Excessive thirst is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Excessive thirst should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Excessive thirst? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Excessive thirst; clinical terminology; patient terminology; suspected Excessive thirst; symptoms associated with Excessive thirst; diagnosis and management of Excessive thirst.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

33. Excessive hunger

DISEASE: Excessive hunger

SYNONYMS: Excessive hunger

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Excessive hunger is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Excessive hunger should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Excessive hunger? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Excessive hunger; clinical terminology; patient terminology; suspected Excessive hunger; symptoms associated with Excessive hunger; diagnosis and management of Excessive hunger.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

34. Vision changes

DISEASE: Vision changes

SYNONYMS: Vision changes

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Vision changes is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Vision changes should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Vision changes? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Vision changes; clinical terminology; patient terminology; suspected Vision changes; symptoms associated with Vision changes; diagnosis and management of Vision changes.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

35. Hearing changes

DISEASE: Hearing changes

SYNONYMS: Hearing changes

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Hearing changes is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Hearing changes should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Hearing changes? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Hearing changes; clinical terminology; patient terminology; suspected Hearing changes; symptoms associated with Hearing changes; diagnosis and management of Hearing changes.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

36. Numbness

DISEASE: Numbness

SYNONYMS: Numbness

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Numbness is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Numbness should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Numbness? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Numbness; clinical terminology; patient terminology; suspected Numbness; symptoms associated with Numbness; diagnosis and management of Numbness.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

37. Tingling

DISEASE: Tingling

SYNONYMS: Tingling

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Tingling is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Tingling should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Tingling? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Tingling; clinical terminology; patient terminology; suspected Tingling; symptoms associated with Tingling; diagnosis and management of Tingling.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

38. Confusion

DISEASE: Confusion

SYNONYMS: Confusion

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Confusion is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Confusion should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Confusion? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Confusion; clinical terminology; patient terminology; suspected Confusion; symptoms associated with Confusion; diagnosis and management of Confusion.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

39. Seizures

DISEASE: Seizures

SYNONYMS: Seizures

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Seizures is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Seizures should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Seizures? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Seizures; clinical terminology; patient terminology; suspected Seizures; symptoms associated with Seizures; diagnosis and management of Seizures.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

40. Sleep problems

DISEASE: Sleep problems

SYNONYMS: Sleep problems

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Sleep problems is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Sleep problems should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Sleep problems? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Sleep problems; clinical terminology; patient terminology; suspected Sleep problems; symptoms associated with Sleep problems; diagnosis and management of Sleep problems.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

41. Anxiety

DISEASE: Anxiety

SYNONYMS: Anxiety

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Anxiety is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Anxiety should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Anxiety? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Anxiety; clinical terminology; patient terminology; suspected Anxiety; symptoms associated with Anxiety; diagnosis and management of Anxiety.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

42. Depression symptoms

DISEASE: Depression symptoms

SYNONYMS: Depression symptoms

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Depression symptoms is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Depression symptoms should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Depression symptoms? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Depression symptoms; clinical terminology; patient terminology; suspected Depression symptoms; symptoms associated with Depression symptoms; diagnosis and management of Depression symptoms.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

43. Nerve pain

DISEASE: Nerve pain

SYNONYMS: Nerve pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Nerve pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Nerve pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Nerve pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Nerve pain; clinical terminology; patient terminology; suspected Nerve pain; symptoms associated with Nerve pain; diagnosis and management of Nerve pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

44. Neuropathic pain

DISEASE: Neuropathic pain

SYNONYMS: Neuropathic pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Neuropathic pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Neuropathic pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Neuropathic pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Neuropathic pain; clinical terminology; patient terminology; suspected Neuropathic pain; symptoms associated with Neuropathic pain; diagnosis and management of Neuropathic pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

45. Chronic pain

DISEASE: Chronic pain

SYNONYMS: Chronic pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Chronic pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Chronic pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Chronic pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Chronic pain; clinical terminology; patient terminology; suspected Chronic pain; symptoms associated with Chronic pain; diagnosis and management of Chronic pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

46. Acute pain

DISEASE: Acute pain

SYNONYMS: Acute pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Acute pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Acute pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Acute pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Acute pain; clinical terminology; patient terminology; suspected Acute pain; symptoms associated with Acute pain; diagnosis and management of Acute pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

47. Cancer-related pain

DISEASE: Cancer-related pain

SYNONYMS: Cancer-related pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Cancer-related pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Cancer-related pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Cancer-related pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Cancer-related pain; clinical terminology; patient terminology; suspected Cancer-related pain; symptoms associated with Cancer-related pain; diagnosis and management of Cancer-related pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

48. Post-surgical pain

DISEASE: Post-surgical pain

SYNONYMS: Post-surgical pain

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Post-surgical pain is a medical condition in the pain, symptoms, and clinical presentations category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Pain, Symptoms, and Clinical Presentations. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Post-surgical pain should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Post-surgical pain? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Post-surgical pain; clinical terminology; patient terminology; suspected Post-surgical pain; symptoms associated with Post-surgical pain; diagnosis and management of Post-surgical pain.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

Category Sources / References

World Health Organization (WHO), Health Topics: https://www.who.int/health-topics

World Health Organization, International Classification of Diseases (ICD): https://www.who.int/standards/classifications/classification-of-diseases

WHO ICD-11 Browser: https://icd.who.int/en

Centers for Disease Control and Prevention (CDC), official clinical and public-health guidance: https://www.cdc.gov/

National health authorities and recognized professional medical organizations should be consulted for current local guidance.

Glossary

Acute: developing over a short period. Chronic: persistent or recurrent over time. Differential diagnosis: comparison of plausible conditions causing a presentation. Diagnostic evaluation: history, examination, and investigations used to assess a patient. Prognosis: expected course and outcome at a population or disease level, not an individual guarantee. Red flag: a feature that may indicate serious disease and requires urgent assessment.

Master References

WHO. Health Topics. Updated information and fact sheets. https://www.who.int/health-topics

WHO. International Classification of Diseases, 11th Revision. https://icd.who.int/en

CDC. Official disease, emergency, prevention, and infection-control guidance. https://www.cdc.gov/

National health ministries/agencies and recognized specialty organizations: use current jurisdiction-specific guidelines.

This knowledge base is intended for educational and information-retrieval purposes. It does not replace evaluation, diagnosis, treatment, or advice from a qualified healthcare professional. Symptoms can have many different causes, and the information presented here cannot determine an individual's diagnosis. Emergency symptoms require immediate professional medical evaluation.