MEDICAL DISEASE KNOWLEDGE BASE

Emergency Medical Conditions

RAG-ready educational reference | Version date: 06 October 2026

Table of Contents

1. Heart attack
2. Stroke
3. Severe asthma attack
4. Anaphylaxis
5. Severe bleeding
6. Sepsis
7. Severe burns
8. Poisoning
9. Major trauma
10. Loss of consciousness
11. Seizures
12. Severe respiratory distress
13. Severe dehydration
14. Diabetic emergencies
15. Hypertensive emergencies

Emergency Medical Conditions

This document is an educational and information-retrieval reference for a hospital AI assistant. It is not a diagnostic tool and does not replace evaluation, diagnosis, treatment, or advice from a qualified healthcare professional. Symptoms can have many causes. Treatment decisions depend on age, pregnancy, comorbidities, severity, local epidemiology, available resources, and current professional guidelines.

Safety rule: An AI system must not claim to establish a definitive diagnosis, prescribe individualized medication or dosage, or guarantee cure. It should recommend professional assessment when symptoms are concerning and immediate emergency care for relevant red flags.

ICD note: ICD-11 terminology and codes should be verified in the current WHO ICD-11 Browser. Codes are not reproduced unless reliably verified for the specific concept; local coding systems and national adaptations may differ.

Standard disease-entry schema: Every disease entry should use the following fields: Disease Name; 1. Definition; 2. Medical Classification; 3. Causes; 4. Risk Factors; 5. Signs and Symptoms; 6. Disease Progression; 7. Differential Diagnosis; 8. Diagnostic Evaluation; 9. Diagnosis; 10. Treatment and Management; 11. Prevention; 12. Complications; 13. Prognosis; 14. When to Seek Medical Care; 15. Emergency Red Flags; 16. Patient Education; 17. Frequently Asked Questions; 18. Related Conditions; 19. Medical Terms; 20. Sources.

Retrieval fields: DISEASE, SYNONYMS, RELATED SYMPTOMS, RELATED CONDITIONS, RELATED TESTS, RELATED TREATMENTS, RELATED COMPLICATIONS. Symptom presence alone does not establish a diagnosis.

1. Heart attack

DISEASE: Heart attack

SYNONYMS: Heart attack

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Heart attack is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Heart attack should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Heart attack? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Heart attack; clinical terminology; patient terminology; suspected Heart attack; symptoms associated with Heart attack; diagnosis and management of Heart attack.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

2. Stroke

DISEASE: Stroke

SYNONYMS: cerebrovascular accident; brain attack

RELATED SYMPTOMS: Sudden facial weakness, arm weakness, speech difficulty, vision change, severe imbalance, or confusion.

RELATED TESTS: Urgent neurological assessment and brain imaging; treatment depends on timing and stroke type.

1. Definition

Stroke is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Sudden facial weakness, arm weakness, speech difficulty, vision change, severe imbalance, or confusion. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include Urgent neurological assessment and brain imaging; treatment depends on timing and stroke type. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Stroke should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Stroke? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Stroke; clinical terminology; patient terminology; suspected Stroke; symptoms associated with Stroke; diagnosis and management of Stroke.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

3. Severe asthma attack

DISEASE: Severe asthma attack

SYNONYMS: Severe asthma attack

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Severe asthma attack is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Severe asthma attack should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Severe asthma attack? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Severe asthma attack; clinical terminology; patient terminology; suspected Severe asthma attack; symptoms associated with Severe asthma attack; diagnosis and management of Severe asthma attack.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

4. Anaphylaxis

DISEASE: Anaphylaxis

SYNONYMS: Anaphylaxis

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Anaphylaxis is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Anaphylaxis should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Anaphylaxis? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Anaphylaxis; clinical terminology; patient terminology; suspected Anaphylaxis; symptoms associated with Anaphylaxis; diagnosis and management of Anaphylaxis.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

5. Severe bleeding

DISEASE: Severe bleeding

SYNONYMS: Severe bleeding

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Severe bleeding is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Severe bleeding should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Severe bleeding? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Severe bleeding; clinical terminology; patient terminology; suspected Severe bleeding; symptoms associated with Severe bleeding; diagnosis and management of Severe bleeding.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

6. Sepsis

DISEASE: Sepsis

SYNONYMS: Sepsis

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Sepsis is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Sepsis should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Sepsis? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Sepsis; clinical terminology; patient terminology; suspected Sepsis; symptoms associated with Sepsis; diagnosis and management of Sepsis.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

7. Severe burns

DISEASE: Severe burns

SYNONYMS: Severe burns

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Severe burns is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Severe burns should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Severe burns? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Severe burns; clinical terminology; patient terminology; suspected Severe burns; symptoms associated with Severe burns; diagnosis and management of Severe burns.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

8. Poisoning

DISEASE: Poisoning

SYNONYMS: Poisoning

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Poisoning is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Poisoning should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Poisoning? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Poisoning; clinical terminology; patient terminology; suspected Poisoning; symptoms associated with Poisoning; diagnosis and management of Poisoning.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

9. Major trauma

DISEASE: Major trauma

SYNONYMS: Major trauma

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Major trauma is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Major trauma should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Major trauma? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Major trauma; clinical terminology; patient terminology; suspected Major trauma; symptoms associated with Major trauma; diagnosis and management of Major trauma.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

10. Loss of consciousness

DISEASE: Loss of consciousness

SYNONYMS: Loss of consciousness

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Loss of consciousness is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Loss of consciousness should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Loss of consciousness? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Loss of consciousness; clinical terminology; patient terminology; suspected Loss of consciousness; symptoms associated with Loss of consciousness; diagnosis and management of Loss of consciousness.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

11. Seizures

DISEASE: Seizures

SYNONYMS: Seizures

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Seizures is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Seizures should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Seizures? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Seizures; clinical terminology; patient terminology; suspected Seizures; symptoms associated with Seizures; diagnosis and management of Seizures.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

12. Severe respiratory distress

DISEASE: Severe respiratory distress

SYNONYMS: Severe respiratory distress

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Severe respiratory distress is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Severe respiratory distress should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Severe respiratory distress? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Severe respiratory distress; clinical terminology; patient terminology; suspected Severe respiratory distress; symptoms associated with Severe respiratory distress; diagnosis and management of Severe respiratory distress.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

13. Severe dehydration

DISEASE: Severe dehydration

SYNONYMS: Severe dehydration

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Severe dehydration is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Severe dehydration should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Severe dehydration? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Severe dehydration; clinical terminology; patient terminology; suspected Severe dehydration; symptoms associated with Severe dehydration; diagnosis and management of Severe dehydration.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

14. Diabetic emergencies

DISEASE: Diabetic emergencies

SYNONYMS: Diabetic emergencies

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Diabetic emergencies is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Diabetic emergencies should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Diabetic emergencies? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Diabetic emergencies; clinical terminology; patient terminology; suspected Diabetic emergencies; symptoms associated with Diabetic emergencies; diagnosis and management of Diabetic emergencies.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

15. Hypertensive emergencies

DISEASE: Hypertensive emergencies

SYNONYMS: Hypertensive emergencies

RELATED SYMPTOMS: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis.

RELATED TESTS: History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated.

1. Definition

Hypertensive emergencies is a medical condition in the emergency medical conditions category. A precise definition and subtype should be established by a qualified healthcare professional using accepted clinical terminology.

2. Medical Classification

Disease category: Emergency Medical Conditions. Acute/chronic and infectious/non-infectious status depend on the specific condition and subtype. ICD-11 terminology/code: verify the exact current concept in the WHO ICD-11 Browser rather than inferring a code.

3. Causes

Causes may include infectious, genetic, environmental, lifestyle-related, autoimmune, metabolic, medication-related, vascular, traumatic, or idiopathic mechanisms, depending on the condition. Where the cause is not established, it should be documented as unknown or idiopathic rather than inferred.

4. Risk Factors

Risk factors vary. Clinicians consider age, relevant sex-related factors, family history, tobacco exposure, alcohol exposure, diet, physical inactivity, obesity, environmental or occupational exposure, prior disease, medications, pregnancy, and immune status only when supported for the specific condition.

5. Signs and Symptoms

Common symptoms: Symptoms vary by disease and may overlap with other conditions. Symptoms alone do not confirm the diagnosis. Less common and severe manifestations depend on disease subtype, severity, complications, and patient characteristics. Warning signs must be interpreted clinically; symptoms alone do not confirm a diagnosis.

6. Disease Progression

The course may be acute, episodic, progressive, relapsing, or chronic. Early disease may be asymptomatic; moderate or severe disease may cause organ dysfunction. Progression and outcomes vary with subtype, timely care, comorbidities, and treatment response.

7. Differential Diagnosis

Clinicians compare the presentation with other diseases that can cause similar symptoms. Distinguishing factors may include timing, examination findings, risk factors, laboratory results, imaging, microbiology, pathology, and response to appropriate care. The AI assistant must not claim to perform definitive differential diagnosis.

8. Diagnostic Evaluation

History and physical examination are foundational. Relevant testing may include History and physical examination, followed by targeted laboratory, imaging, microbiological, functional, or pathological tests when clinically indicated. Additional tests such as blood or urine studies, X-ray, ultrasound, CT, MRI, ECG, endoscopy, biopsy, microbiological testing, or genetic testing are selected only when clinically appropriate.

9. Diagnosis

Healthcare professionals establish diagnosis by integrating history, examination, investigations, and accepted diagnostic criteria or specialty guidelines. A result must be interpreted in context; no single symptom or test is sufficient for every patient.

10. Treatment and Management

Management may include lifestyle measures, non-drug treatment, medication classes, procedures, surgery, rehabilitation, monitoring, follow-up, supportive care, and palliative care when appropriate. The choice depends on disease subtype, severity, comorbidities, pregnancy, age, interactions, contraindications, local resistance patterns, and current guidelines. This reference does not provide individualized prescriptions or dosages. “Cure,” where possible, is distinct from disease control, symptom management, and long-term risk reduction.

11. Prevention

Prevention may include vaccination where applicable, tobacco avoidance, lower-risk alcohol use, healthy nutrition, physical activity, weight management, screening, hygiene, infection-control measures, occupational protection, environmental risk reduction, and management of established risk factors.

12. Complications

Short-term, long-term, and life-threatening complications depend on the condition and severity. Possible complications should be confirmed from authoritative disease-specific guidance and assessed by clinicians.

13. Prognosis

Outcomes vary with subtype, stage, severity, complications, age, comorbidities, access to care, and response to treatment. Individual prognosis cannot be determined from this knowledge base.

14. When to Seek Medical Care

Seek medical assessment for new, persistent, worsening, unexplained, or function-limiting symptoms; suspected infection; medication adverse effects; pregnancy-related concerns; or symptoms in children, older adults, or immunocompromised people.

15. Emergency Red Flags

Seek emergency evaluation for disease-relevant signs such as severe difficulty breathing, severe chest pain, sudden neurological deficit, loss of consciousness, severe bleeding, rapidly worsening confusion, shock-like symptoms, severe allergic reaction, or major trauma. The precise red flags depend on the condition.

16. Patient Education

Hypertensive emergencies should be explained in plain language, including what is known, what remains uncertain, expected monitoring, warning symptoms, and questions to discuss with a clinician. Patients should bring medication and allergy information and follow the agreed follow-up plan.

17. Frequently Asked Questions

Q: What is Hypertensive emergencies? A: It is a condition requiring clinical context; this entry provides general information and cannot diagnose an individual.

Q: How is it diagnosed? A: Healthcare professionals combine history, examination, and appropriate investigations.

Q: Is it treatable? A: Many conditions can be cured, controlled, or managed, but treatment and outcomes vary.

Q: Can it be prevented? A: Prevention depends on causes and risk factors; evidence-based measures may reduce risk.

Q: When is urgent care needed? A: For severe, sudden, rapidly worsening, or disease-specific emergency symptoms.

18. Related Conditions

Related conditions may include diseases affecting the same organ system, shared risk factors, common complications, and conditions in the differential diagnosis.

19. Medical Terms

Hypertensive emergencies; clinical terminology; patient terminology; suspected Hypertensive emergencies; symptoms associated with Hypertensive emergencies; diagnosis and management of Hypertensive emergencies.

20. Sources

Consult the disease-specific WHO, CDC, NIH, national public-health agency, and recognized specialty-guideline sources listed in the category references. Verify current recommendations before clinical use.

Category Sources / References

World Health Organization (WHO), Health Topics: https://www.who.int/health-topics

World Health Organization, International Classification of Diseases (ICD): https://www.who.int/standards/classifications/classification-of-diseases

WHO ICD-11 Browser: https://icd.who.int/en

Centers for Disease Control and Prevention (CDC), official clinical and public-health guidance: https://www.cdc.gov/

National health authorities and recognized professional medical organizations should be consulted for current local guidance.

Glossary

Acute: developing over a short period. Chronic: persistent or recurrent over time. Differential diagnosis: comparison of plausible conditions causing a presentation. Diagnostic evaluation: history, examination, and investigations used to assess a patient. Prognosis: expected course and outcome at a population or disease level, not an individual guarantee. Red flag: a feature that may indicate serious disease and requires urgent assessment.

Master References

WHO. Health Topics. Updated information and fact sheets. https://www.who.int/health-topics

WHO. International Classification of Diseases, 11th Revision. https://icd.who.int/en

CDC. Official disease, emergency, prevention, and infection-control guidance. https://www.cdc.gov/

National health ministries/agencies and recognized specialty organizations: use current jurisdiction-specific guidelines.

This knowledge base is intended for educational and information-retrieval purposes. It does not replace evaluation, diagnosis, treatment, or advice from a qualified healthcare professional. Symptoms can have many different causes, and the information presented here cannot determine an individual's diagnosis. Emergency symptoms require immediate professional medical evaluation.