

Topological Phase Dynamics of the Relativistic Vacuum

A Complete Physical Theory of Matter, Gravity, and Quantum Behavior

Branden Lee Friend¹

¹*Private work. All rights reserved.*

(Dated: February 20, 2026)

This theory presents a complete physical description of the universe based on a single continuous relativistic medium. The medium’s state is described by a real scalar phase field $\varphi(x^\mu)$ with intrinsic energy density, mechanical stiffness, and topological structure. All observed phenomena—matter, gravity, electromagnetism, quantum behavior, and gauge interactions—arise from the dynamics and topology of this medium. General Relativity and the Standard Model are recovered as effective macroscopic descriptions in the appropriate limits. The theory excludes singularities via a finite maximum vacuum tension scale, provides a mechanical origin of mass as stored field energy and reconfiguration resistance, predicts quantized charge and spin as topological constraints, identifies dark energy with quiescent vacuum pressure, and attributes dark-matter phenomenology to large-scale vacuum stress structure. A quantitative closure program is specified: particle masses, couplings, and mixing parameters are computed as defect-energy and overlap integrals of solutions of the master field equation under topological boundary conditions.

I. FUNDAMENTAL PHYSICAL POSTULATE

There exists a single continuous relativistic medium filling all regions of spacetime. There is no empty space, no void, and no background grid separate from this medium. Its state at any spacetime point is fully described by a real scalar phase field

$$\varphi(x^\mu), \quad x^\mu = (ct, x, y, z). \quad (1)$$

A. Key properties

- **Quiescent energy density:** the medium has a nonzero base energy density

$$\rho_{\text{vac}} \approx 5.35 \times 10^{-10} \text{ J/m}^3. \quad (2)$$

- **Mechanical stiffness:** finite stiffness characterized by coupling constant λ .
- **Maximum tension:** finite maximum compression/tension scale

$$T_{\text{vac}} \sim 10^{113} \text{ J/m}^3. \quad (3)$$

- **Topological support:** the medium supports stable finite-energy topological defects (knotted configurations).
- **Continuity:** there is no ontological “nothingness” underneath the field: the vacuum is the physical substrate.

II. LAGRANGIAN DYNAMICS

A. Action and Lagrangian density

The medium is governed by a Lorentz-invariant Lagrangian density

$$\mathcal{L} = \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - V(\varphi), \quad (4)$$

with action

$$S[\varphi] = \int d^4x \mathcal{L}. \quad (5)$$

B. Stabilizing potential

The stabilizing potential is

$$V(\varphi) = \lambda(\varphi^2 - \varphi_0^2)^2. \quad (6)$$

Physical meaning: φ_0 is the quiescent vacuum value; λ sets resistance to deformation; the potential stores elastic energy and stabilizes finite-core defects.

C. Master field equation

Applying Euler–Lagrange variation yields

$$\square \varphi + \frac{dV}{d\varphi} = J_{\text{top}}, \quad (7)$$

i.e.

$$\square \varphi + 4\lambda \varphi(\varphi^2 - \varphi_0^2) = J_{\text{top}}, \quad (8)$$

where J_{top} represents defect-supported topological source terms encoding boundary conditions and nontrivial winding.

III. STRESS-ENERGY TENSOR AND VACUUM PRESSURE

The mechanical stress is described by the canonical stress-energy tensor

$$T_{\mu\nu} = \partial_\mu \varphi \partial_\nu \varphi - g_{\mu\nu} \left(\frac{1}{2} \partial_\alpha \varphi \partial^\alpha \varphi - V(\varphi) \right). \quad (9)$$

For a quiescent vacuum where $\varphi = \varphi_0$ and $\partial_\mu \varphi = 0$,

$$T_{\mu\nu}^{\text{vac}} = -g_{\mu\nu} V(\varphi_0) \equiv -g_{\mu\nu} \rho_{\text{vac}} c^2. \quad (10)$$

The vacuum pressure is

$$P_{\text{vac}} = -\rho_{\text{vac}} c^2, \quad (11)$$

corresponding to dark-energy equation of state $w = -1$.

IV. GRAVITY AS VACUUM STRESS GRADIENT

Gravity is not curvature of empty space; it is mechanical acceleration caused by gradients in vacuum pressure.

A. Force density and acceleration

Define force density

$$f^\mu = -\partial^\mu P_{\text{vac}}. \quad (12)$$

For a test configuration, the acceleration is

$$\mathbf{a} = -\frac{1}{\rho_{\text{vac}}} \nabla P_{\text{vac}}. \quad (13)$$

B. Physical mechanism

A massive object is a topological defect that displaces the vacuum field from φ_0 . This produces a region of reduced vacuum pressure. The surrounding quiescent vacuum at higher pressure pushes configurations toward the defect. This is the mechanical origin of gravitational attraction.

C. Connection to General Relativity

The metric tensor is an effective field describing the stress state of the medium:

$$g_{\mu\nu}^{\text{eff}} = \eta_{\mu\nu} + h_{\mu\nu}(\varphi, \partial\varphi, \dots), \quad (14)$$

where $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$. Einstein-type equations are recovered as the hydrodynamic/macrosopic limit of vacuum stress conservation.

V. MATTER AS TOPOLOGICAL DEFECTS

Matter is not separate from the vacuum. A particle is a stable, localized topological defect of φ .

A. Defect energy and mass

The defect energy is

$$E = \int \left[\frac{1}{2} (\nabla \varphi)^2 + V(\varphi) \right] d^3x, \quad (15)$$

and the mass is

$$m = \frac{E}{c^2}. \quad (16)$$

B. Inertial mass

Mass is the inertial resistance of a defect to acceleration through the medium. Operationally,

$$m = \frac{1}{c^2} \left. \frac{d^2 E}{dv^2} \right|_{v=0}. \quad (17)$$

Accelerating a defect requires continuous deformation of the surrounding vacuum; resistance to that reconfiguration is inertia.

C. Higgs reinterpretation

The Higgs field is reinterpreted as an effective description of vacuum stiffness λ and medium response at electroweak scales, rather than a fundamental mass-giving substance.

VI. DISCRETE MOTION AND UNIVERSAL SPEED LIMIT

Motion is a sequence of discrete state transitions of the vacuum medium: dissolution at one region and re-emergence at the next. This occurs at Planck-scale time resolution

$$t_P \approx 5.39 \times 10^{-44} \text{ s}. \quad (18)$$

The universal speed limit is the maximum speed of phase propagation:

$$c = \sqrt{\frac{T_{\text{vac}}}{\rho_{\text{vac}}}}. \quad (19)$$

VII. CHARGE QUANTIZATION

Charge arises from quantized circulation of the phase field around defects. Define

$$A_\mu = \partial_\mu \varphi. \quad (20)$$

The circulation around a closed loop is

$$\oint A_\mu dx^\mu = \oint \partial_\mu \varphi dx^\mu = \Delta \varphi. \quad (21)$$

Because φ is a phase variable, single-valued consistency demands

$$\Delta \varphi = 2\pi n, \quad n \in \mathbb{Z}, \quad (22)$$

and charge is quantized as $q = ne$.

VIII. SPIN

Spin is intrinsic angular momentum of vacuum circulation around a defect core. Define angular momentum density schematically

$$\mathbf{L} = \epsilon_{ijk} x_j (\partial_0 \varphi) (\partial_k \varphi), \quad (23)$$

and total spin

$$\mathbf{S} = \int \mathbf{L} d^3x. \quad (24)$$

Quantization follows from topological constraints on stable circulation classes:

$$S = n \frac{\hbar}{2}, \quad n \in \mathbb{Z}. \quad (25)$$

Fermions vs. bosons correspond to distinct topological symmetry classes of defect configurations.

IX. QUANTUM MECHANICS

The wavefunction is the physical phase structure of the medium, not a probability amplitude in a void:

$$\psi(\mathbf{x}, t) = \sqrt{\rho(\mathbf{x}, t)} e^{i\varphi(\mathbf{x}, t)/\hbar}. \quad (26)$$

A. Schrödinger limit

In the nonrelativistic/WKB regime, the effective dynamics reproduce Schrödinger-type evolution for defect-center motion in an external environment:

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi + V_{\text{ext}} \psi. \quad (27)$$

B. Quantum potential

The Bohm quantum potential

$$Q = -\frac{\hbar^2}{2m} \frac{\nabla^2 \sqrt{\rho}}{\sqrt{\rho}} \quad (28)$$

is interpreted as a mechanical curvature/stress term of the medium.

C. Entanglement and measurement

Entanglement corresponds to shared global phase topology of the continuous medium. Measurement is mechanical coupling of a microscopic defect to macroscopic detector boundary conditions. The transition from coherent superposition to a definite state is physical medium evolution governed by Eq. (8). In this interpretation, “collapse” is not an axiom: it is the mechanical convergence of the coupled system (defect + apparatus) to a stable boundary-constrained phase configuration.

Nonlocal correlations are attributed to shared global topological constraints of a single continuous substrate. No signal propagates superluminally; constraint updates are mediated by physical phase-stress propagation bounded by the medium wave speed c .

X. GAUGE STRUCTURE

A. U(1) electromagnetism as defect-supported connection

Define an effective potential

$$A_\mu \equiv \kappa \partial_\mu \varphi, \quad (29)$$

where κ is a constant fixing physical units. Away from defect cores where φ is smooth and globally single-valued, A_μ is locally pure gauge and

$$F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu = 0. \quad (30)$$

Physical electromagnetic field strength arises when the vacuum phase is *not* globally trivial: defects make φ effectively multi-valued (or undefined) at cores, producing distributional curvature supported on defect worldlines/worldsheets. In that case one writes schematically

$$\partial_{[\mu} \partial_{\nu]} \varphi \neq 0 \quad (\text{distributionally at defect support}), \quad (31)$$

and Maxwell-type equations are interpreted as macroscopic conservation laws for defect-supported circulation current.

B. Topological current and effective Maxwell equations

Introduce an effective topological current J^ν associated with defect circulation/transport such that

$$\partial_\mu F^{\mu\nu} = J^\nu, \quad (32)$$

and $\partial_\nu J^\nu = 0$ expresses conservation of defect-supported circulation number.

C. $SU(2)$ weak sector as internal circulation doublet

The weak interaction is attributed to coupled internal circulation modes of a defect, represented effectively by a two-component internal orientation degree of freedom. For macroscopic modeling, introduce a doublet Φ and an $SU(2)$ connection $W_\mu = W_\mu^a \tau^a / 2$ encoding how the defect's internal phase-frame rotates through spacetime:

$$D_\mu \Phi = (\partial_\mu - igW_\mu)\Phi. \quad (33)$$

Parity violation is attributed to chiral topology of admissible defect configurations in the vacuum medium: only one handedness of stable circulation couples in the low-energy limit.

D. $SU(3)$ strong sector and Yang–Mills emergence

Color is identified with three independent internal circulation orientations required for stable baryonic composite defects. Introduce an $SU(3)$ connection $G_\mu = G_\mu^a T^a$ (with generators T^a) and field strength

$$\mathcal{G}_{\mu\nu} = \partial_\mu G_\nu - \partial_\nu G_\mu + ig_s [G_\mu, G_\nu]. \quad (34)$$

A macroscopic effective Yang–Mills + matter form is

$$\mathcal{L}_{\text{YM}} = -\frac{1}{2} \text{Tr} (\mathcal{G}_{\mu\nu} \mathcal{G}^{\mu\nu}) + (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi), \quad (35)$$

leading to Yang–Mills equations

$$D_\mu \mathcal{G}^{\mu\nu} = J_{\text{top}}^\nu, \quad (36)$$

interpreted as stress-equilibrium conditions for non-Abelian internal twist of the vacuum medium sourced by defect currents.

E. Confinement and flux tubes

An isolated fractional circulation (quark-like) defect class produces divergent energy in the surrounding medium because the phase cannot relax to the quiescent state without completing an integer topological class. Separation of fractional defects produces a narrow tube of stressed vacuum (flux tube). The tube tension σ is an emergent mechanical constant of the medium set by stiffness and defect-core structure, consistent with the empirical scale $\sim \text{GeV/fm}$.

XI. DERIVATION PROGRAM FOR CONSTANTS AND STANDARD MODEL PARAMETERS

This section states the end-to-end computational closure as explicit inputs/outputs of the master equation.

A. Constitutive identifications and vacuum scales

The theory uses the vacuum scales ρ_{vac} and T_{vac} . The wave speed identification is

$$c = \sqrt{\frac{T_{\text{vac}}}{\rho_{\text{vac}}}}. \quad (37)$$

The potential stiffness scale is parameterized by λ and φ_0 via Eq. (6).

B. Defect spectrum and particle masses

Each particle species corresponds to a topological class $\mathcal{C}$ and a stable solution $\varphi_{\mathcal{C}}$ of Eq. (8) under the associated boundary conditions. The predicted mass is

$$m_{\mathcal{C}} = \frac{1}{c^2} \int \left[\frac{1}{2} (\nabla \varphi_{\mathcal{C}})^2 + V(\varphi_{\mathcal{C}}) \right] d^3x. \quad (38)$$

Closure requirement: the theory is fully numerically closed when the defect classes and boundary conditions are uniquely specified and the solutions are computed without fitting.

C. Charge, spin, and coupling constants

Charge is fixed by the integer circulation condition Eq. (??); spin is fixed by the intrinsic circulation class. Dimensionless couplings (e.g. α) are identified with ratios of energy stored in transverse circulation modes to longitudinal phase-wave modes; their computation reduces to energy minimization for the relevant defect topology under the vacuum constitutive parameters.

D. Mixing matrices (CKM and PMNS) as overlap integrals

Flavor/generation structure is attributed to distinct stable defect harmonics. Mixing elements are defined as overlap functionals between normalized defect modes:

$$V_{ij} = \langle \Phi_i | \Phi_j \rangle \equiv \int d^3x \Phi_i^\dagger(\mathbf{x}) \Phi_j(\mathbf{x}), \quad (39)$$

with Φ_i constructed from the internal orientation content of the defect solutions. **Closure requirement:** specify the mode basis and inner product induced by the medium's kinetic term, then compute V_{ij} from the numerically obtained defect fields.

TABLE I. Characteristic energy scales used in the theory.

Property	Symbol	Value (as used)
Quiescent energy density	ρ_{vac}	$5.35 \times 10^{-10} \text{ J/m}^3$
Vacuum stiffness (max tension)	T_{vac}	$\sim 10^{113} \text{ J/m}^3$
Planck time	t_P	$5.39 \times 10^{-44} \text{ s}$
Speed of light	c	$2.998 \times 10^8 \text{ m/s}$

E. Gravitational constant

A mechanical derivation of G requires a defined vacuum correlation length L (set by the linearized spectrum about φ_0 and/or defect-core regularization). An effective scaling used in this theory is

$$G \sim \frac{c^4}{8\pi\rho_{\text{vac}}L^2}, \quad (40)$$

with L computed from the vacuum constitutive parameters rather than inserted phenomenologically.

F. Planck constant

$\hbar$ is identified as the minimum action associated with a stable 2π phase transition of the medium. Its computation reduces to evaluating the action for the minimal topological transition event (instanton/phase-slip) supported by the vacuum potential and stiffness:

$$\hbar \sim S_{\text{min}}/(2\pi), \quad (41)$$

where S_{min} is the minimal stable phase-action of the substrate.

XII. ENERGY SCALE TABLE

XIII. COSMOLOGY

A. Dark energy

Dark energy is identified with quiescent vacuum pressure:

$$\rho_{\text{DE}} = \rho_{\text{vac}}, \quad P_{\text{DE}} = -\rho_{\text{vac}}c^2. \quad (42)$$

B. Dark matter

Dark matter phenomenology is attributed to large-scale topological stress structure and flow patterns of the vacuum medium that couple gravitationally (pressure gradients) but do not carry electromagnetic circulation.

C. Inflation as phase transition and vacuum crystallization

Inflation is interpreted as rapid relaxation of the vacuum field from a high-energy initial configuration toward the quiescent state φ_0 . Domain formation and topological freezing during this transition imprint primordial perturbations. In slow-roll form one writes

$$\ddot{\varphi} + 3H\dot{\varphi} + \frac{dV}{d\varphi} = 0, \quad (43)$$

and the near-scale-invariant spectrum arises from finite stiffness and a finite number of e-folds required for topological stabilization.

D. No singularities and Planck-density cores

The theory forbids singularities by imposing a finite saturation scale T_{vac} . Black-hole interiors are replaced by finite-radius Planck-density cores in which compression is bounded and repulsive stiffness terms balance inward pressure.

E. Bekenstein–Hawking entropy

Entropy is interpreted as counting vacuum topological microstates supported on the boundary of a saturated core. The area scaling is recovered by counting Planck-scale phase cells on the boundary:

$$S \sim \frac{k_B A}{4\ell_P^2}. \quad (44)$$

F. Arrow of time

The arrow of time is attributed to topological decay: complex high-energy knots transition irreversibly toward lower-energy configurations, corresponding to global relaxation toward the vacuum minimum.

G. Redshift as medium dissipation (tired-light claim)

A specific cosmological claim of the theory is that observed redshift may be interpreted as dissipation/attenuation of propagating phase waves through a vacuum medium with extremely small effective viscosity. This is an explicit empirical fork: the theory must satisfy all observational constraints currently supporting expansion (time dilation in supernova light curves, Tolman surface-brightness test, CMB consistency). This section is therefore a primary target for decisive observational validation or falsification.

XIV. TESTABLE PREDICTIONS (BENCHMARKS)

This theory makes empirically distinguishing predictions:

1. **Gravitational-wave dispersion:** frequency-dependent propagation speed due to finite stiffness of the medium.
2. **Vacuum birefringence:** polarization rotation in ultra-strong fields from induced anisotropic vacuum stress.
3. **Departures from GR in saturation regimes:** near neutron stars/compact objects where the effective-metric description breaks down as T_{vac} is approached.
4. **Quantized angular momentum emission steps:** discrete defect state transitions in high-energy scattering/decay.
5. **CMB polarization signatures:** patterns tied to vacuum crystallization/topological freezing.

XV. WHY I REJECT “NORMAL PHYSICS” (ONTOLOGY STATEMENT)

The core rejection is ontological: modern physics often treats the vacuum as empty and then introduces mathematical objects as placeholders for missing physical substrate.

A. Metric as placeholder versus stress summary

The metric tensor is not taken as a fundamental substance of “nothing.” In this theory it is an effective macroscopic summary of vacuum stress state. Geometry is a language for medium stress, not a replacement for physical substance.

B. Point particles versus finite-core defects

Point particles are rejected as unphysical infinite-density idealizations. Matter is finite-core topological defect structure of a real medium; infinities and singularities are excluded by stiffness and saturation.

C. Gauge bosons and Higgs as effective descriptions

Gauge bosons are interpreted as collective excitation modes and alignment dynamics of the vacuum; the Higgs description is an effective stiffness parameterization of the medium, not a separate mass-giving substance.

D. Wavefunction realism

Probability amplitudes are rejected as fundamental ontology. The wavefunction is physical phase structure of the vacuum medium; “collapse” is deterministic mechanical coupling under boundary constraints.

E. Dark sector

Dark energy and dark matter are not separate substances. They are vacuum properties: quiescent pressure and large-scale stress/topological structure.

F. Virtual particles and extra dimensions

Virtual particles are treated as perturbative bookkeeping artifacts rather than literal particles. Extra dimensions are rejected as unnecessary if the vacuum medium in $(3 + 1)$ dimensions supplies the missing physical substrate.

XVI. STATUS AND PATH TO FULL QUANTITATIVE CLOSURE

The theory is ontologically closed by one medium, one field, one potential, and one master equation. Full quantitative completion requires:

- explicit classification of defect topology classes corresponding to particle species,
- numerical solution of Eq. (8) for each class,
- evaluation of mass integrals Eq. (38),
- computation of overlap integrals defining mixing matrices,
- derivation of dispersion relations for the benchmark predictions,
- direct confrontation with cosmological expansion tests if tired-light dissipation is maintained.

XVII. CONCLUSION

Topological Phase Dynamics of the Relativistic Vacuum asserts a single physical substrate: a continuous relativistic vacuum medium described by a real scalar phase field. Matter is stable topological defect structure; inertia is reconfiguration resistance; gravity is vacuum pressure-gradient mechanics; and quantum behavior is physical phase dynamics. General Relativity and gauge field theories are recovered as macroscopic effective descriptions. The theory is rendered scientific by explicit computational closure tasks and falsifiable benchmark predictions.

-
- [1] J. A. Wheeler, Phys. Rev. **97**, 511 (1955).
 - [2] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, 1973).
 - [3] T. H. R. Skyrme, Proc. Roy. Soc. A **260**, 127 (1961).
 - [4] L. de Broglie, J. Phys. Radium **8**, 225 (1927).
 - [5] D. Bohm, Phys. Rev. **85**, 166 (1952).
 - [6] A. Einstein, *Ether and the Theory of Relativity* (Leiden Lecture, 1920).
 - [7] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford University Press, 2003).
 - [8] R. B. Laughlin, *A Different Universe: Reinventing Physics from the Bottom Down* (Basic Books, 2005).
 - [9] A. D. Sakharov, Sov. Phys. Dokl. **12**, 1040 (1968).
 - [10] D. Hestenes, Found. Phys. **20**, 1213 (1990).
 - [11] D. Hestenes, *New Foundations for Classical Mechanics*, 2nd ed. (Springer, 1999).
 - [12] S. Weinberg, *The Quantum Theory of Fields, Vol. I* (Cambridge University Press, 1995).