

Canonical Equations Across Civilizations and Historical Figures

Executive summary

This report assembles a cross-civilizational catalog of “canonical” equations and equation-like rules—spanning early mensuration, number theory, algebra, optics, mechanics, field theory, thermodynamics, and quantum/relativistic physics—and then analyzes how later formalisms *embed* earlier structures (as limits, special cases, or re-expressions under new invariants). The catalog is necessarily selective: it aims for breadth across the requested civilizations while focusing on results that (a) are explicitly attested in primary texts or high-quality critical editions/translations, and (b) became structurally foundational in later mathematical physics. ¹

A central finding is that, across cultures, many “equations” first appear not as symbolic equalities but as **procedures, proportional rules, or geometric constructions**—often tuned to practical tasks (surveying, construction, taxation, astronomical prediction, lens design, etc.). For example, the Rhind Mathematical Papyrus ² encodes an area-of-circle rule operationally as “subtract 1/9 of the diameter; square the remainder,” rather than naming π as a constant. ³ Likewise, Muhammad ibn Musa al-Khwarizmi ⁴’s algebra is framed as “completion and reduction” with worked word-problems and explicit type-cases, not a general quadratic formula in modern notation. ⁵ In optics, Ibn Sahl ⁶’s refraction work expresses a constant ratio (equivalent to Snell’s law) via segment geometry and sine ratios, anticipating a later “law” formulation. ⁷

The relationship mapping shows repeated *structural motifs* reappearing with increasing abstraction:

- **Geometric invariants → algebraic invariants → variational invariants → symmetry invariants.**
- Inner-product geometry generalizes the Pythagorean relation; variational calculus unifies “least” principles in optics and mechanics; gauge and geometric formulations recast Maxwell and Einstein equations into coordinate-free objects (forms, tensors), clarifying what is physically invariant versus notational artifact. ⁸

Translation into modern standards introduces predictable anachronisms. Most importantly: (i) assigning modern concepts (angle, function, equation, field) to texts that used different conceptual primitives; and (ii) importing modern “constants” (π , refractive index n , $\hbar$) into contexts where they were not explicit objects. The report identifies such mismatches and gives “least-distorting” modern reformulations that preserve the original task-structure while making assumptions explicit. ⁹

Finally, the unification section prioritizes targets where reconciliation is mathematically explicit (by derivation or limiting process), and flags cases where reconciliation is inherently speculative (e.g., disputed claims about what a tablet was *for*, or uncertain transmission routes of a method across language spheres).

Catalog of canonical equations and formulas

Scope note: the table emphasizes (1) attested primary statements (or critical translations where the original is non-symbolic), (2) equations that later became “nodes” in physics/mathematics curricula, and (3) coverage across the requested civilizations. Dates are approximate and often bracket a manuscript’s copying vs. composition. ¹¹

Catalog table

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
Egypt	Circle area rule (RMP mensuration)	Rhind Mathematical Papyrus ²	c. 1550 BCE (copy)	Operationally: <i>subtract 1/9 of diameter; square remainder</i> (used to get “64” from diameter 9). ³	$A \approx \left(\frac{8}{9}D\right)^2 = \frac{64}{81}D^2$, i.e. $A \approx \pi r^2$ with $\pi \approx 256/81$. ³	Geometric mensuration
Egypt	Cylinder/bin volume via circle rule	Same as above	c. 1550 BCE (copy)	Compute base area by “(diameter – 1/9 diameter) ² ” then multiply by height for contents. ³	$V \approx \left(\frac{8}{9}D\right)^2 h$. ³	Geometric applied
Egypt	Pyramid slope parameter (“seked”)	Same as above	c. 1550 BCE (copy)	Non-symbolic ratio “palms per cubit” describing the relation of horizontal run to vertical rise. ³	$\text{seked} \sim \cot(\theta) = \frac{\Delta x}{\Delta z}$ (up to unit conversion). ³	Geometric trigonometry
Egypt	Frustum of pyramid volume (MMP Problem 14)	Moscow Mathematical Papyrus ¹²	c. 1850 BCE (copy date often cited)	Worked example: square 4 → 16; double 4 → 8; square 2 → 4; sum 28; take 1/3 of 6 → 2; twice 28 → 56. ¹³	$V = \frac{h}{3}(a^2 + ab + b^2)$ for square frustum with top side b , bottom side a . ¹³	Geometric volume

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
Mesopotamia	$\sqrt{2}$ approximation (diagonal of unit square)	Cuneiform Digital Library Initiative ¹⁴ artifact YBC 07289	Old Babylonian (ca. 1900–1600 BCE)	Sexagesimal entry: 1; 24, 51, 10 labeled as Babylonian $\sqrt{2}$, with halved value 0; 42, 25, 35. ¹⁵	$\sqrt{2} \approx 1.41421296$ (very close to modern). ¹⁵	Arithmetic geometry
Mesopotamia	Right-triangle relation via tabulation (Plimpton-type triples)	CDLI record for Plimpton tablet + critical reassessment	Old Babylonian (ca. 1900–1600 BCE)	Tabular data lists pairs corresponding to integer sides/diagonals; modern reconstructions show systematic generation by parameter pairs. ¹⁶	Special cases of $s^2 + \ell^2 = d^2$ (right triangles); table supports generating/using triples. ¹⁷	Number theory geometry
Greece	Pythagorean theorem (axiomatic proof tradition)	Euclid ¹⁹ , Elements ²⁰	c. 300 BCE	Proposition form (Book I, Prop. 47): squares on legs equal square on hypotenuse (in Euclidean geometry). ²¹	For right triangle: $a^2 + b^2 = c^2$. ²¹	Geometry
Greece	Euclidean algorithm (GCD)	Same source as above	c. 300 BCE	Algorithmic subtraction/division process to find “greatest measure” (common divisor). ²¹	$\gcd(a, b) = \gcd(b, a \bmod b)$. ²¹	Number theory algorithm

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
Greece	Sphere volume / area relations	Archimedes ²² (via Heath's collected translation)	3rd century BCE	Results in geometric ratio form: sphere vs circumscribed cylinder relationships, expressed without algebraic symbols. ²³	$V_{\text{sphere}} = \frac{4}{3}\pi r^3$, $A_{\text{sphere}} = 4\pi r^2$, and $V_{\text{sphere}} = \frac{2}{3}V_{\text{cyl}}$. ²³	Geometric integration
India	Diagonal-area principle (Śulba tradition; "rope rule")	Katyayana Sulba Sutra ²⁴	often dated ~800–200 BCE (text strata)	Discusses diagonal/square relation as area-doubling or rectangle-composition principle (stated as a construction rule). ²⁵	Right-triangle relation $a^2 + b^2 = c^2$ (interpreting "areas" of squares on sides). ²⁵	Geometric construction
India	"Square on diagonal is double" (special case)	Same as above	~800–200 BCE (text strata)	Commentary explicitly links diagonal of a square to area doubling. ²⁵	For square side s : diagonal $d = s\sqrt{2}$ and $d^2 = 2s^2$. ²⁵	Geometric
India	Power series / infinite series for π (Kerala school attention point)	Secondary discussion of Kerala series + archival Madhava text	c. 14th–16th century CE (attribution varies)	Series representations for π and trig (infinite procedures). ²⁶	Prototype: $\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \dots$ and related expansions for $\sin, \cos$. ²⁶	Analysis

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
China	Chinese remainder theorem (canonical puzzle form)	Historical development of the Chinese remainder theorem ²⁸	Text tradition often dated 3rd–5th c. CE (Sunzi)	Classic residues: “÷3 remainder 2; ÷5 remainder 3; ÷7 remainder 2”; solution constructed additively. ²⁹	Solve $x \equiv 2 \pmod{3}$, $x \equiv 3 \pmod{5}$, $x \equiv 2 \pmod{7} \Rightarrow x \equiv 23 \pmod{105}$. ²⁹	Number modular arithmetic
China	Fangcheng elimination (rectangular arrays)	Solving a System of Linear Equations Using Ancient Chinese Methods ³⁰	Core text assembled by early CE; commentary 3rd c. CE	Coefficients set on counting board; operations eliminate unknowns column/row-wise. ³¹	Gaussian-elimination-like reduction of $A\mathbf{x} = \mathbf{b}$. ³¹	Linear algebra
Islamic Golden Age	Canonical quadratic solving by “completion and reduction”	The algebra of Mohammed ben Musa ³³	c. 820 CE (composition)	Example type: “a square and ten roots amount to thirty-nine”; solve by halving roots, squaring, adding, square root, subtract. ³⁴	For $x^2 + bx = c$: $x = \frac{\sqrt{(b/2)^2 + c} - b/2}{1}$. ⁵	Algebra
Islamic Golden Age	Taxonomy of linear/quadratic equation types	Same as above	c. 820 CE	Enumerates cases like “squares equal roots,” “roots equal numbers,” etc. ³⁵	Modern classification of degree-1/2 polynomial equations; proto-normal forms under allowed operations. ³⁵	Algebra foundations

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
Islamic Golden Age	Refraction ratio equivalent to Snell's law	A Pioneer in Anaclastics: Ibn Sahl on Burning Mirrors and Lenses ³⁶	Treatise dated ~983–985 CE	For plane interface, gives CE/CH as constant ("reciprocal of refractive index"), with sine relation shown. ⁷	$n_1 \sin \theta_1 = n_2 \sin \theta_2$ (in the paper's notation: $CE/CH = \sin i_1 / \sin i_2 = 1/n$). ⁷	Optics physics
Islamic Golden Age	Experimental constraints on refraction (qualitative laws)	Alhacen on Refraction ³⁷	11th century CE (original), 2010 ed.	Experimental synopsis: light travels straight within each medium, refraction occurs in a plane, bending toward normal entering water, no refraction at normal incidence. ³⁸	Qualitative boundary behavior; modern form is Snell + piecewise straight rays. ³⁸	Optics experim methods
Europe	Cartesian analytic geometry (coordinates and equations for curves)	René Descartes ³⁹ , The geometry of René Descartes ⁴⁰	1637	Uses algebra to represent curves; early explicit coordinate usage is documented in extracts. ⁴¹	Coordinate representation of loci, e.g. line $y = mx + b$, conics as quadratic forms. ⁴²	Geometric algebra

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
Europe	Uniform acceleration (Galilean kinematics)	Galileo Galilei ⁴³ , Dialogues Concerning Two New Sciences ⁴⁴	1638	Treatise addresses acceleration and projectile motion; the falling-body law appears in the development of motion theory. ⁴⁵	For constant a : $s = \frac{1}{2}at^2$ (from rest). ⁴⁶	Mechanics
Europe	Kepler's harmonic law	Johannes Kepler ⁴⁷ , Harmonices mundi libri V ⁴⁸	1619	Published as a proportional relation among orbital period and "mean distance." ⁴⁹	$T^2 \propto a^3$ for bodies orbiting same primary (modern semi- major axis a). ⁵⁰	Celestial mechanics
Europe	Newtonian mechanics + gravitation	Isaac Newton ⁵² , The mathematical principles of natural philosophy ⁵³	1687 (orig.), 1729 trans.	Foundational laws and inverse-square gravitation stated/ proved geometrically. ⁵⁴	$\mathbf{F} = m\mathbf{a}$; $F = G \frac{m_1 m_2}{r^2}$. ⁵⁴	Mechanics gravitation

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
Europe	Hamiltonian dynamics (canonical equations)	William Rowan Hamilton ⁵⁶ , On a General Method in Dynamics ⁵⁷	1834	Presented as a “general method in dynamics” with a characteristic function framework. ⁵⁸	Canonical equations: $\dot{q}_i = \partial H / \partial p_i$, $\dot{p}_i = -\partial H / \partial q_i$. ⁵⁸	Analytical mechanics
Modern	Maxwell field equations (classical EM)	James Clerk Maxwell ⁶⁰ , A Dynamical Theory of the Electromagnetic Field ⁶¹	1865	Field-theoretic framework culminating in EM wave propagation at light speed. ⁶²	$\nabla \cdot \mathbf{E} = \rho / \epsilon_0$, $\nabla \cdot \mathbf{B} = 0$, $\nabla \times \mathbf{E} = -\partial_t \mathbf{B}$, $\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \partial_t \mathbf{E}$. ⁶³	Electrodynamics
Modern	Planck radiation law + quantization seed	Max Planck ⁶⁵ , On the Law of Distribution of Energy in the Normal Spectrum ⁶⁶	1900–1901	Derives blackbody spectrum consistent with data by assuming energy elements. ⁶⁷	$u(\nu, T) = \frac{8\pi h \nu^3}{c^3} \frac{1}{e^{h\nu/kT} - 1}$ (one standard form). ⁶⁸	Thermodynamics, quantum precursors
Modern	Special relativity kinematics	Albert Einstein ⁶⁹ , On the Electrodynamics of Moving Bodies ⁷⁰	1905	Reformulates space/time for consistency with electrodynamics; translation notes adjust notation. ⁷¹	Lorentz transform; invariant interval $s^2 = c^2 t^2 - x^2 - y^2 - z^2$; energy–momentum relations. ⁷²	Relativity

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
Modern	General relativity field equations	The Field Equations of Gravitation ⁷⁴	1915	Published formulation of gravitational field equations (translation editions exist). ⁷⁵	$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$. ⁷⁶	Relativistic gravitation
Modern	Noether's theorem backbone (symmetry $\rightarrow$ conservation)	Emmy Noether ⁷⁸ , Invariant Variation Problems ⁷⁹	1918	Proves systematic link between continuous symmetries of an action and conservation laws. ⁸⁰	If $S = \int L dt$ invariant under parameter ϵ , then $\exists$ conserved current/quantity. ⁸⁰	Foundational mechanics theory
Modern	Schrödinger wave equation	Erwin Schrödinger ⁸¹ , Quantisation as an eigenvalue problem ⁸²	1926	Frames quantization as eigenvalue problem for wavefunctions. ⁸³	$i\hbar\partial_t\psi = \hat{H}\psi$; stationary $\hat{H}\psi = E\psi$. ⁸³	Quantum mechanics
Modern	Matrix mechanics principle (observable quantities)	Werner Heisenberg ⁸⁵ , On the quantum-theoretical reinterpretation of kinematical and mechanical relationships ⁸⁶	1925	Proposes mechanics built from relations among observables; early matrix structure. ⁸⁷	Canonical commutator (later form): $[x, p] = i\hbar$. ⁸⁷	Quantum mechanics

Tradition	Equation / rule (canonical label)	Original source / author	Approx. date	Original notation / statement (translated or paraphrased)	Modern standardized notation	Domain
Modern	Uncertainty inequality (modern standard form)	Historical analyses of 1927–1928 formalization	1927–1928	Heisenberg gives intuitive tradeoffs; later derivation yields variance inequality. ⁸⁹	$\sigma_x \sigma_p \geq \hbar/2.$ ⁸⁸	Quantum statistical
Modern	Dirac relativistic electron equation	Paul Dirac ⁹⁰ , The Quantum Theory of the Electron ⁹¹	1928	Relativistic wave equation for electron; paper digitizations exist via scholarly archives. ⁹²	$(i\gamma^\mu \partial_\mu - m)\psi = 0.$ ⁹²	Quantum relativistic
Modern	Path-integral formulation (equivalent quantum mechanics)	Richard Feynman ⁹⁴ , Space-Time Approach to Non-Relativistic Quantum Mechanics ⁹⁵	1948	Reformulates quantum mechanics as sum over histories; stated as mathematically equivalent to familiar formulations. ⁹⁶	$\langle x_f, t_f x_i, t_i \rangle = \int \mathcal{D}x e^{\frac{i}{\hbar} S[x]}.$ ⁹⁶	Quantum variational methods

Structural relationships and influence mappings

The strongest “structural” through-line is not a single genealogy but a repeated pattern: **procedures** → **proofs** → **symbol systems** → **invariants** → **coordinate-free objects**. Across the catalog, later theories tend to *embed* earlier rules as (i) special cases, (ii) limiting regimes, or (iii) reparameterizations under a new invariant quantity. ⁹⁷

```

graph TD
  A[Mensuration & ratios as procedures] --> B[Axiomatic geometry & proof]
  A --> C[Algebra as transformations on types]
  B --> D[Analytic geometry & coordinates]
  C --> D
  D --> E[Calculus & variational principles]
  E --> F[Analytical mechanics: Lagrange/Hamilton]
  F --> G[Field theory: Maxwell]
  F --> H[Statistical/thermo potentials]
  G --> I[Special relativity constraints]
  I --> J[General relativity geometry]
  F --> K[Quantum mechanics operator/sum-over-paths]
  E --> K
  K --> L[Quantum-field methods]

```

This dependency graph is concretely anchored by examples in the sources: procedural mensuration in Egyptian papyri, rhetorical algebra and quadratic completion, variational/symmetry formalisms, and the field-theoretic turn from Newtonian forces to Maxwellian fields and Einsteinian geometry. ⁹⁸

A particularly clear mapping is optics → variational mechanics → quantum:

- 1) In refraction/lens construction, Ibn Sahl encodes the medium-pair constant via sine/segment geometry, functionally equivalent to Snell's law. ⁷
- 2) In later European science, least-time/least-action ideas are reframed as a general variational calculus (leading into Hamiltonian methods and, with Noether, symmetry as the generator of conservation structure). ⁵⁹
- 3) Feynman's formulation reuses the action functional as the phase of a path amplitude, making "variational structure" a backbone that survives quantization (with important measure/rigor caveats). ⁹⁹

A second mapping—geometry/number theory → algebra/linear algebra—is visible when comparing the conceptual units of expression. The Chinese "rectangular array" method is procedurally close to elimination but is not a matrix concept originally; modern matrix notation is best viewed as a **lossless encoding of operations**, not evidence of identical conceptual objects. ³² The same caution applies to Babylonian tables: different modern reconstructions treat Plimpton-style data as (i) generating right-triangle integer triples (Robson's context-based critique) or (ii) a kind of ratio-table with trigonometric flavor (Mansfield/Wildberger's thesis). ¹⁸

```

timeline
  title Threaded timeline of selected canonical equations
  2000 BCE : Egypt frustum volume algorithm (MMP) ; Egypt circle area rule (RMP)
  1800 BCE : Old Babylonian sqrt(2) tablet ; Plimpton 322 parameter table
  300 BCE : Euclid axiomatic Pythagorean theorem; Euclid GCD algorithm
  250 BCE : Archimedes sphere/cylinder volume relations
  500 CE : Sunzi CRT puzzle form ; Nine Chapters elimination tradition

```

820 CE : al-Khwarizmi completion/reduction quadratics
 984 CE : Ibn Sahl refraction ratio (Snell-equivalent)
 1020 CE : Ibn al-Haytham experimental refraction regularities
 1619 CE : Kepler harmonic law
 1638 CE : Galileo accelerated motion
 1687 CE : Newton laws + inverse-square gravity
 1834 CE : Hamilton canonical dynamics
 1865 CE : Maxwell field equations and EM waves
 1905 CE : Special relativity kinematics
 1915 CE : Einstein field equations
 1918 CE : Noether symmetry-conservation
 1925–1928 CE : Heisenberg/Schrödinger/Dirac quantum formalisms
 1948 CE : Feynman path integral

Dating anchors and primary textual points in this timeline come from the cited document metadata and translations. ¹⁰⁰

Translation issues, anachronisms, and modern reformulations

A recurring anachronism is the temptation to project modern *objects* (constants, functions, abstract equation-forms) onto sources that used **actions and constructions** instead. The least-distorting modern reformulation is therefore often: “identify the invariant being computed” (area, volume, ratio, residue class) and then express the procedure as an equation in a modern algebraic encoding.

Egyptian π vs “8/9 diameter square.” The Rhind rule is frequently rewritten as $A = \pi r^2$ with $\pi \approx 256/81$, which is mathematically correct, but historically misleading if it implies the explicit concept of π as a constant already in use. A better reformulation is to treat the rule as a *map* $D \mapsto A$: $A := ((8/9)D)^2$. Only secondarily do we infer an “effective π .” ³

Ambiguity of geometric terms. Even within the same translated source, geometry terms can be unclear: the Rhind discussion notes uncertainty about whether a “meret” line is an altitude or a side in triangle-area computations; a modern re-expression $A = \frac{1}{2}bh$ is only correct when the “height” parameter is indeed the altitude. The least-anachronistic fix is to write a conditional statement: “If the given segment is the altitude, then $A = \frac{1}{2}bh$; otherwise the computed area is approximate and shape-constrained.” ³

Mesopotamian right triangles vs “Pythagorean theorem” and “trigonometry.” The safest translation is to say the tablets support computations consistent with the right-triangle relation (and table generation of integer triples), without assuming the same theorem/proof status or angle-based trigonometric concept. Robson argues that reading Plimpton 322 as a trigonometric table conflicts with what is known about Old Babylonian conceptions of angle; Mansfield/Wildberger argue for a ratio-based “sexagesimal trigonometry” not using angles. A conservative reformulation is: “tabulated exact rational ratios related to right triangles,” leaving the interpretive layer explicit. ¹⁰¹

Chinese elimination vs matrix theory. Modern linear algebra writes $A\mathbf{x} = \mathbf{b}$ and row-reduces A . The “Fangcheng” method can be encoded that way, but that does not mean the original concept was “matrices”

as abstract objects. Cullen emphasizes the danger of treating symbolic algebra as the thinking itself; modern notation should be treated as scaffolding, not the historical cognition. ¹⁰²

Algebra: “completion and reduction” vs quadratic formula. Translating al-Khwarizmi as if he “knew the quadratic formula” collapses a genuine structural difference: he frames problems through a taxonomy of cases (squares/roots/numbers) and only permits transformations that keep quantities positive and meaningful as magnitudes. A faithful modern reformulation is to present the algorithm as completing the square for a constrained normal form, rather than as an unconditional polynomial identity over $\mathbb{R}$. ¹⁰³

Quantum uncertainty: inequality vs interpretive claims. Modern textbooks often present $\sigma_x \sigma_p \geq \hbar/2$ as “Heisenberg’s uncertainty principle,” but historical analyses emphasize that early presentations mixed several notions: preparation uncertainty, joint measurement limits, and disturbance tradeoffs. A careful translation distinguishes (i) the mathematically general variance inequality from (ii) specific measurement-disturbance statements that require extra modeling assumptions. ⁸⁹

Unification and generalization proposals with derivations

This section prioritizes unifications that (a) cover many catalog items as special cases, and (b) have explicit derivations demonstrating how “earlier” equations embed into “later” formalisms. Where the historical record is interpretively contested, the unification is presented as a *mathematical embedding* (modern structural map), not as a claim about what an ancient author intended. ¹⁰⁴

Unification proposal: inner-product geometry as the umbrella for right-triangle relations.

Earlier: right-triangle rules (Egyptian seked geometry, Babylonian diagonal computation, Sulba diagonal-area laws, Euclid’s theorem). ¹⁰⁵

Modern unification: define an inner product $\langle \cdot, \cdot \rangle$ and norm $\|v\|^2 = \langle v, v \rangle$. Orthogonality $\langle u, v \rangle = 0$ implies

$$\|u + v\|^2 = \langle u + v, u + v \rangle = \|u\|^2 + \|v\|^2 + 2\langle u, v \rangle = \|u\|^2 + \|v\|^2,$$

which is the Pythagorean relation. This generalizes from Euclidean triangles to Hilbert spaces, clarifying that the “square” structure is a property of quadratic norms, not a special property of geometric squares.

Unification proposal: “completion of the square” as a cross-cultural normal-form engine.

Earlier: al-Khwarizmi’s canonical example $x^2 + 10x = 39$ solved by halving the root coefficient, squaring, adding, square-rooting, then subtracting. ⁵

Modern derivation: for $x^2 + bx = c$, add $(b/2)^2$ to both sides:

$$x^2 + bx + (b/2)^2 = c + (b/2)^2 \Rightarrow (x + b/2)^2 = c + (b/2)^2,$$

so $x = \sqrt{c + (b/2)^2} - b/2$ (positive root in the historical constraint set). This unifies ancient quadratic procedures with modern polynomial algebra, while preserving the original constraint that solutions correspond to meaningful magnitudes. ⁵

Unification proposal: variational calculus connects optics, mechanics, symmetries, and quantum amplitudes.

Anchor nodes: Hamilton’s dynamics program, Noether’s invariance theorems, and Feynman’s action-phase

formulation. ¹⁰⁶

Modern derivation sketch: start from an action functional

$$S[q] = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt.$$

Setting $\delta S = 0$ yields the Euler-Lagrange equation

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}} - \frac{\partial L}{\partial q} = 0.$$

Define conjugate momentum $p = \partial L / \partial \dot{q}$ and Hamiltonian $H = p\dot{q} - L$; then Hamilton's canonical equations follow:

$$\dot{q} = \frac{\partial H}{\partial p}, \quad \dot{p} = -\frac{\partial H}{\partial q}.$$

Noether adds: if S is invariant under a continuous transformation, a conserved quantity/current follows. Feynman reuses S not as a minimand but as a phase factor $e^{iS/\hbar}$, making classical mechanics emerge in a stationary-phase limit. ¹⁰⁶

Unification proposal: geometric optics refraction law $\rightarrow$ constitutive parameter as invariant.

Ibn Sahl expresses refraction through a constant ratio CE/CH and an equivalent sine-ratio. ⁷ Modern optics rewrites this as Snell's law $n_1 \sin \theta_1 = n_2 \sin \theta_2$, where "refractive index" is precisely the invariant parameter linking two media. The unification here is conceptual: segment geometry, sine geometry, and modern n -based law encode the same invariant mapping at an interface.

Unification proposal: Maxwell's equations as differential-forms identities (coordinate-free EM).

Maxwell's field unification is already explicit in the 19th-century move from forces to fields. ⁶² Modern differential forms provide a compact embedding: let F be the electromagnetic 2-form, $*F$ its Hodge dual. Then

$$dF = 0, \quad d*F = J$$

(where J is the current 3-form) encodes the homogeneous and inhomogeneous Maxwell equations, respectively. This reformulation clarifies what is invariant (forms, exterior derivative) and what is coordinate artifact (vector calculus component expressions). Background on differential forms as a systematic alternative calculus is given in standard notes. ¹⁰⁷

Unification proposal: thermodynamic potentials as Legendre-transform family.

A large fraction of "thermodynamics equations" are not independent; they are generated by Legendre transforms among conjugate variables. A pedagogical modern summary explicitly connects mechanics' $L \leftrightarrow H$ transform and thermodynamics' U to free-energy potentials. ¹⁰⁸

Concretely, if $U(S, V, \dots)$ then define $F(T, V, \dots) = U - TS$ and $G(T, P, \dots) = U + PV - TS$. The unification is that each "new" potential is the same underlying state function reparameterized to natural variables; many "equations" become differential identities under this mapping. ¹⁰⁹

Unification proposal: gravity as geometry that reproduces Newton as a limit.

Einstein's field equations embed Newtonian gravity as a weak-field, low-velocity approximation (recovering

classical predictions in the appropriate regime). ¹¹⁰ This is a mathematically explicit embedding: in the limit $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ with small h , the Einstein equations linearize and yield a Poisson-like equation for the gravitational potential (under standard gauge choices), matching Newtonian gravity's structure. (Full derivations are standard; the key validation point is that the embedding is central to the physical acceptability of GR.) ¹¹⁰

Validation against contemporary mathematics and physics

Where reconciliation is robust (non-speculative).

Many catalog items are *exactly* consistent with modern mathematics when interpreted as procedures or theorems in their natural domain: Euclid's Pythagorean theorem is valid in Euclidean geometry; Archimedes' sphere/cylinder relations are correct within classical geometry of exhaustion; al-Khwarizmi's completing-the-square algorithm is algebraically correct for the constrained positive-magnitude cases he treats; Ibn Sahl's sine-ratio expression is equivalent to the modern Snell law parameterization for refraction; Maxwell's field formalism matches modern classical electromagnetism; Hamilton's canonical equations are a standard modern backbone for mechanics; and Noether's theorem is a cornerstone for conservation laws in modern physics. ¹¹¹

Where reconciliation is only approximate.

Egyptian mensuration is internally consistent as a computation system but uses approximations (notably the implied $\pi \approx 256/81$), so "validation" must be interpreted as "correct algorithm for their approximation scheme," not "exact analytic identity." ³ Similarly, any finite truncation of infinite series for π or trig requires an explicitly stated error bound to be meaningfully "validated" in modern analysis terms; this is often present as a practical remainder-control technique rather than a later formal convergence proof. ²⁶

Where reconciliation is structurally correct but historically delicate.

Encoding Fangcheng elimination as matrix row reduction is mathematically exact as a representation of operations, but it can misstate the historical conceptual content if presented as "they had matrices." The correct reconciliation is a *representation theorem*: modern matrices encode the same elimination operations, without implying identity of conceptual ontology. ³²

Where reconciliation becomes interpretive or speculative.

Plimpton 322 is the clearest case: one can map the tablet's numbers into modern right-triangle relations and even into ratio tables that resemble trigonometric data, but scholars disagree over whether "trigonometry" is an appropriate historical description. Therefore, any claim of "first trigonometric table" is interpretively loaded; a safer validation is that the tablet data are consistent with exact rational relationships of right triangles in sexagesimal arithmetic, and multiple modern reconstructions exist. ¹⁰¹ Transmission claims can also become speculative: for example, discussions of how certain medieval Islamic astronomical devices or methods may have influenced later European work often note manuscript evidence but may not establish a definitive chain. Where the record is uncertain, this report treats such links as plausible hypotheses rather than confirmed lineages. ¹¹²

¹ ³ ⁹ ¹⁹ ²² ³⁶ ⁷⁴ ⁷⁹ ⁹⁷ ⁹⁸ https://ia903207.us.archive.org/2/items/the-rhind-papyrus/The%20Rhind%20Papyrus_text.pdf
https://ia903207.us.archive.org/2/items/the-rhind-papyrus/The%20Rhind%20Papyrus_text.pdf

2 14 26 27 91 <https://mathshistory.st-andrews.ac.uk/Strick/brahmagupta.pdf>
<https://mathshistory.st-andrews.ac.uk/Strick/brahmagupta.pdf>

4 13 100 <https://tildesites.geneseo.edu/~johannes/egypt.pdf>
<https://tildesites.geneseo.edu/~johannes/egypt.pdf>

5 34 35 43 69 90 103 https://archive.org/stream/algebraofmohamme00khuwuoft/algebraofmohamme00khuwuoft_djvu.txt
https://archive.org/stream/algebraofmohamme00khuwuoft/algebraofmohamme00khuwuoft_djvu.txt

6 107 <https://www.math.purdue.edu/~arapura/preprints/diffforms.pdf>
<https://www.math.purdue.edu/~arapura/preprints/diffforms.pdf>

7 57 78 85 95 https://files.backyardbrains.com/retired/files/Rashed_1990_Ibn_Sahl.pdf
https://files.backyardbrains.com/retired/files/Rashed_1990_Ibn_Sahl.pdf

8 21 66 82 94 111 <https://hypertextbook.com/eworld/chords/>
<https://hypertextbook.com/eworld/chords/>

10 17 18 24 28 33 44 65 81 101 <https://mathcs.holycross.edu/~little/Mont201617/RobsonNeither.pdf>
<https://mathcs.holycross.edu/~little/Mont201617/RobsonNeither.pdf>

11 https://www.britishmuseum.org/collection/object/Y_EA10057
https://www.britishmuseum.org/collection/object/Y_EA10057

12 23 70 https://commons.warburg.sas.ac.uk/concern/published_works/r494vk17h?locale=en
https://commons.warburg.sas.ac.uk/concern/published_works/r494vk17h?locale=en

15 105 <https://cdli.ucla.edu/P255048>
<https://cdli.ucla.edu/P255048>

16 <https://cdli.earth/artifacts/254790>
<https://cdli.earth/artifacts/254790>

20 31 32 https://digitalcommons.ursinus.edu/context/triumphs_linear/article/1001/viewcontent/F3_Solving_a_System_of_Linear_Equations_Using_Ancient_Chinese_Methods_12.11.2018.pdf
https://digitalcommons.ursinus.edu/context/triumphs_linear/article/1001/viewcontent/F3_Solving_a_System_of_Linear_Equations_Using_Ancient_Chinese_Methods_12.11.2018.pdf

25 40 https://ia601403.us.archive.org/9/items/in.ernet.dli.2015.133882/2015.133882.Katyayana-Sulba-Sutra_text.pdf
https://ia601403.us.archive.org/9/items/in.ernet.dli.2015.133882/2015.133882.Katyayana-Sulba-Sutra_text.pdf

29 <https://people.math.harvard.edu/~knill/crt/lib/Kangsheng.pdf>
<https://people.math.harvard.edu/~knill/crt/lib/Kangsheng.pdf>

30 60 75 76 77 86 110 https://reprints.gravitywaves.com/Gravitation/Einstein-1915_TheFieldEquationsOfGravitation_EFEs_Wikisource.pdf
https://reprints.gravitywaves.com/Gravitation/Einstein-1915_TheFieldEquationsOfGravitation_EFEs_Wikisource.pdf

37 87 https://www.neo-classical-physics.info/uploads/3/4/3/6/34363841/heisenberg_-_qm_interp_of_kin_and_mech.pdf
https://www.neo-classical-physics.info/uploads/3/4/3/6/34363841/heisenberg_-_qm_interp_of_kin_and_mech.pdf

38 52 53 https://archive.org/stream/TAPhS100-3/TAPhS%20100%2C%203%2C%20Secc.%202%2C%20A.%20Mark%20Smith%20-%20Alhacen%20on%20Refraction%20Volume%202%2C%20English%20Text_djvu.txt
https://archive.org/stream/TAPhS100-3/TAPhS%20100%2C%203%2C%20Secc.%202%2C%20A.%20Mark%20Smith%20-%20Alhacen%20on%20Refraction%20Volume%202%2C%20English%20Text_djvu.txt

39 42 <https://www.math.stonybrook.edu/~tony/archive/336s20/documents/from%20La%20Geometrie.pdf>
<https://www.math.stonybrook.edu/~tony/archive/336s20/documents/from%20La%20Geometrie.pdf>

41 <https://archive.org/download/geometryofrene00desc/geometryofrene00desc.pdf>
<https://archive.org/download/geometryofrene00desc/geometryofrene00desc.pdf>

45 <https://oll.libertyfund.org/titles/galilei-dialogues-concerning-two-new-sciences>
<https://oll.libertyfund.org/titles/galilei-dialogues-concerning-two-new-sciences>

46 https://sites.pitt.edu/~jdnorton/papers/Galileo_fall_final.pdf
https://sites.pitt.edu/~jdnorton/papers/Galileo_fall_final.pdf

47 56 80 104 https://web.stanford.edu/~cantwell/AA218_Course_Material/Resources/Invariant_Variation_Problems_Emma_Noether.pdf
https://web.stanford.edu/~cantwell/AA218_Course_Material/Resources/Invariant_Variation_Problems_Emma_Noether.pdf

48 83 84 https://ofp.cosmo-ufes.org/uploads/1/3/7/0/13701821/quantisation_as_an_eigenvalue_problem.pdf
https://ofp.cosmo-ufes.org/uploads/1/3/7/0/13701821/quantisation_as_an_eigenvalue_problem.pdf

49 <https://library.si.edu/digital-library/book/ioanniskeplerih00kepl>
<https://library.si.edu/digital-library/book/ioanniskeplerih00kepl>

50 <https://inters.org/Giostra-Third-Law-Kepler>
<https://inters.org/Giostra-Third-Law-Kepler>

51 https://en.wikipedia.org/wiki/Kepler%27s_laws_of_planetary_motion
https://en.wikipedia.org/wiki/Kepler%27s_laws_of_planetary_motion

54 55 https://archive.org/details/bim_eighteenth-century_principia-english-th_newton-sir-isaac_1729_1
https://archive.org/details/bim_eighteenth-century_principia-english-th_newton-sir-isaac_1729_1

58 59 106 <https://www.maths.tcd.ie/pub/HistMath/People/Hamilton/Dynamics/GenMeth.pdf>
<https://www.maths.tcd.ie/pub/HistMath/People/Hamilton/Dynamics/GenMeth.pdf>

61 62 64 <https://royalsocietypublishing.org/rstl/article/doi/10.1098/rstl.1865.0008/118816/VIII-A-dynamical-theory-of-the-electromagnetic>
<https://royalsocietypublishing.org/rstl/article/doi/10.1098/rstl.1865.0008/118816/VIII-A-dynamical-theory-of-the-electromagnetic>

63 <https://www.bem.fi/library/1865-001.pdf>
<https://www.bem.fi/library/1865-001.pdf>

67 68 <https://strangepaths.com/files/planck1901.pdf>
<https://strangepaths.com/files/planck1901.pdf>

71 72 73 https://sites.pitt.edu/~jdnorton/teaching/Einstein_graduate/pdfs/Einstein_STR_1905_English_reset.pdf
https://sites.pitt.edu/~jdnorton/teaching/Einstein_graduate/pdfs/Einstein_STR_1905_English_reset.pdf

88 89 <https://eprints.whiterose.ac.uk/id/eprint/3610/1/Buschp2.pdf>
<https://eprints.whiterose.ac.uk/id/eprint/3610/1/Buschp2.pdf>

92 93 <https://wwwhome.lorentz.leidenuniv.nl/~boyarsky/media/Proc.R.Soc.Lond.-1928-Dirac-610-24.pdf>
<https://wwwhome.lorentz.leidenuniv.nl/~boyarsky/media/Proc.R.Soc.Lond.-1928-Dirac-610-24.pdf>

96 99 <https://jontalle.web.engr.illinois.edu/uploads/Feynman.48.pdf>
<https://jontalle.web.engr.illinois.edu/uploads/Feynman.48.pdf>

102 https://people.math.harvard.edu/~knill/teaching/math22a2018/exhibits/ninechapters/Cullen_NineChapters.pdf
https://people.math.harvard.edu/~knill/teaching/math22a2018/exhibits/ninechapters/Cullen_NineChapters.pdf

108 109 https://web.physics.wustl.edu/alford/physics/Legendre_introduction.pdf
https://web.physics.wustl.edu/alford/physics/Legendre_introduction.pdf

112 <https://blogs.loc.gov/loc/2016/01/newly-acquired-arabic-manuscript-on-early-astronomy-and-mathematics-2/>
<https://blogs.loc.gov/loc/2016/01/newly-acquired-arabic-manuscript-on-early-astronomy-and-mathematics-2/>