

The Andromeda Simultaneity Problem and Phase-State Projection Model of Reality

Branden Lee Friend

November 7, 2025

Abstract

This paper presents a complete mathematical framework for the Phase-State Projection (PSP) Model, which resolves fundamental questions about the nature of simultaneity in special relativity. By treating observers as active projectors with phase-state vectors operating on a universal informational field Ψ , we derive both classical relativistic effects and novel testable predictions. The Andromeda Simultaneity Problem serves as a motivating example for the deeper informational substrate proposed by the PSP framework.

1 The Andromeda Simultaneity Problem

1.1 Overview

The Andromeda Simultaneity Problem, proposed by Roger Penrose, demonstrates the relativity of simultaneity in special relativity. Two observers walking past each other on Earth may disagree about what is happening "now" in the Andromeda galaxy—separated by several days—despite only moving at walking speed relative to each other.

1.2 Key Variables

Symbol	Meaning
v	Relative velocity between observers
c	Speed of light in vacuum
L	Distance to distant event (e.g., Andromeda)
$\Delta t'$	Time difference in simultaneity at distant location
γ	Lorentz factor: $\gamma = 1/\sqrt{1 - v^2/c^2}$

1.3 Classical Formula: Relativity of Simultaneity

The shift in simultaneity for a distant event is given by:

$$\Delta t' = \frac{vL}{c^2}$$

1.4 Derivation from Lorentz Transformation

Starting from the Lorentz transformation for time:

$$t' = \gamma \left(t - \frac{vx}{c^2} \right)$$

At local time $t = 0$ and distant position $x = L$ (distance to Andromeda):

$$t'(x = L) = -\gamma \frac{vL}{c^2}$$

For $v \ll c$, we have $\gamma \approx 1$, yielding:

$$\Delta t \approx \frac{vL}{c^2}$$

1.5 Example Calculation

Consider two observers with:

$$\begin{aligned} v &= 1 \text{ m/s} \\ L &= 2.5 \times 10^6 \text{ ly} = 2.365 \times 10^{22} \text{ m} \\ c &= 3 \times 10^8 \text{ m/s} \end{aligned}$$

Then:

$$\Delta t = \frac{(1)(2.365 \times 10^{22})}{(3 \times 10^8)^2} \approx 2.63 \times 10^5 \text{ s} \approx 3 \text{ days}$$

Result: Two people walking past each other with just 1 m/s velocity difference will disagree by approximately 3 days about what is happening "now" in Andromeda.

2 The Phase-State Projection (PSP) Model

2.1 Motivation

While special relativity describes *how* simultaneity changes between frames, it does not explain *why* relative motion should alter what is "real" at distant locations. The PSP model proposes that reality emerges from observer-dependent projections of a universal informational field Ψ , with motion modulating the observer's phase-state and thus their accessible "slice" of reality.

2.2 Fundamental Definitions

2.2.1 Universal Informational Field

We define a universal complex-valued informational field over extended spacetime:

$$\Psi : \mathcal{M}^4 \times \mathbb{R}^n \rightarrow \mathbb{C}, \quad \Psi = \Psi(x^\mu, \phi_j)$$

where:

- $x^\mu = (t, x, y, z) \in \mathcal{M}^4$: standard spacetime coordinates
- $\phi_j \in \mathbb{R}^n$: phase-state dimensions encoding kinematic, quantum, thermodynamic, and gravitational observer properties

2.2.2 Observer Phase-State Vector

Each observer is characterized by a time-dependent phase-state vector:

$$\Phi_{\text{obs}}(t) = \begin{bmatrix} \phi_1(t) \\ \phi_2(t) \\ \vdots \\ \phi_n(t) \end{bmatrix}$$

The kinematic phase component is given by:

$$\phi_{\text{kin}}(v) = \phi_{\text{kin}}^{(0)} + \int_0^v \beta(u) du$$

with $\beta(u) \approx u/c^2$ for $u \ll c$.

2.2.3 Reality Projection Operator

Observed reality emerges as a projection from the universal field:

$$R(x^\mu) = P[\Psi, \Phi_{\text{obs}}] = \int_{\mathbb{R}^n} \Psi(x^\mu, \phi) \cdot K(\phi | \Phi_{\text{obs}}(t)) d^n \phi$$

The projection kernel is defined as:

$$K(\phi | \Phi) = \exp \left(i \sum_{j=1}^n \phi_j \cdot F_j(x^\mu) \right)$$

where $F_j(x^\mu)$ are field operators coupling phase dimensions to spacetime events.

2.3 Field Dynamics

2.3.1 Action Functional

The universal field Ψ obeys an action principle:

$$S[\Psi] = \int d^4x d^n \phi \left[\frac{1}{2} |\partial_\mu \Psi|^2 - \frac{1}{2} |\partial_{\phi_j} \Psi|^2 - V(\Psi) \right]$$

2.3.2 Field Equation

Variation of the action yields the generalized informational Klein-Gordon equation:

$$\boxed{\square\Psi - \nabla_\phi^2\Psi + \frac{\delta V}{\delta\Psi^*} = 0}$$

where:

$$\square = \partial^\mu\partial_\mu = \frac{\partial^2}{\partial t^2} - \nabla^2$$

$$\nabla_\phi^2 = \sum_{j=1}^n \frac{\partial^2}{\partial\phi_j^2}$$

2.4 Modified Simultaneity Relation

For an observer moving with velocity $\vec{v}$, their effective simultaneity time for a distant event at position $\vec{r}$ is:

$$\boxed{t'_{\text{effective}} = t_0 - \frac{\vec{v} \cdot \vec{r}}{c^2} - \frac{\Delta\phi_{\text{kin}}}{\omega_0}}$$

The phase shift is:

$$\Delta\phi_{\text{kin}} = \phi_{\text{kin}}(v) - \phi_{\text{kin}}^{(0)}$$

and ω_0 is a characteristic universal frequency scale.

Physical interpretation: The first term is the standard Lorentz simultaneity shift; the second term represents the novel PSP correction due to observer phase-state modulation.

2.5 Quantum Correspondence

The standard quantum mechanical wavefunction emerges as a marginal projection:

$$\psi(x^\mu) = \int \Psi(x^\mu, \phi) \cdot \rho(\phi) d^n\phi$$

where $\rho(\phi)$ is the observer-specific phase distribution function.

2.6 Constraints on Projection Freedom

To preserve objective reality and prevent arbitrary manipulation of simultaneity:

$$\left| \frac{d\phi_{\text{kin}}}{dt} \right| \leq \omega_c \quad (\text{phase coherence bandwidth constraint}) \quad (1)$$

$$|\nabla_\phi R(x^\mu)| \leq \Lambda_{\text{coh}} \quad (\text{causal projection constraint}) \quad (2)$$

These constraints ensure:

- Phase-state changes are limited by fundamental coherence timescales
- Observer projections remain causally consistent
- Arbitrary "construction of reality" via engineered ϕ -fields is prevented

2.7 Experimental Predictions

2.7.1 Enhanced Simultaneity Shift

The apparent simultaneity difference between two observers includes both relativistic and phase-state contributions:

$$\Delta t_{\text{apparent}} = -\frac{vd}{c^2} - \frac{\Delta\phi_{\text{kin}}}{\omega_0}$$

For precision measurements, the second term may be detectable in:

- Atomic interferometry with controlled velocity modulation
- Quantum entanglement timing measurements between moving platforms
- Gravitational wave detector phase correlations

2.7.2 Observable Quantities

The kinematic phase shift can be operationally defined via detector-field coupling (Unruh-DeWitt formalism):

$$\Delta\phi_{\text{kin}} = \arg [\langle 0|U(t)|\Phi_{\text{obs}}\rangle] - \arg [\langle 0|U(t)|\Phi_0\rangle]$$

where $U(t)$ is the system evolution operator and $|\Phi_0\rangle$ represents a reference rest-frame observer state.

3 Comparison: Classical SR vs PSP Model

Feature	Special Relativity	PSP Model
Simultaneity	Frame-dependent geometric	Phase-dependent informational
Observer	Passive coordinate frame	Active information projector
Wavefunction	Abstract probability amplitude	Slice of universal field Ψ
Motion	Alters coordinates	Alters phase-state $\rightarrow$ reality access
QM-Relativity Link	Largely unresolved	Unified via projection dynamics
Testability	High (confirmed)	Medium (requires phase detection)

4 Conclusions

The Phase-State Projection model:

1. Provides a mechanistic explanation for relativistic simultaneity
2. Subsumes special relativity as a geometric limit
3. Unifies quantum measurement and relativistic observation
4. Makes novel testable predictions via phase-state corrections
5. Constrains observer freedom to prevent paradoxes

The Andromeda Simultaneity Problem, rather than being merely a curious consequence of Lorentz transformations, reveals the deeper structure of reality as an informational field accessible through observer-dependent phase projections.

Author: Branden Lee Friend

Framework: Phase-State Projection (PSP) Model