

Assignment #4

題目範圍: section 4.5~4.9

1. Show that the following polynomials form a basis for P_3 .
 $1+x, 1-x, 1-x^2, 1-x^3$
2. In each part, let $T_A : R^3 \rightarrow R^3$ be multiplication by A , and let $u = (1, -2, -1)$. Find the coordinate vector of $T_A(u)$ relative to the basis $S = \{(1, 1, 0), (0, 1, 1), (1, 1, 1)\}$ for R^3 .
 - a. (5 points) $A = \begin{bmatrix} 2 & -1 & 0 \\ 1 & 1 & 1 \\ 0 & -1 & 2 \end{bmatrix}$
 - b. (5 points) $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$
3. In each part, find a basis for the given subspace of R^3 , and state its dimension.
 - a. (2.5 points) The plane $3x - 2y + 5z = 0$.
 - b. (2.5 points) The plane $x - y = 0$.
 - c. (2.5 points) The line $x = 2t, y = -t, z = 4t$.
 - d. (2.5 points) All vectors of the form (a, b, c) , where $b = a + c$.
4. In each part, let T_A be multiplication by A and find the dimension of the subspace R^4 consisting of all vectors x for which $T_A(x) = 0$.
 - a. (5 points) $A = \begin{bmatrix} 1 & 0 & 2 & -1 \\ -1 & 4 & 0 & 0 \end{bmatrix}$
 - b. (5 points) $A = \begin{bmatrix} 0 & 0 & 1 & 1 \\ -1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 \end{bmatrix}$
5. Consider the bases $B = \{\mathbf{u}_1, \mathbf{u}_2\}$ and $B' = \{\mathbf{u}'_1, \mathbf{u}'_2\}$ for R^2 , where
$$\mathbf{u}_1 = \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 4 \\ -1 \end{bmatrix}, \mathbf{u}'_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \mathbf{u}'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$
 - a. (2.5 points) Find the transition matrix from B' to B .
 - b. (2.5 points) Find the transition matrix from B to B' .
 - c. (2.5 points) Compute the coordinate vector $[\mathbf{w}]_B$, where $\mathbf{w} = \begin{bmatrix} 3 \\ -5 \end{bmatrix}$ and use (11) to compute $[\mathbf{w}]_{B'}$.
 - d. (2.5 points) Check your work by computing $[\mathbf{w}]_{B'}$ directly.

Note: use(11) please to check the reference at the bottom of this assignment.

6. If B_1 , B_2 , and B_3 are bases for R^2 , and if

$$P_{B_1 \rightarrow B_2} = \begin{bmatrix} 3 & 1 \\ 5 & 2 \end{bmatrix} \quad \text{and} \quad P_{B_2 \rightarrow B_3} = \begin{bmatrix} 7 & 2 \\ 4 & -1 \end{bmatrix},$$

then $P_{B_3 \rightarrow B_1} =$ _____.

7. Suppose that $x_1 = -1$, $x_2 = 2$, $x_3 = 4$, $x_4 = -3$ is a solution of a nonhomogeneous linear system $A\mathbf{x} = \mathbf{b}$ and that the solution set of the homogeneous system $A\mathbf{x} = \mathbf{0}$ is given by the formulas

$$x_1 = -3r + 4s, \quad x_2 = r - s, \quad x_3 = r, \quad x_4 = s$$

- a.(5 points) Find a vector form of the general solution of $A\mathbf{x} = \mathbf{0}$.
b.(5 points) Find a vector form of the general solution of $A\mathbf{x} = \mathbf{b}$

8. Consider the linear systems

$$\begin{bmatrix} 1 & -2 & 3 \\ 2 & 1 & 4 \\ 1 & -7 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

and

$$\begin{bmatrix} 1 & -2 & 3 \\ 2 & 1 & 4 \\ 1 & -7 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 7 \\ -1 \end{bmatrix}$$

- a. (2.5 points) Find a general solution of the homogeneous system.
b.(2.5 points) Confirm that $x_1 = 1$, $x_2 = 1$, $x_3 = 1$ is a solution of the nonhomogeneous system.
c. (2.5 points) Use the results in parts (a) and (b) to find a general solution of the nonhomogeneous system.
d. (2.5 points) Check your result in part (c) by solving the nonhomogeneous system directly

9. Let $T : R^5 \rightarrow R^3$ be the linear transformation defined by the formula

$$T(x_1, x_2, x_3, x_4, x_5) = (x_1 + x_2, x_2 + x_3 + x_4, x_4 + x_5)$$

- a. (5 points) Find the rank of the standard matrix for T .
b. (5 points) Find the nullity of the standard matrix for T .

- 10.a. (2.5 points) If A is a 3×5 matrix, then the rank of A is at most _____.

Why?

b. (2.5 points) If A is a 3×5 matrix, then the nullity of A is at most _____.

Why?

c. (2.5 points) If A is a 3×5 matrix, then the rank of A^T is at most _____.

Why?

d. (2.5 points) If A is a 3×5 matrix, then the nullity of A^T is at most _____. Why?

評分標準：

每題 10 分，每小題配分已標注，答錯即 0 分。

每題都需寫心得，不單獨算分，但缺一題心得-2 point, 最多-20.

繳交期限：11/8（週二）00:00 遲交分數*0.8

Reference:

Transforming Coordinates

Suppose now that B and B' are bases for a finite-dimensional vector space V . Since multiplication by $P_{B \rightarrow B'}$ maps coordinate vectors relative to the basis B into coordinate vectors relative to a basis B' , and $P_{B' \rightarrow B}$ maps coordinate vectors relative to B' into coordinate vectors relative to B , it follows that for every vector \mathbf{v} in V we have

$$[\mathbf{v}]_{B'} = P_{B \rightarrow B'} [\mathbf{v}]_B \quad (11)$$

$$[\mathbf{v}]_B = P_{B' \rightarrow B} [\mathbf{v}]_{B'} \quad (12)$$