

PHY 831: Statistical Mechanics

Homework 6

Due Friday Nov 12, 2021

1. Consider a gas of massless identical particles with spin degeneracy g in three dimensions. Assuming $\mu = 0$, derive $P = AgT^4$ and $E/V = BgT^4$ and evaluate A and B for a) Fermions, and b) Bosons.
2. **Stoner ferromagnetism problem.** In a metal, the conduction electrons can be treated as a gas with an effective spin-spin interaction energy that mocks up the effects of the Coulomb interactions. More precisely, the Coulomb repulsion amongst the conduction electrons favors a *spatial* wave function that is antisymmetric in the coordinates, so that the electrons are kept from overlapping strongly. Since the *full* wave function (spatial and spin variables) is antisymmetric, this interaction can be approximated by an effective spin-spin coupling that favors states with parallel spins. In this simple approximation, the net effect is described by an interaction energy

$$U = \alpha \frac{N_+ N_-}{V},$$

where N_{\pm} represent the number of spin-up and spin-down electrons, and α is a parameter related to the zero-energy scattering amplitude between two electrons.

- (a) The $T = 0$ ground state consists of two fermi seas filled by the spin-up and spin-down electrons. Express the corresponding Fermi wave vectors k_F^{\pm} in terms of the densities $n_{\pm} = N_{\pm}/V$.
 - (b) Calculate the kinetic energy density of the ground state as a function of n_{\pm} and various fundamental constants.
 - (c) Assuming small deviations $n_{\pm} = n/2 \pm \delta$ from the symmetric state, expand the kinetic energy to fourth-order in the small parameter δ .
 - (d) Express the spin-spin interaction density U/V in terms of n and δ . Find the critical value of α_c such that for $\alpha > \alpha_c$, the electron gas can lower its total energy by developing a magnetization. (This is known as the Stoner instability.)
 - (e) Explain qualitatively and sketch the behavior of the spontaneous magnetization as a function of α .
3. **A generalized quantum ideal gas.** Consider a gas of non-interacting identical spinless quantum particles in d -dimensions with a weird single particle energy dispersion $\epsilon_p = \alpha p^s$, where α is some constant so that ϵ_p has dimensions of energy.

- (a) Calculate the grand potential and density at a chemical potential of μ , expressing your answer in terms of s , d , and $f_m^\eta(z)$ where $z = e^{\beta\mu}$.
 - (b) Find the ratio of PV/E .
 - (c) For fermions, calculate the dependence of E/N and P on the density $n = N/V$ at zero temperature. (Hint: Recall that $f_m^-(z) \rightarrow (\log z)^m/m!$ as $z \rightarrow \infty$.)
 - (d) For Bosons, find the dimension $d(s)$ below which there is no Bose-Einstein condensation. Is there condensation at $d = 2$ for $s = 2$? Why or why not?
4. Consider an ultrarelativistic gas of bosons in d -dimensions. By mimicking what we did in class for the non-relativistic gas, find the lower critical dimension— i.e., highest dimension d_c for which N diverges as you take $\mu \rightarrow 0^-$ indicating no macroscopic condensation in the lowest single particle level. For $d > d_c$, there is condensation. Find an expression for the critical temperature $T_c(N, V)$.