

Problem Set 2

Brandon Henke¹

¹Michigan State University - Department of Physics & Astronomy

Last updated: January 31, 2023

I PERTURBATIVE HYDROGEN FINE STRUCTURE

A

B

C

II SAKURAI 5.6

The energy eigenstates of the infinite square well in two dimensions are:

$$\psi(\mathbf{r}) = \frac{2}{L} \sin\left(x \frac{n_x \pi}{L}\right) \sin\left(y \frac{n_y \pi}{L}\right). \quad (\text{II.1})$$

The energies of these states are

$$E_{n_x, n_y} = \frac{\hbar^2}{2m} \left(\frac{(n_x + n_y)\pi}{L} \right)^2 \quad (\text{II.2})$$

For a total $N = n_x + n_y$, there are $N + 1$ states with the same energy. The ground state is $n_x = n_y = 1$, and the first excited state is $n_x = 2$ and $n_y = 1$ or vice versa. Denote the energy states as $|n_x, n_y\rangle$.

$$\langle 1, 1 | \hat{V}_1 | 1, 1 \rangle = \lambda \langle 1, 1 | \hat{x} \hat{y} | 1, 1 \rangle, \quad (\text{II.3})$$

$$= \lambda \frac{L^2}{4}. \quad (\text{II.4})$$

$$V = \lambda \begin{pmatrix} \langle 1, 2 | xy | 1, 2 \rangle & \langle 1, 2 | xy | 2, 1 \rangle \\ \langle 2, 1 | xy | 1, 2 \rangle & \langle 2, 1 | xy | 2, 1 \rangle \end{pmatrix}, \quad (\text{II.5})$$

$$= \frac{\lambda L^2}{4\pi^4} \begin{pmatrix} \pi^4 & 1024/81 \\ 1024/81 & \pi^4 \end{pmatrix}, \quad (\text{II.6})$$

$$\therefore \Delta_{1\pm}^{(1)} = \frac{1}{81} (81\pi^4 \pm 1024). \quad (\text{II.7})$$

The eigenvectors (zeroth-order energy eigenkets) are

$$|\Delta_{1\pm}^{(1)}\rangle = \frac{1}{\sqrt{2}} (|1, 2\rangle \pm |2, 1\rangle). \quad (\text{II.8})$$

III SAKURAI 5.19

$$H = \begin{pmatrix} E_2 + h\delta & 3ea_0\mathcal{E} \\ 3ea_0\mathcal{E} & E_2 \end{pmatrix}. \quad (\text{III.1})$$

The eigenvalues, E , are

$$E_{\pm} = -\frac{h\delta}{2} \pm \frac{1}{2} \sqrt{h^2\delta^2 + 36e^2\mathcal{E}^2 a_0^2} + E_2. \quad (\text{III.2})$$

Taking the limit where $\delta \ll 3ea_0\mathcal{E}$,

$$E_{\pm} \approx -\frac{h\delta}{2} \pm 3ea_0\mathcal{E} + E_2, \quad (\text{III.3})$$

which is indeed linear in \mathcal{E} .

In the limit where $\delta \gg 3ea_0\mathcal{E}$,

$$E_{\pm} = -\frac{h\delta}{2} \pm \frac{1}{2} \left(h\delta + \frac{9e^2 a_0^2 \mathcal{E}^2}{2h\delta} \right) + E_2, \quad (\text{III.4})$$

which is quadratic in \mathcal{E} .