

# Complex Analysis Chapter 1 Section 2

Brandyn Tucknott

Last Updated: 24 September 2025

## 2 Functions on the Complex Plane

### 2.1 Continuous Functions

Let  $f$  be a function on a set  $\Omega$  of complex numbers. We say that  $f$  is **continuous** at a point  $z_0 \in \Omega$  if for every  $\epsilon > 0$  there exists a  $\delta > 0$  such that whenever  $z \in \Omega$  and  $|z - z_0| < \delta$  then  $|f(z) - f(z_0)| < \epsilon$ . Equivalently, we can say for every sequence  $\{z_1, z_2, \dots\} \subset \Omega$  such that  $\lim z_n = z_0$ , then  $\lim f(z_n) = f(z_0)$ . The function  $f$  is continuous on  $\Omega$  if it is continuous at every point in  $\Omega$ . Sums and products of continuous functions are also continuous.

It is worth noting that the function  $f$  of the complex argument  $z = x + iy$  is continuous if and only if it is continuous viewed as a function of the two real variables  $x, y$ .

By the triangle inequality, we see that if  $f$  is continuous, then the real-valued function defined by  $z \rightarrow |f(z)|$  is continuous. We say that  $f$  attains a **maximum** at a point  $z_0 \in \Omega$  if

$$|f(z)| \leq |f(z_0)| \text{ for all } z \in \Omega,$$

with the inequality reversed for the definition of a **minimum**.

**Theorem 2.1.** *A continuous function on a compact set  $\Omega$  attains a maximum and minimum on  $\Omega$ .*  $\square$

### 2.2 Holomorphic Functions

Let  $\Omega \subset \mathbb{C}$  be open and  $f$  a complex-valued function on  $\Omega$ . The function  $f$  is **holomorphic at the point**  $z_0 \in \Omega$  if

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}$$

converges. Here  $h \in \mathbb{C}$  and  $h \neq 0$  with  $z_0 + h \in \Omega$ , so that the quotient is well-defined. The limit of the quotient, when it exists, is denoted by  $f'(z_0)$  and is called the **derivative of  $f$  at  $z_0$** :

$$f'(z_0) = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}.$$

Take note that  $h$  is complex and can approach 0 from any direction.

The function  $f$  is **holomorphic on  $\Omega$**  if it is holomorphic at every point of  $\Omega$ . If  $C$  is a closed subset of  $\mathbb{C}$ , we say that  $f$  is **holomorphic on  $C$**  if  $f$  is holomorphic in some open set containing  $C$ . If  $f$  is holomorphic on  $\mathbb{C}$ , we say that  $f$  is **entire**.

**Proposition 2.2.** *If  $f$  and  $g$  are holomorphic in  $\Omega$ , then:*

- $f + g$  is holomorphic in  $\Omega$  and  $(f + g)' = f' + g'$ .
- $fg$  is holomorphic in  $\Omega$  and  $(fg)' = f'g + fg'$ .
- If  $g(z_0) \neq 0$ , then  $f/g$  is holomorphic at  $z_0$  and

$$(f/g)' = \frac{gf' - fg'}{g^2}.$$

Moreover, if  $f : \Omega \rightarrow U$  and  $g : U \rightarrow \mathbb{C}$  are holomorphic, then the chain rule holds;

$$(g \circ f)'(z) = g'(f(z))f'(z) \text{ for all } z \in \Omega.$$

## Complex-Valued Functions as Mappings

Recall that a function  $F(x, y) = (u(x, y), v(x, y))$  is said to be differentiable at a point  $P_0 = (x_0, y_0)$  if there exists a linear transformation  $J : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that

$$\frac{|F(P_0 + H) - F(P_0) - J(H)|}{|H|} \rightarrow 0 \text{ as } |H| \rightarrow 0, H \in \mathbb{R}^2$$

Equivalently we can write

$$F(P_0 + H) - F(P_0) = J(H) + |H| \Psi(H),$$

with  $|\Psi(H)| \rightarrow 0$  as  $H \rightarrow 0$ . The linear transformation  $J$  is unique and is called the derivative of  $F$  at  $P_0$ . If  $F$  is differentiable, the partial derivatives of  $u$  and  $v$  exist and  $J$  is described with the standard basis in  $\mathbb{R}^2$  by the Jacobian of  $F$

$$J = J_F(x, y) = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix}.$$

With complex-differentiation, the derivative is a complex number  $f'(z_0)$ , and with the reals it is a matrix. However, there is a relation to be found involving the partials of  $u$  and  $v$ .

Given the following equations

$$\frac{f(z_0 + h) - f(z_0)}{h} \tag{1}$$

$$f(z_0 + h) - f(z_0) - ah = h\psi(h) \tag{2}$$

$$\frac{|F(P_0 + H) - F(P_0) - J(H)|}{|H|} \rightarrow 0 \text{ as } |H| \rightarrow 0 \tag{3}$$

consider the limit when  $h$  is real. That is,  $h = h_1 + ih_2$  with  $h_2 = 0$ . Then if we write  $z = x + iy$ ,  $z_0 = x_0 + iy_0$ , and  $f(z) = f(x, y)$ , we find that

$$\begin{aligned} f'(z_0) &= \lim_{h_1 \rightarrow 0} \frac{f(x + h_1, y_0) - f(x_0, y_0)}{h_1} \\ &= \frac{\partial f}{\partial x}(z_0). \end{aligned}$$

Now taking  $h$  to be purely imaginary with  $h = ih_2$ , a similar argument shows that

$$\begin{aligned} f'(z_0) &= \lim_{h_2 \rightarrow 0} \frac{f(x_0, y_0 + h_2) - f(x_0, y_0)}{ih_2} \\ &= \frac{1}{i} \frac{\partial f}{\partial y}(z_0). \end{aligned}$$

Therefore, if  $f$  is holomorphic, we have shown that

$$\frac{\partial f}{\partial x} = \frac{1}{i} \frac{\partial f}{\partial y}.$$

Writing  $f = u + iv$  after separating real and imaginary parts as well as using  $1/i = -i$ , we find that the partials of  $u$  and  $v$  exist, and they satisfy the following relations:

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

These are the **Cauchy-Riemann equations**. We can take this further and define two differential operators:

$$\frac{\partial}{\partial z} = \frac{1}{2} \left( \frac{\partial}{\partial x} + \frac{1}{i} \frac{\partial}{\partial y} \right) \text{ and } \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial}{\partial x} - \frac{1}{i} \frac{\partial}{\partial y} \right).$$

**Proposition 2.3.** *If  $f$  is holomorphic at  $z_0$ , then*

$$\frac{\partial f}{\partial \bar{z}} = 0 \text{ and } f'(z_0) = \frac{\partial f}{\partial z}(z_0) = 2 \frac{\partial u}{\partial z}(z_0).$$

*Also, if we write  $F(x, y) = f(z)$ , then  $F$  is differentiable in the sense of real variables, and*

$$\det J_F(x_0, y_0) = |f'(z_0)|^2.$$

*Proof.* Taking real and imaginary parts, it is easy to see the Cauchy-Riemann equations are equivalent to  $\partial f / \partial \bar{z} = 0$ . By our earlier observation

$$\begin{aligned} f'(z_0) &= \frac{1}{2} \left( \frac{\partial f}{\partial x}(z_0) + \frac{1}{i} \frac{\partial f}{\partial y}(z_0) \right) \\ &= \frac{\partial f}{\partial z}(z_0) \end{aligned}$$

and the Cauchy-Riemann equations give  $\partial f / \partial z = 2 \partial u / \partial z$ . To prove that  $F$  is differentiable, it suffices to show that if  $H = (h_1, h_2)$  and  $h = h_1 + ih_2$ , then the Cauchy-Riemann equations imply

$$J_F(x_0, y_0)(H) = \left( \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} \right) (h_1 + ih_2) = f'(z_0)h,$$

where we have identified a complex number with the pair of real and imaginary parts. Another application of the Cauchy-Riemann equations give

$$\det J_F(x_0, y_0) = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial v}{\partial x} \frac{\partial u}{\partial y} = \left( \frac{\partial u}{\partial x} \right)^2 + \left( \frac{\partial v}{\partial y} \right)^2 = \left| 2 \frac{\partial u}{\partial z} \right|^2 = |f'(z_0)|^2.$$

□

**Theorem 2.4.** *Suppose  $f = u + iv$  is a complex-valued function defined on an open set  $\Omega$ . If  $u$  and  $v$  are continuously differentiable and satisfy the Cauchy-Riemann equations on  $\Omega$ , then  $f$  is holomorphic on  $\Omega$  and  $f'(z) = \partial f / \partial z$ .*

*Proof.* Write

$$u(x + h_1, y + h_2) - u(x, y) = \frac{\partial u}{\partial x} h_1 + \frac{\partial u}{\partial y} h_2 + |h| \psi_1(h)$$

and

$$v(x + h_1, y + h_2) - v(x, y) = \frac{\partial v}{\partial x} h_1 + \frac{\partial v}{\partial y} h_2 + |h| \psi_2(h),$$

where  $\psi_j(h) \rightarrow 0$  (for  $j = 1, 2$ ) as  $|h| \rightarrow 0$  and  $h = h_1 + ih_2$ . Using the Cauchy-Riemann equations, we find that

$$f(z+h) - f(z) = \left( \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} \right) (h_1 + ih_2) + |h|\psi(h),$$

where  $\psi(h) = \psi_1(h) + \psi_2(h) \rightarrow 0$  as  $|h| \rightarrow 0$ . Therefore  $f$  is holomorphic and

$$f'(z) = 2 \frac{\partial u}{\partial z} = \frac{\partial f}{\partial z}.$$

□

## 2.3 Power Series

A classic example of a power series is the complex **exponential** function, defined for all  $z \in \mathbb{C}$  by

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

The above series converges absolutely for all  $z \in \mathbb{C}$ , which can be seen by observing

$$\left| \frac{z^n}{n!} \right| = \frac{|z|^n}{n!},$$

which allows us to compare  $|e^z|$  with the power series  $\sum_{n=0}^{\infty} |z|^n / n! = e^{|z|} < \infty$ . Moreover, this shows that the power series of  $e^z$  converges uniformly in all discs in  $\mathbb{C}$ .

**Theorem 2.5.** *Given a power series  $\sum_{n=0}^{\infty} z^n$ , there exists  $0 \leq R \leq \infty$  such that:*

- *If  $|z| < R$ , the series converges absolutely.*
- *If  $|z| > R$ , the series diverges.*

Moreover, if we use the convention that  $1/0 = \infty$  and  $1/\infty = 0$ , then  $R$  is given by Hadamard's formula

$$\frac{1}{R} = \limsup |a_n|^{1/n}.$$

**Remark.** Notice this theorem says nothing about when  $|z| = R$ . This situation can either converge or diverge.

The number  $R$  is called the **radius of convergence** of the power series, and the region  $|z| < R$  the **disc of convergence**. Moreover, the radius of convergence for the exponential function and geometric series are  $R = \infty$  and  $R = 1$  respectively.

More examples of power series in the complex plane are given below in the form of **trigonometric functions**. These are defined as

$$\cos z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!} \quad \text{and} \quad \sin z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!}.$$

A connection between these and the exponential function can be expressed as

$$\cos z = \frac{e^{iz} + e^{-iz}}{2} \quad \text{and} \quad \sin z = \frac{e^{iz} - e^{-iz}}{2i}.$$

These are called **Euler formulas** for the sine and cosine functions.

**Theorem 2.6.** *The power series  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  defines a holomorphic function in its disc of convergence. The derivative of  $f$  is also a power series obtained by differentiating the term by term power series for  $f$ , that is,*

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}.$$

Moreover,  $f'$  has the same radius of convergence as  $f$ .

*Proof.* Since  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ , by Hadamard's formula we know that

$$\frac{1}{R} = \lim_{n \rightarrow \infty} |n|^{1/n} = \lim_{n \rightarrow \infty} |n a_n|^{1/n},$$

thus  $\sum a_n z^n$  and  $\sum n a_n z^n$  have the same radius of convergence.

To prove the first assertion, we must show that

$$g(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}$$

gives the derivative of  $f$ .

Let  $R$  denote the radius of convergence of  $f$ , and let  $|z_0| < r < R$ . Denote

$$S_N(z) = \sum_{n=0}^N a_n z^n \text{ and } E_N(z) = \sum_{n=N+1}^{\infty} a_n z^n$$

and write

$$f(z) = S_N(z) + E_N(z).$$

If  $h$  is chosen such that  $|z_0 + h| < r$ , we have

$$\frac{f(z_0 + h) - f(z_0)}{h} - g(z_0) = \left( \frac{S_N(z_0 + h) - S_N(z_0)}{h} - S'_N(z_0) \right) + (S'_N(z_0) - g(z_0)) + \left( \frac{E_N(z_0 + h) - E_N(z_0)}{h} \right).$$

Since  $a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + \dots + ab^{n-2} + b^{n-1})$ , we see that

$$\left| \frac{E_N(z_0 + h) - E_N(z_0)}{h} \right| \leq \sum_{n=N+1}^{\infty} |a_n| \left| \frac{(z_0 + h)^n - z_0^n}{h} \right| \leq \sum_{n=N+1}^{\infty} |a_n| n r^{n-1},$$

where we use the fact that  $|z_0| < r$  and  $|z_0 + h| < r$ . The expression on the right is the tail end of a convergent series, since  $g$  converges absolutely on  $|z| < R$ . Therefore, given  $\varepsilon > 0$ , we can find  $N_1$  such that  $N > N_1$  implies

$$\left| \frac{E_N(z_0 + h) - E_N(z_0)}{h} \right| < \varepsilon.$$

Also, since  $\lim_{n \rightarrow \infty} S'_N(z_0) = g(z_0)$ , we can find  $N_2$  such that  $N > N_2$  implies

$$|S'_N(z_0) - g(z_0)| < \varepsilon.$$

If we fix  $N$  so that  $N > N_1, N_2$ , we can find  $\delta > 0$  so that  $|h| < \delta$  implies

$$\left| \frac{S_N(z_0 + h) - S_N(z_0)}{h} - S'_N(z_0) \right| < \varepsilon,$$

since the derivative of a polynomial is obtained by differentiating it term by term. Therefore

$$\left| \frac{f(z_0 + h) - f(z_0)}{h} - g(z_0) \right| < 3\varepsilon$$

whenever  $|h| < \delta$ , thereby concluding the proof of the theorem.  $\square$

**Corollary 2.7.** *A power series is infinitely complex differentiable in its disc of convergence, and the higher derivatives are also obtained by term-wise differentiation.*  $\square$

So far, we have only dealt with power series centered about the origin, but we more generally express a power series centered at  $z_0 \in \mathbb{C}$  as

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n.$$

The disc of convergence of  $f$  is now centered at  $z_0$ , but the radius is still given by Hadamard's formula. In fact, if

$$g(z) = \sum_{n=0}^{\infty} a_n z^n,$$

$f$  is obtained by simply translating  $g$ .

A function  $f$  defined on an open set  $\Omega$  is said to be **analytic** (having a power series expansion) at a point  $z_0 \in \Omega$  if there exists a power series  $\sum a_n (z - z_0)^n$  centered at  $z_0$  with positive radius of convergence such that

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n \text{ for all } z \text{ in a neighborhood of } z_0.$$

If  $f$  has a power series expansion point at every point in  $\Omega$ , we say that  $f$  is **analytic on  $\Omega$** . By Theorem 2.6, an analytic function is also holomorphic. In the future we will prove the converse is true: every holomorphic function is analytic. For that reason we use the terms holomorphic and analytic interchangeably.

## 2.4 Integration along curves

pg 38