


# FORECASTING LIPID BINDING AND MODULATION OF AN ION CHANNEL FROM SIMULATION



EZRY SANTIAGO-MCRAE<sup>1</sup>, THOMAS T. JOSEPH<sup>2</sup>, WAYLAND CHENG<sup>3</sup>, JÉRÔME HÉNIN<sup>4</sup>, GRACE BRANNIGAN<sup>1,5</sup>

1) CENTER FOR COMPUTATIONAL AND INTEGRATIVE BIOLOGY, RUTGERS UNIVERSITY, CAMDEN 2) DEPT. OF ANESTH. CRIT. CARE, UNIVERSITY OF PENNSYLVANIA PERELMAN SCHOOL OF MEDICINE  
3) SCHOOL OF MEDICINE IN ST. LOUIS, WASHINGTON UNIVERSITY 4) LABORATOIRE DE BIOCHIMIE THÉORIQUE, CNRS 5) CAMDEN, DEPT. PHYSICS, RUTGERS UNIVERSITY




## ABSTRACT

Membrane proteins are very sensitive (>5x gain of function) to their local lipid environment. The mechanisms that cause this sensitivity are not well understood. Our work involves computationally modeling a membrane protein to study how specific lipid binding may cause this sensitivity. Our system of interest is Erwinia ligand-gated ion channel (ELIC), a bacterial homolog of many neuronal proteins including n-acetylcholine receptor and GABA<sub>A</sub> receptor. We have extended free energy perturbation (FEP) to phospholipid binding in a methodology we call SAFE. From these simulations, we have been able to estimate the binding free energy of POPG to ELIC as well as make predictions of POPG modulation at a single-protein level. Our results are consistent with, and clarify, the experimental data available.

## LIPID DIVERSITY AFFECTS MEMBRANE FUNCTION<sup>1</sup>

- Cell membranes have diverse compositions
- Membrane lipids differ widely




**Fig 1: Lipid Diversity.** Phospholipids can differ in their tail length and saturation (e.g. oleoyl, palmitoyl), or in their head group chemistry (e.g. choline, glycerol). POPC is a common mammalian lipid, while POPG is more common in bacterial membranes.

- Many membrane proteins have specific membrane requirements:
  - Cytochromes
  - ATPases
  - Ion Channels

## pLGICs ARE ESSENTIAL NEURONAL PROTEINS<sup>2</sup>

- Pentameric Ligand-gated Ion Channels
- Examples include<sup>2</sup>:
  - N-Acetylcholine receptor
  - GABA<sub>A</sub> receptor
  - 5-HT<sub>3</sub> (serotonin) receptor
- Key role in neuron-neuron communication<sup>2</sup>:
  - Reside in post-synaptic membrane
  - Detect neurotransmitters
  - Initiate depolarization (action potential)
- ELIC is a bacterial model pLGIC [Fig 3]<sup>2,3</sup>


## ELIC IS SENSITIVE TO POPG



**Fig 2: ELIC is sensitive to POPG concentration.** Patch clamp recordings of ELIC in a POPC model membrane normalized to peak current. As POPG concentration is increased, desensitization is delayed. Peak currents also increase (Data not shown). Adapted from citation 4.


## OPEN OR CLOSED? WHEN AND WHY?

### ELIC'S PORE CAN BE OPEN OR CLOSED TO IONS



**Fig 3: Cross-sections of the open and closed states of ELIC.** After 600ns of simulation, the closed state (left) maintains a narrow pore with little hydration (blue surface). The open state (right), in contrast, is well-hydrated with a wide pore.

### THE OPEN STATE HAS A HIGHER AFFINITY FOR POPG




**Fig 4: Site occupancy by POPG in a primarily POPC membrane.** Occupancy probability of ELIC5/open (blue) and wild-type (orange). Shaded regions indicate 95% confidence intervals.

**Eqn 1:** Occupancy probability as a function of POPG mole fraction and binding affinity.

$$p_{occ} = \frac{x_{PG}}{e^{-\frac{\Delta G_{bind}}{RT}} + x_{PG}}$$

- Binding affinity for each state was computed by SAFE simulations
- The open state has a much higher affinity for POPG than the closed state

### POPG INCREASES THE RELATIVE OPEN PROBABILITY BY UP TO 30x




**Fig 5: Log modulation of ELIC versus mole fraction of POPG.** Calculated by eqn 2. Greater values correspond to gain of function. Shaded region indicates 95% confidence interval.

**Eqn 2:** The relative open probability as a function of binding free energies.

$$\frac{p_{occ}^{open}}{p_{occ}^{closed}} = \frac{\Delta G_{bind}^{open} \cdot x_{PG} + 1}{\Delta G_{bind}^{closed} \cdot x_{PG} + 1}$$

- State-dependence suggests positive allosteric modulation
- Relative open probability is estimated by eqn 2
- Consistent with experiment:
  - Undetectable flux without POPG<sup>4</sup>
- Compare with Fig 2

## METHODS - SAFE



**Fig 6: The thermodynamic cycle for moving a POPG molecule from the bulk to the site.** POPG (red) is achemically removed from each state: bound (left) and bulk (right). The free energy of each transformation is combined to estimate the free energy of transfer. Solid lines are computed directly. Dashed lines are derived.

$$\Delta G_{bind} = \Delta G_{bulk} + \Delta G_{restraints} + \Delta G_{DBC} - \Delta G_{site}$$

- A molecular dynamics system is transformed into another over a simulation
- The free energy of transformation can be computed<sup>5</sup>
- SAFE makes FEP in membranes possible:
  - Removes the need of a standard state
  - Introduces restraints that improve numerical convergence (not illustrated above)

## SUMMARY

### 1. Novel use of FEP on phospholipids via SAFE

### 2. Quantitative Predictions:

- Under physiologically relevant conditions
- Increasing POPG mol fraction
  - Increases site occupancy
- Affinity is state-dependent
  - Increases open probability
- Strong positive allostery
- Consistent with experimental results

## REFERENCES

- Yeagle. The Membranes of Cells. 2016
- Jaiteh, Taly, Hénin. PLOS ONE. 2014
- Petroff, [...], Santiago-McRae, [...] Hénin, Brannigan, Cheng. Nat Comms. 2022.
- Tong, [...], Brannigan, Cheng. eLife. 2019
- Salari, Joseph, Lohia, Hénin, Brannigan. JCTC. 2018.
- Santiago-McRae, Ebrahimi, Sandberg, Brannigan, Hénin. Live-CoMS. 2023

### SAFE PROTOCOL<sup>5</sup>



## ACKNOWLEDGMENTS

- Computation resources provided by the Rutgers Office of Advanced Research Computing and ACCESS (allocation BIO220103)
- Funding provided by NSF DGE2152059 & NIH K08 GM139031