

Understanding the Mechanisms of Decorated Nanoparticle Aggregation In Lipid Membranes



Jahmal Ennis¹, Jesse Sandberg¹, Ezry St.Iago-McRae¹, Julianne Griepenburg^{1,2}, and Grace Brannigan^{1,2}
¹Center for Computational and Integrative Biology and ²Department of Physics, Rutgers University, Camden, NJ



Abstract

Gold nanoparticles are a ubiquitous photosensitizer with a broad range of applications in microscopy and targeted drug delivery. Vesicles photosensitized with gold nanoparticles are promising as targeted drug delivery vehicles due to their non-invasive rupture mechanism. Controlling the vesicle properties increases control of the spatial-temporal release of cargo and drug dosage. The aggregation of gold nanoparticles affects the photoporation of these vesicles by interfering with peak SPR wavelengths. These nanoparticles cause leaflet bending deformations on the order of the nanoparticle size, while ligand chains may also disrupt local packing of lipid chains. Aggregation may be driven by the need to minimize one or both effects, but their relative contributions are unknown. In order to test the contributions of these two perturbations to membrane structure, we simulated multi-nanoparticle systems in lipid membranes of varying compositions. We used coarse-grained molecular dynamics simulation via the MARTINI forcefield to simulate simple spherical nanoparticles with a decorated ligand exterior. We found that large-scale nanoparticle aggregation depends more on ligand chain length than nanoparticle size, suggesting that aggregation of gold nanoparticles is dominated by microscopic perturbations to lipid packing.

Hydrophobic nanoparticle size regime determines aggregation mechanism




Fig 2. Visualization of a lipid membrane with hydrophobic 5nm nanoparticles. Lipids are in cyan, phosphate bead in orange, and nanoparticles in blue. Ligands and nanoparticle surface beads are not shown




Fig 4. Fraction of nanoparticle monomers in systems with varying NP diameters. Each system has 3 replicas that ran for 5 μ s, analysis was averaged over the final 1 μ s. Systems contain 10 nanoparticles, lipid-nanoparticle ratio is held constant.




Fig 5. Lipid order parameter dependence on nanoparticle ligand length. Systems contain single, 2nm nanoparticles and ran for 20 μ s, analysis was averaged over the final 1 μ s. A negative monotonic trend is observed as ligand length increases.




Fig 6. Fraction of nanoparticles in the largest aggregate in systems of varying ligand length. Each system has 3 replicas that ran for 5 μ s, analysis was averaged over the final 1 μ s. Systems contain sets of 10, 2nm nanoparticles held at constant lipid-nanoparticle ratio.

Background

- Ligand coated nanoparticles (NPs, Figure 1) are multipurpose tools used in biosensing, biolabelling, and controlled drug delivery¹




Fig 1. Visualization of 2 nm charged nanoparticles aggregating. Ligands in blue, nanoparticles in ochre and blue, lipid heads in green, lipids in grey.

- Nanoparticle aggregation in lipid membranes poses a challenge for many applications
- The mechanism of ligand coated nanoparticle aggregation is not well understood.
- We use coarse grained molecular dynamics to study the aggregation mechanism due to two types of deformations, membrane bending and acyl chain bending

Charge influences small NP aggregation




Fig 7. Solvent accessible surface area at varying ligand length in single nanoparticle systems. Systems contain 2nm nanoparticles and ran for 20 μ s, analysis was averaged over the final 5 μ s. A negative monotonic trend is observed with increasing ligand length.





Fig 8. Fraction of nanoparticles in the largest aggregate dependence on the ligand length of charged nanoparticles. System contain 10 nanoparticles held at a fixed ratio with lipids. 3 replicas of each system ran for 5 μ s, analysis was averaged over the final 1 μ s.



Summary

- Bare hydrophobic nanoparticle aggregation in lipid membranes has been shown in other works to be primarily driven by membrane deformations, however, ligand coated nanoparticle aggregation is driven by both elastic membrane deformations (large NP regime) and microscopic packing deformations.^{5,6}
- Nanoparticle aggregation is dependent on the degree of local lipid chain disorder.
- At short ligand lengths, the charged nanoparticle surface is more exposed.
- Charged nanoparticles coated by short (but not long) form larger aggregates than the equivalent hydrophobic nanoparticles, likely reflecting increased exposure.

Research Questions

- What is the primary mechanism of aggregation for charged and hydrophobic ligand coated nanoparticles in phosphatidylcholine membranes?
- Is the nanoparticle aggregation mechanism dependent on the nanoparticle regime?
- How do charged interactions affect nanoparticle aggregation?



Methods

- Simulation software: Gromacs 2016³, Force Field: Martini 2.2²
- POPC Membranes constructed with insane.py⁴
- # of NP's in largest aggregate (n)
- Total # of nanoparticles in the system (n_{tot})
- Total number of single nanoparticle in the system (n_s)

$$F_a = \frac{n}{n_{tot}} \quad M_f = \frac{n_s}{n_{tot}} \quad S = \frac{3\langle \cos^2 \theta \rangle - 1}{2}$$

Approach

- CG-MD simulations of hydrophobic and charged nanoparticles.
- Simulating nanoparticles at 4 different diameters to understand nanoparticle induced membrane deformation impact on aggregation.
- Simulating nanoparticles at 5 different ligand length to understand nanoparticle induced lipid deformations impact on aggregation.



References

- Vargas, K. M., & Shon, Y. S. (2019). Hybrid lipid-nanoparticle c complexes for biomedical applications. *Journal of materials chemistry. B*, 7(18), 695-708.
- The MARTINI Force Field: Coarse Grained Model for Biomolecular Simulations. Sievert J. Marrink, H. Jelger Risselada, Serge Yefimov, D. Peter Tieleman, and Alex H. de Vries. *The Journal of Physical Chemistry B* 2007 111 (27), 7812-7824.
- Mark James Abraham, Teemu Murtola, Roland Schulz, Szilard Páll, Jeremy C. Smith, Berk Hess, Erik Lindahl, GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* 1-2, 2015, Pages 19-25, ISSN 2352-7110
- Computational Lipidomics with insane: A Versatile Tool for Generating Custom Membranes for Molecular Simulations Tjerk A. Wassenaar, Helgi I. Ingólfsson, Rainer A. Böckmann, D. Peter Tieleman, and Sievert J. Marrink. *Journal of Chemical Theory and Computation* 2015 11 (5), 2144-2155
- Contini, Claudio and Hindley, James W. and Macdonald, Thomas J. and Barritt, Joseph D. and Ces, Oscar and Quirk, Nick. *Communications Chemistry* Size dependency of gold nanoparticles interacting with model membranes. 2020, 2399-3668, 1, 130, 3
- Lavagna, Enrico and Barnoud, Jonathan and Rossi, Giulia and Monticelli, Luca. *Nanoscale* Size-dependent aggregation of hydrophobic nanoparticles in lipid membranes. 2020, 2040-2049, 12, 9452-9461, 12

Acknowledgements

- Rutgers Office of Research and Computing (OARC)
- Pittsburgh Supercomputing Center (Bridges)
- NRT Award: NSF DGE 2152059