


Alchemy-based predictions for binding affinities of lipophilic ligands and membrane proteins: *obstacles, solutions, and outcomes*

Grace Brannigan
Center for Computational & Integrative Biology
Rutgers University - Camden




main target: pentameric gated ligand channels (pLGICs)



main target: pentameric gated ligand channels (pLGICs)



recreation





main target: pentameric gated ligand channels (pLGICs)



recreation



nicotine



propofol



pentobarbital



chloroform

general anesthetics



main target: pentameric gated ligand channels (pLGICs)



recreation



nicotine



propofol pentobarbital chloroform

general anesthetics



poisons

rat poison (TETS)

strychnine

curare

Fishberry (picrotoxin)



main target: pentameric gated ligand channels (pLGICs)



recreation



nicotine



propofol pentobarbital chloroform

general anesthetics

sedatives



benzodiazepines



methaqualone



rat poison (TETS)



strychnine



curare



Fishberry (picrotoxin)

poisons

pLGIcs : extremely lipid sensitive

pLGICs : extremely lipid sensitive

Most membrane proteins



pLGICs : extremely lipid sensitive

Most membrane proteins



pLGICs

pLGICs : extremely lipid sensitive

Most membrane proteins



pLGICs




pLGICs : extremely lipid sensitive

Most membrane proteins



pLGICs




pLGICs : extremely lipid sensitive

Most membrane proteins



pLGICs





Henin...Brannigan, 2014, Biophys J

pLGICs : extremely lipid sensitive

Most membrane proteins



pLGICs




pLGICs : extremely lipid sensitive


Most membrane proteins



pLGICs



Criado...Barrantes, 1984, J.Biol.Chem.



General Motivation

1. Lipid-protein interactions: specific binding of

- cholesterol
- polyunsaturated fatty acids
- anionic lipids


to transmembrane receptors

2. Pharmacology of central nervous system proteins

- general anesthetics
- steroids
- hormones
- alcohol
- neurotoxins

General Motivation

1. Lipid-protein interactions: specific binding of
 - cholesterol
 - polyunsaturated fatty acids
 - anionic lipidsto transmembrane receptors




Hanson, ... Stevens, 2009, Structure

2. Pharmacology of central nervous system proteins
 - general anesthetics
 - steroids
 - hormones
 - alcohol
 - neurotoxins

General Motivation


1. Lipid-protein interactions: specific binding of
 - cholesterol
 - polyunsaturated fatty acids
 - anionic lipidsto transmembrane receptors



Hanson, ... Stevens, 2009, Structure


2. Pharmacology of central nervous system proteins

- general anesthetics
- steroids
- hormones
- alcohol
- neurotoxins




Woll, Murlidaran...Brannigan, Garcia, Eckenhoff, 2016, J Biol Chem



ligand-binding in a lamellar phase






ligand-binding in a lamellar phase






ligand-binding in a lamellar phase



ligand-binding in a lamellar phase






ligand-binding in a lamellar phase






typical approach for coarse-grained systems: pathway dependent method
(ABF/Umbrella Sampling/Metadynamics/Steered MD)

ligand-binding in a lamellar phase



typical approach for coarse-grained systems: pathway dependent method
(ABF/Umbrella Sampling/Metadynamics/Steered MD)




ligand-binding in a lamellar phase




typical approach for coarse-grained systems: pathway dependent method
(ABF/Umbrella Sampling/Metadynamics/Steered MD)

Can we do this using
Alchemical Free Energy
Perturbation (AFEP)?

ligand-binding in a lamellar phase






typical approach for coarse-grained systems: pathway dependent method
(ABF/Umbrella Sampling/Metadynamics/Steered MD)




Can we do this using
Alchemical Free Energy
Perturbation (AFEP)?

ligand-binding in a lamellar phase




typical approach for coarse-grained systems: pathway dependent method
(ABF/Umbrella Sampling/Metadynamics/Steered MD)




Can we do this using
Alchemical Free Energy
Perturbation (AFEP)?

"Classic" Alchemical Free Energy Perturbation


Double Decoupling



Classical Binding Assumptions




Classical Binding Assumptions



ligand is **dilute** : no interactions between ligand


Classical Binding Assumptions



ligand is **dilute** : no interactions between ligand

ligand is **abundant** : most ligand is free ligand


Classical Binding Assumptions



ligand is **dilute** : no interactions between ligand

ligand is **abundant** : most ligand is free ligand

Classical Binding Assumptions




ligand is **dilute** : no interactions between ligand

ligand is **abundant** : most ligand is free ligand

homogenous bulk, **well-defined volume**
concentration

Classical Binding Assumptions



ligand is **dilute** : no interactions between ligand

ligand is **abundant** : most ligand is free ligand

homogenous bulk, **well-defined volume**
concentration

ligand binding has no effect on **receptor-receptor
interactions**

Classical Binding Assumptions

ligand is **dilute** : no interactions between ligand




ligand is **abundant** : most ligand is free ligand

homogenous bulk, **well-defined volume**
concentration

ligand binding has no effect on **receptor-receptor
interactions**

Classical Binding Assumptions




ligand is **dilute** : no interactions between ligand X

ligand is **abundant** : most ligand is free ligand

homogenous bulk, **well-defined volume**
concentration

ligand binding has no effect on **receptor-receptor
interactions**

Classical Binding Assumptions




ligand is **dilute** : no interactions between ligand X

ligand is **abundant** : most ligand is free ligand X

homogenous bulk, **well-defined volume**
concentration

ligand binding has no effect on **receptor-receptor
interactions**

Classical Binding Assumptions



ligand is **dilute** : no interactions between ligand



ligand is **abundant** : most ligand is free ligand



homogenous bulk, **well-defined volume**
concentration



ligand binding has no effect on **receptor-receptor
interactions**

Goal

Goal

An implementation that is :

Goal

An implementation that is :

1. applicable for a full range of ligand concentrations

Goal

An implementation that is :

1. applicable for a full range of ligand concentrations
2. tractable for phase-separated systems

Goal

An implementation that is :

1. applicable for a full range of ligand concentrations
2. tractable for phase-separated systems
3. Avoids REST

Goal

An implementation that is :

1. applicable for a full range of ligand concentrations
2. tractable for phase-separated systems
3. Avoids REST
4. Avoids a complicated restraint scheme

Goal

An implementation that is :

1. applicable for a full range of ligand concentrations
2. tractable for phase-separated systems
3. Avoids REST
4. Avoids a complicated restraint scheme
5. Yields probabilities rather than constants of arbitrary dimension

Obstacles : Interpretation & Implementation

Obstacles : Interpretation & Implementation

Interpretation : predicting ligand titration given non-ideality and ambiguous concentration

Obstacles : Interpretation & Implementation

Interpretation : predicting ligand titration given non-ideality and ambiguous concentration

Implementation : Amplified ACRES obstacles
Accuracy **C**onvergence **R**estraints **E**ndpoint **S**olvation

INTERPRETATION : Invalid Assumptions

$$L + R \rightarrow RL$$

INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}}$$

probability
of site being
occupied


INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$

probability
of site being
occupied

INTERPRETATION : Invalid Assumptions




$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$

probability
of site being
occupied

In infinitely dilute limit:

INTERPRETATION : Invalid Assumptions




$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$

probability
of site being
occupied

In infinitely dilute limit:

$$\frac{[RL]}{[R]} = K_A [L] = K_A p_{\text{free}} [L]_{\text{tot}}$$

INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$


probability
of site being
occupied

In infinitely dilute limit:

$$\frac{[RL]}{[R]} = K_A [L] = K_A p_{\text{free}} [L]_{\text{tot}}$$



INTERPRETATION : Invalid Assumptions




$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$


probability
of site being
occupied

In infinitely dilute limit:

$$\frac{[RL]}{[R]} = K_A [L] = K_A p_{\text{free}} [L]_{\text{tot}}$$



INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$


probability
of site being
occupied

In infinitely dilute limit:

$$\frac{[RL]}{[R]} = K_A [L] = K_A p_{free} [L]_{tot}$$

total ligand

INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$


|
probability
of site being
occupied

In infinitely dilute limit:

$$\frac{[RL]}{[R]} = K_A [L] = K_A p_{free} [L]_{tot}$$

prob of free ligand total ligand

INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$

|
probability
of site being
occupied


In infinitely dilute limit:

$$\frac{[RL]}{[R]} = K_A [L] = K_A p_{free} [L]_{tot}$$

prob of free ligand total ligand

$$p_{occ} = \frac{1}{1 + \frac{1}{K_A [L]}}$$

INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$

probability
of site being
occupied

In infinitely dilute limit:


$$\frac{[RL]}{[R]} = K_A [L] = K_A p_{free} [L]_{tot}$$

prob of free ligand total ligand

$$p_{occ} = \frac{1}{1 + \frac{1}{K_A [L]}}$$

Otherwise

INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$

probability
of site being
occupied

In infinitely dilute limit:

$$\frac{[RL]}{[R]} = K_A [L]$$


p_{free} [L]_{tot}
prob of free ligand

$$p_{occ} = \frac{1}{1 + \frac{1}{K_A [L]}}$$

Otherwise

$$\frac{[RL]}{[R]} \sim \frac{\{RL\}}{\{R\}} = K_A \{L\} = K_A \gamma_L [L]$$

INTERPRETATION : Invalid Assumptions



$$p_{occ} = \frac{1}{1 + \left(\frac{p_{occ}}{p_{unocc}}\right)^{-1}} = \frac{1}{1 + \left(\frac{[RL]}{[R]}\right)^{-1}}$$

probability
of site being
occupied

In infinitely dilute limit:

$$\frac{[RL]}{[R]} = K_A [L] = K_A p_{free} [L]_{tot}$$

prob of free ligand total ligand

$$p_{occ} = \frac{1}{1 + \frac{1}{K_A [L]}}$$

Otherwise

$$\frac{[RL]}{[R]} \sim \frac{\{RL\}}{\{R\}} = K_A \{L\} = K_A \gamma_L [L] = K_A \gamma_L p_{free} [L]_{tot}$$

INTERPRETATION : Invalid Assumptions

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

INTERPRETATION : Invalid Assumptions

$$\frac{[RL]}{[R]} = \underbrace{K_A \gamma_L p_{\text{free}}}_{\downarrow} [L]_{\text{tot}}$$

INTERPRETATION : Invalid Assumptions

Sigmoidal binding assumes constant across concentrations

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

INTERPRETATION : Invalid Assumptions

Sigmoidal binding assumes constant across concentrations

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

↓

only constant for large $[L]_{\text{tot}}$

INTERPRETATION : Invalid Assumptions

Sigmoidal binding assumes constant across concentrations

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

↓

constant for small $[L]_{\text{tot}}$

only constant for large $[L]_{\text{tot}}$

INTERPRETATION : Invalid Assumptions

Sigmoidal binding assumes constant across concentrations

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

↓

constant by definition constant for small $[L]_{\text{tot}}$ only constant for large $[L]_{\text{tot}}$

INTERPRETATION : Invalid Assumptions

Sigmoidal binding assumes constant across concentrations

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

↓

constant by definitionconstant for small $[L]_{\text{tot}}$ only constant for large $[L]_{\text{tot}}$

problem: Only true for intermediate concentrations. Rarely confirmed experimentally.
KA not useful outside infinitely dilute limit.

INTERPRETATION : Invalid Assumptions

Sigmoidal binding assumes constant across concentrations

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

↓

constant by definitionconstant for small $[L]_{\text{tot}}$ only constant for large $[L]_{\text{tot}}$

problem: Only true for intermediate concentrations. Rarely confirmed experimentally.
KA not useful outside infinitely dilute limit.

solution: Combine all this stuff together into one concentration dependent term, drop assumptions, forget about infinitely dilute limit.

INTERPRETATION : Invalid Assumptions

Sigmoidal binding assumes constant across concentrations

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

↓

constant by definition constant for small $[L]_{\text{tot}}$ only constant for large $[L]_{\text{tot}}$

problem: Only true for intermediate concentrations. Rarely confirmed experimentally.
KA not useful outside infinitely dilute limit.

solution: Combine all this stuff together into one concentration dependent term, drop assumptions, forget about infinitely dilute limit.

$$\frac{[RL]}{[R]} = \kappa_L [L]_{\text{tot}}$$

INTERPRETATION : Invalid Assumptions

Sigmoidal binding assumes constant across concentrations

$$\frac{[RL]}{[R]} = K_A \gamma_L p_{\text{free}} [L]_{\text{tot}}$$

↓

constant by definitionconstant for small $[L]_{\text{tot}}$ only constant for large $[L]_{\text{tot}}$

problem: Only true for intermediate concentrations. Rarely confirmed experimentally.
KA not useful outside infinitely dilute limit.

solution: Combine all this stuff together into one concentration dependent term, drop assumptions, forget about infinitely dilute limit.

$$\frac{[RL]}{[R]} = \kappa_L [L]_{\text{tot}}$$


$$p_{\text{occ}} = \frac{1}{1 + \frac{1}{\kappa_L [L]_{\text{tot}}}}$$

INTERPRETATION : Ambiguous Concentration

problem: What is the concentration of  in



INTERPRETATION : Ambiguous Concentration

problem: What is the concentration of  in






INTERPRETATION : Ambiguous Concentration

problem: What is the concentration of  in







INTERPRETATION : Ambiguous Concentration

problem: What is the concentration of  in








INTERPRETATION : Ambiguous Concentration

problem: What is the concentration of  in








INTERPRETATION : Ambiguous Concentration

problem: What is the concentration of  in



INTERPRETATION : Ambiguous Concentration

problem: What is the concentration of  in



What about standard state?

INTERPRETATION : Ambiguous Concentration

Solution part 1: Stay anchored by occupation probabilities

$$\frac{[RL]}{[R]} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \kappa_L [L]_{\text{tot}}$$

INTERPRETATION : Ambiguous Concentration


Solution part 1: Stay anchored by occupation probabilities

$$\frac{[RL]}{[R]} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \kappa_L [L]_{\text{tot}}$$


physically unambiguous

INTERPRETATION : Ambiguous Concentration

Solution part 1: Stay anchored by occupation probabilities


$$\frac{[RL]}{[R]} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \kappa_L [L]_{\text{tot}}$$


physically unambiguous

how we define $[L]_{\text{tot}}$

INTERPRETATION : Ambiguous Concentration

Solution part 1: Stay anchored by occupation probabilities

$$\frac{[RL]}{[R]} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \kappa_L [L]_{\text{tot}}$$


physically unambiguous

how we define $[L]_{\text{tot}}$

determines scale and dimensions of κ_L

INTERPRETATION : Ambiguous Concentration

Solution part 1: Stay anchored by occupation probabilities

$$\frac{[RL]}{[R]} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \kappa_L [L]_{\text{tot}}$$


physically unambiguous

how we define $[L]_{\text{tot}}$

determines scale and dimensions of κ_L

but quantity $\kappa_L [L]_{\text{tot}}$ remains **unambiguous.**

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

total number of ligand molecules



$$\mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
total number of receptor molecules

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
number of ligand molecules **per receptor** (still intensive!)
total number of receptor molecules

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
 \mathcal{L} \equiv $\frac{N_L}{N_R} \frac{1}{\mathcal{V}}$ = $\frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}}$ = $\frac{n_L}{\mathcal{V}}$
|
total number of receptor molecules

number of ligand molecules **per receptor** (still intensive!)
generalized volume that determines concentration units/scale

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\text{generalized concentration } - \mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
number of ligand molecules **per receptor** (still intensive!)
total number of receptor molecules
|
generalized volume that determines concentration units/scale

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\text{generalized concentration } - \mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
total number of receptor molecules

number of ligand molecules **per receptor** (still intensive!)

generalized volume that determines concentration units/scale

$$= \begin{cases} [L]_{\text{tot}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}} \\ [L]_{\text{tot}}^{\text{area}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}^{\text{area}}} \\ n_L & \text{if } \mathcal{V} \equiv 1 \end{cases}$$

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\text{generalized concentration } - \mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
total number of receptor molecules

number of ligand molecules **per receptor** (still intensive!)

generalized volume that determines concentration units/scale

$$= \begin{cases} [L]_{\text{tot}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}} \\ [L]_{\text{tot}}^{\text{area}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}^{\text{area}}} \\ n_L & \text{if } \mathcal{V} \equiv 1 \end{cases}$$

standard generalized “volume”

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\text{generalized concentration } - \mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
total number of receptor molecules

number of ligand molecules **per receptor** (still intensive!)

generalized volume that determines concentration units/scale

$$= \begin{cases} [L]_{\text{tot}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}} \\ [L]_{\text{tot}}^{\text{area}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}^{\text{area}}} \\ n_L & \text{if } \mathcal{V} \equiv 1 \end{cases}$$

standard generalized “volume”

volume **per receptor**

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\text{generalized concentration } - \mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
total number of receptor molecules

number of ligand molecules **per receptor** (still intensive!)

generalized volume that determines concentration units/scale

$$= \begin{cases} [L]_{\text{tot}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}} \\ [L]_{\text{tot}}^{\text{area}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}^{\text{area}}} \\ n_L & \text{if } \mathcal{V} \equiv 1 \end{cases}$$

standard generalized “volume”

volume **per receptor**

area **per receptor**

INTERPRETATION : Ambiguous Concentration

Solution part 2: Represent intensive ligand abundance via “per receptor” units

$$\text{generalized concentration } - \mathcal{L} \equiv \frac{N_L}{N_R} \frac{1}{\mathcal{V}} = \frac{[L]_{\text{tot}}}{[R]_{\text{tot}}} \frac{1}{\mathcal{V}} = \frac{n_L}{\mathcal{V}}$$

total number of ligand molecules
|
total number of receptor molecules

number of ligand molecules **per receptor** (still intensive!)

generalized volume that determines concentration units/scale

$$= \begin{cases} [L]_{\text{tot}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}} \quad \text{volume } \mathbf{per receptor} \\ [L]_{\text{tot}}^{\text{area}} & \text{if } \mathcal{V} \equiv \frac{1}{[R]_{\text{tot}}^{\text{area}}} \quad \text{area } \mathbf{per receptor} \\ n_L & \text{if } \mathcal{V} \equiv 1 \quad \text{receptors } \mathbf{per receptor} \end{cases}$$

|
standard generalized “volume”


ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site




ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site



ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site




Our typical **solution**: flat-well translational restraint



ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site




Our typical **solution**: flat-well translational restraint

$$U_R(r) = \begin{cases} 0 & \text{if } r \leq r_{\max}, \\ \frac{1}{2} r_d (r - r_{\max})^2 & \text{if } r > r_{\max} \end{cases}$$




ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site



Our typical **solution**: flat-well translational restraint




$$U_R(r) = \begin{cases} 0 & \text{if } r \leq r_{\max}, \\ \frac{1}{2} r_d (r - r_{\max})^2 & \text{if } r > r_{\max} \end{cases}$$




ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site



Our typical **solution**: flat-well translational restraint




$$U_R(r) = \begin{cases} 0 & \text{if } r \leq r_{\max}, \\ \frac{1}{2} r_d (r - r_{\max})^2 & \text{if } r > r_{\max} \end{cases}$$




ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site



Our typical **solution**: flat-well translational restraint




$$U_R(r) = \begin{cases} 0 & \text{if } r \leq r_{\max}, \\ \frac{1}{2} r_d (r - r_{\max})^2 & \text{if } r > r_{\max} \end{cases}$$

Correction: Negligible contribution in coupled state.
Volume-based translational contribution in decoupled state.




ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site



Our typical **solution**: flat-well translational restraint



$$U_R(r) = \begin{cases} 0 & \text{if } r \leq r_{\max}, \\ \frac{1}{2} r_d (r - r_{\max})^2 & \text{if } r > r_{\max} \end{cases}$$


Correction: Negligible contribution in coupled state.
Volume-based translational contribution in decoupled state.

problem: Works well for binding sites that can be fully enclosed with a regular shape. Not superficial and/or lipid sites!




ACRES : Convergence Restraints

problem: Annihilated/decoupled ligand samples outside binding site




Our typical **solution**: flat-well translational restraint



$$U_R(r) = \begin{cases} 0 & \text{if } r \leq r_{\max}, \\ \frac{1}{2} r_d (r - r_{\max})^2 & \text{if } r > r_{\max} \end{cases}$$

Correction: Negligible contribution in coupled state.
Volume-based translational contribution in decoupled state.

problem: Works well for binding sites that can be fully enclosed with a regular shape. Not superficial and/or lipid sites!



ACRES : Anisotropic Restraints

“Canonical” anisotropic approach: 7 restraints


Published in final edited form as:

J Chem Theory Comput. 2013 January 8; 9(1): 794–802. doi:10.1021/ct3008099.

**Standard binding free energies from computer simulations:
What is the best strategy?**

James C. Gumbart[†], Benoît Roux^{†,¶*}, and Christophe Chipot^{‡,§,*}

Biosciences Division, Argonne National Laboratory, Argonne, IL 60439, and Beckman Institute,
University of Illinois at Urbana-Champaign, Urbana IL 61801



ACRES : Anisotropic Restraints

“Canonical” anisotropic approach: 7 restraints


Published in final edited form as:

J Chem Theory Comput. 2013 January 8; 9(1): 794–802. doi:10.1021/ct3008099.

**Standard binding free energies from computer simulations:
What is the best strategy?**

James C. Gumbart[†], Benoît Roux^{†,¶*}, and Christophe Chipot^{‡,§,*}

Biosciences Division, Argonne National Laboratory, Argonne, IL 60439, and Beckman Institute,
University of Illinois at Urbana-Champaign, Urbana IL 61801



Our **solution**: 1 anisotropic restraint
RMSD restraint in protein reference frame

ACRES : Anisotropic Restraints

“Canonical” anisotropic approach: 7 restraints


Published in final edited form as:

J Chem Theory Comput. 2013 January 8; 9(1): 794–802. doi:10.1021/ct3008099.

**Standard binding free energies from computer simulations:
What is the best strategy?**

James C. Gumbart[†], Benoît Roux^{†,¶*}, and Christophe Chipot^{‡,§,*}

Biosciences Division, Argonne National Laboratory, Argonne, IL 60439, and Beckman Institute,
University of Illinois at Urbana-Champaign, Urbana IL 61801



Our **solution**: 1 anisotropic restraint
RMSD restraint in protein reference frame

$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$


ACRES : Anisotropic Restraints

“Canonical” anisotropic approach: 7 restraints

Published in final edited form as:
J Chem Theory Comput. 2013 January 8; 9(1): 794–802. doi:10.1021/ct3008099.

**Standard binding free energies from computer simulations:
What is the best strategy?**

James C. Gumbart[†], Benoît Roux^{†,¶*}, and Christophe Chipot^{‡,§,*}
Biosciences Division, Argonne National Laboratory, Argonne, IL 60439, and Beckman Institute,
University of Illinois at Urbana-Champaign, Urbana IL 61801



Our **solution**: 1 anisotropic restraint
RMSD restraint in protein reference frame

$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$

transformation that
minimizes **protein** RMSD

ACRES : Anisotropic Restraints

“Canonical” anisotropic approach: 7 restraints


Published in final edited form as:

J Chem Theory Comput. 2013 January 8; 9(1): 794–802. doi:10.1021/ct3008099.

**Standard binding free energies from computer simulations:
What is the best strategy?**

James C. Gumbart[†], Benoît Roux^{†,¶*}, and Christophe Chipot^{‡,§,*}

Biosciences Division, Argonne National Laboratory, Argonne, IL 60439, and Beckman Institute,
University of Illinois at Urbana-Champaign, Urbana IL 61801



Our **solution**: 1 anisotropic restraint
RMSD restraint in protein reference frame

$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$

$$d = \left[\sum_{l,\text{lig}} (\mathbf{x}'_l - \mathbf{x}_l^{\text{ref}})^2 \right]^{\frac{1}{2}}$$

transformation that
minimizes **protein** RMSD


ACRES : Anisotropic Restraints

“Canonical” anisotropic approach: 7 restraints

Published in final edited form as:
J Chem Theory Comput. 2013 January 8; 9(1): 794–802. doi:10.1021/ct3008099.

**Standard binding free energies from computer simulations:
What is the best strategy?**

James C. Gumbart[†], Benoît Roux^{†,¶*}, and Christophe Chipot^{‡,§,*}
Biosciences Division, Argonne National Laboratory, Argonne, IL 60439, and Beckman Institute,
University of Illinois at Urbana-Champaign, Urbana IL 61801



Our **solution**: 1 anisotropic restraint
RMSD restraint in protein reference frame

$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$

transformation that

minimizes **protein** RMSD

$$d = \left[\sum_{l,\text{lig}} (\mathbf{x}'_l - \mathbf{x}_l^{\text{ref}})^2 \right]^{\frac{1}{2}}$$

“Displacement from Bound Configuration” (DBC):
RMSD in **protein** reference
frame

ACRES : Anisotropic Restraints


“Canonical” anisotropic approach: 7 restraints

Published in final edited form as:
J Chem Theory Comput. 2013 January 8; 9(1): 794–802. doi:10.1021/ct3008099.

**Standard binding free energies from computer simulations:
What is the best strategy?**

James C. Gumbart[†], Benoît Roux^{†,¶*}, and Christophe Chipot^{‡,§,*}

Biosciences Division, Argonne National Laboratory, Argonne, IL 60439, and Beckman Institute,
University of Illinois at Urbana-Champaign, Urbana IL 61801



Our **solution**: 1 anisotropic restraint
RMSD restraint in protein reference frame

$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$

transformation that

minimizes **protein** RMSD

$$d = \left[\sum_{l,\text{lig}} (\mathbf{x}'_l - \mathbf{x}_l^{\text{ref}})^2 \right]^{\frac{1}{2}} \quad U_{DBC}(d) = \begin{cases} 0 & \text{if } d \leq d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d > d_{\max} \end{cases}$$

“Displacement from Bound Configuration” (DBC):
RMSD in **protein** reference
frame

ACRES : Anisotropic Restraints


“Canonical” anisotropic approach: 7 restraints

Published in final edited form as:
J Chem Theory Comput. 2013 January 8; 9(1): 794–802. doi:10.1021/ct3008099.

**Standard binding free energies from computer simulations:
What is the best strategy?**

James C. Gumbart[†], Benoît Roux^{†,¶*}, and Christophe Chipot^{‡,§,*}

Biosciences Division, Argonne National Laboratory, Argonne, IL 60439, and Beckman Institute,
University of Illinois at Urbana-Champaign, Urbana IL 61801



Our **solution**: 1 anisotropic restraint
RMSD restraint in protein reference frame

$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$

transformation that

minimizes **protein** RMSD

$$d = \left[\sum_{l,\text{lig}} (\mathbf{x}'_l - \mathbf{x}_l^{\text{ref}})^2 \right]^{\frac{1}{2}}$$


“Displacement from Bound Configuration” (DBC):
RMSD in **protein** reference frame

$$U_{DBC}(d) = \begin{cases} 0 & \text{if } d \leq d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d > d_{\max} \end{cases}$$

Flat-well potential on
DBC Collective Variable

ACRES: DBC Restraints


calculated using blue
protein atoms



$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$

sum over orange ligand atoms

$$d = \left[\sum_{l,\text{lig}} (\mathbf{x}'_l - \mathbf{x}_l^{\text{ref}})^2 \right]^{\frac{1}{2}}$$




$$U_{DBC}(d) = \begin{cases} 0 & \text{if } d \leq d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d > d_{\max} \end{cases}$$

restraint potential
 $d_{\max} \sim 2\text{\AA}$
 $k_d \sim 100 \text{ kcal/mol}/\text{\AA}^2$

Salari, Joseph, Lohia, Henin, Brannigan,
JCTC 2018 submitted
<https://arxiv.org/abs/1801.04901>

ACRES: DBC Restraints

calculated using blue
protein atoms



$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$

sum over orange ligand atoms

A ribbon diagram of a protein structure. Blue sticks represent protein atoms, and orange sticks represent ligand atoms. An orange line points from the text "sum over orange ligand atoms" to one of the orange sticks. A red stick represents a reference ligand atom.

$$d = \left[\sum_{l,\text{lig}} (\mathbf{x}'_l - \mathbf{x}_l^{\text{ref}})^2 \right]^{\frac{1}{2}}$$

$$U_{DBC}(d) = \begin{cases} 0 & \text{if } d \leq d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d > d_{\max} \end{cases}$$


restraint potential
 $d_{\max} \sim 2\text{\AA}$
 $k_d \sim 100 \text{ kcal/mol}/\text{\AA}^2$

correction restraint free energy perturbation (TI)

Salari, Joseph, Lohia, Henin, Brannigan,
JCTC 2018 submitted
<https://arxiv.org/abs/1801.04901>

ACRES: DBC Restraints


calculated using blue
protein atoms



$$\mathbf{x}'_i = \rho(\mathbf{x}_i - \bar{\mathbf{x}}_R) + \bar{\mathbf{x}}_R^{\text{ref}},$$

sum over orange ligand atoms

$$d = \left[\sum_{l,\text{lig}} (\mathbf{x}'_l - \mathbf{x}_l^{\text{ref}})^2 \right]^{\frac{1}{2}}$$



$$U_{DBC}(d) = \begin{cases} 0 & \text{if } d \leq d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d > d_{\max} \end{cases}$$


restraint potential
 $d_{\max} \sim 2\text{\AA}$
 $k_d \sim 100 \text{ kcal/mol}/\text{\AA}^2$

*Implemented in NAMD2.12
Collective Variables Module*

correction restraint free energy perturbation (TI)


Salari, Joseph, Lohia, Henin, Brannigan,
JCTC 2018 submitted
<https://arxiv.org/abs/1801.04901>

ACRES: Exclusion Restraints




- problem** high affinity bulk ligand replaces decoupled ligand
- solution** exclusion restraint potential (opposite of DBC restraint potential) on non-test ligand molecules
- correction** *none needed*: by definition, unoccupied state has no ligand bound

$$U_{DBC}(d) = \begin{cases} 0 & \text{if } d \leq d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d > d_{\max} \end{cases}$$




$$U_{EX}(d) = \begin{cases} 0 & \text{if } d > d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d \leq d_{\max} \end{cases}$$

ACRES: Exclusion Restraints





- problem** high affinity bulk ligand replaces decoupled ligand
- solution** exclusion restraint potential (opposite of DBC restraint potential) on non-test ligand molecules
- correction** *none needed*: by definition, unoccupied state has no ligand bound

$$U_{DBC}(d) = \begin{cases} 0 & \text{if } d \leq d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d > d_{\max} \end{cases}$$




$$U_{EX}(d) = \begin{cases} 0 & \text{if } d > d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d \leq d_{\max} \end{cases}$$

ACRES: Exclusion Restraints



- problem** high affinity bulk ligand replaces decoupled ligand
- solution** exclusion restraint potential (opposite of DBC restraint potential) on non-test ligand molecules
- correction** *none needed*: by definition, unoccupied state has no ligand bound

$$U_{DBC}(d) = \begin{cases} 0 & \text{if } d \leq d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d > d_{\max} \end{cases}$$



$$U_{EX}(d) = \begin{cases} 0 & \text{if } d > d_{\max}, \\ \frac{1}{2} k_d (d - d_{\max})^2 & \text{if } d \leq d_{\max} \end{cases}$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}}$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_b^- Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} Z_b^-}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{Z_r^{\Delta}} \right)$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_b^- Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{Z_r^{\Delta}} \right)$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{Z_b^=} \right) \left(\frac{Z_g^\circ}{Z_b^- Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^\circ} \right) \left(\frac{Z_g^\Delta Z_b^-}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^\Delta}{Z_g^\Delta Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{Z_r^\Delta} \right)$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^- Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{\cancel{Z_g^\Delta} \cancel{Z_b^-}}{\cancel{Z_g^{\circ\Delta}}} \right) \left(\frac{Z_r^\Delta}{Z_g^\Delta Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{Z_r^\Delta} \right)$$

Cycle for Lipophilic Ligand


$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^{=}}{Z_b^{\square}} \right) \left(\frac{Z_b^{-} Z_g^{=}}{Z_b^{=}} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^{-} Z_g^{=}}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{\cancel{Z_g^{\Delta}} \cancel{Z_b^{-}}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^{-}} \right) \left(\frac{\cancel{Z_r^{\bullet}}}{\cancel{Z_r^{\Delta}}} \right)$$

Cycle for Lipophilic Ligand

$$\begin{aligned}\kappa_{\mathcal{L}} \mathcal{L} &= \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^{=}}{Z_b^{\square}} \right) \left(\frac{Z_b^{-} Z_g^{=}}{Z_b^{=}} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^{-} Z_g^{=}}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^{-}}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^{-}} \right) \left(\frac{\cancel{Z_r^{\bullet}}}{Z_g^{\Delta}} \right) \\ &= \left(\frac{Z_b^{=}}{Z_b^{\square}} \right) \left(\frac{Z_b^{-} Z_g^{=}}{Z_b^{=}} \right) \left(\frac{Z_g^{\circ}}{Z_g^{=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^{-}} \right)\end{aligned}$$


Cycle for Lipophilic Ligand

$$\begin{aligned}\kappa_{\mathcal{L}} \mathcal{L} &= \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^- Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\bullet}}{\cancel{Z_r^{\Delta}}} \right) \\ &= \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)\end{aligned}$$



Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^- Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{\cancel{Z_r^{\Delta}}} \right)$$
$$= \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




$n_L \alpha$

$\sim 10^2 x_{\text{CHOL}}$


Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^- Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{\cancel{Z_r^{\Delta}}} \right)$$
$$= \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^- Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{\cancel{Z_r^{\Delta}}} \right)$$
$$= \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$



Cycle for Lipophilic Ligand


$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^- Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{\cancel{Z_g^{\circ\Delta}}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{\cancel{Z_r^{\Delta}}} \right)$$
$$= \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$



Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_b^- Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} Z_b^-}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\bullet}}{Z_g^{\Delta}} \right)$$

$$= \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




$$\frac{2\pi r_R^3}{3z_R A^=} \frac{1}{1 - \cos \theta_R} \sim 10^{-1}$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^-} \cancel{Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{\cancel{Z_g^{\circ\Delta}}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\bullet}}{\cancel{Z_g^{\Delta}}} \right)$$

$$= \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




$$\frac{2\pi r_R^3}{3z_R A^=} \frac{1}{1 - \cos \theta_R} \sim 10^{-1}$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^-} \cancel{Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{\cancel{Z_g^{\circ\Delta}}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\bullet}}{\cancel{Z_g^{\Delta}}} \right)$$

$$= \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




$$\frac{2\pi r_R^3}{3z_R A^=} \frac{1}{1 - \cos \theta_R} \sim 10^{-1}$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^- Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{\cancel{Z_g^{\Delta} Z_b^-}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{\cancel{Z_r^{\Delta}}} \right)$$

$$= \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




$$\frac{2\pi r_R^3}{3z_R A^=} \frac{1}{1 - \cos \theta_R} \sim 10^{-1}$$

Cycle for Lipophilic Ligand

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\blacksquare}}{Z_b^{\square}} = \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{\cancel{Z_b^- Z_g^=}} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_g^{\Delta} \cancel{Z_b^-}}{Z_g^{\circ\Delta}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right) \left(\frac{Z_r^{\blacksquare}}{Z_r^{\Delta}} \right)$$

$$= \left(\frac{Z_b^=}{}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=}{}{Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




$$\frac{2\pi r_R^3}{3z_R A^=} \frac{1}{1 - \cos \theta_R} \sim 10^{-1}$$

Application 1 : GPCR-Bound Cholesterol

bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$




Application 1 : GPCR-Bound Cholesterol

bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$

Quadratic
Mixture
Model




Application 1 : GPCR-Bound Cholesterol

bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$

Quadratic
Mixture
Model

$$\mu_g^0 + RT \ln x_g = \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2$$



Application 1 : GPCR-Bound Cholesterol


bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$

Quadratic
Mixture
Model

$$\mu_g^0 + RT \ln x_g = \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2$$

enthalpy of mixing



Application 1 : GPCR-Bound Cholesterol


bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$

Quadratic
Mixture
Model

$$\begin{aligned}\mu_g^0 + RT \ln x_g &= \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2 \\ P_x &= \frac{x_b}{x_g} = P_0 e^{-h^0(1-x_b)^2/RT}\end{aligned}$$

enthalpy of mixing



Application 1 : GPCR-Bound Cholesterol


bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$

Quadratic
Mixture
Model

$$\begin{aligned}\mu_g^0 + RT \ln x_g &= \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2 \\ P_x &= \frac{x_b}{x_g} = P_0 e^{-h^0(1-x_b)^2/RT}\end{aligned}$$

enthalpy of mixing



AFEP decouple single cholesterol molecule from cholesterol:POPC bilayer

Application 1 : GPCR-Bound Cholesterol


bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$


Quadratic
Mixture
Model

$$\begin{aligned} \mu_g^0 + RT \ln x_g &= \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2 \\ P_x &= \frac{x_b}{x_g} = P_0 e^{-h^0(1-x_b)^2/RT} \end{aligned}$$

enthalpy of mixing



AFEP decouple single cholesterol molecule from cholesterol:POPC bilayer



Application 1 : GPCR-Bound Cholesterol


bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$


Quadratic
Mixture
Model

$$\begin{aligned} \mu_g^0 + RT \ln x_g &= \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2 \\ P_x &= \frac{x_b}{x_g} = P_0 e^{-h^0(1-x_b)^2/RT} \end{aligned}$$

enthalpy of mixing



AFEP decouple single cholesterol molecule from cholesterol:POPC bilayer



Application 1 : GPCR-Bound Cholesterol


bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$


Quadratic
Mixture
Model

$$\begin{aligned} \mu_g^0 + RT \ln x_g &= \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2 \\ P_x &= \frac{x_b}{x_g} = P_0 e^{-h^0(1-x_b)^2/RT} \end{aligned}$$

enthalpy of mixing



AFEP decouple single cholesterol molecule from cholesterol:POPC bilayer



$$P^0 = 3 \times 10^{14}$$

$$h^0 = 1.6 \text{ kcal/mol}$$

Application 1 : GPCR-Bound Cholesterol


bulk/gas
partition
coefficient

$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$


Quadratic
Mixture
Model

$$\begin{aligned}\mu_g^0 + RT \ln x_g &= \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2 \\ P_x &= \frac{x_b}{x_g} = P_0 e^{-h^0(1-x_b)^2/RT}\end{aligned}$$

enthalpy of mixing



AFEP decouple single cholesterol molecule from cholesterol:POPC bilayer



$$P_0 = 3 \times 10^{14}$$

$$h^0 = 1.6 \text{ kcal/mol}$$

$$h^0 \sim \langle u_{\text{chol-popc}} \rangle - \langle u_{\text{chol-chol}} + u_{\text{popc-popc}} \rangle / 2 > 0$$

non-ideality


Application 1 : GPCR-Bound Cholesterol

bulk/gas
partition
coefficient


$$\frac{1}{P_x} \equiv \frac{Z_b^- Z_g^=}{Z_b^=} = \text{AFEP1}$$

Quadratic
Mixture
Model

$$\begin{aligned}\mu_g^0 + RT \ln x_g &= \mu_b^0 + RT \ln x_b + h^0(1 - x_b)^2 \\ P_x &= \frac{x_b}{x_g} = P_0 e^{-h^0(1-x_b)^2/RT}\end{aligned}$$



AFEP decouple single cholesterol molecule from cholesterol:POPC bilayer



$$P^0 = 3 \times 10^{14}$$


$$h^0 = 1.6 \text{ kcal/mol}$$

$$h^0 \sim \langle u_{\text{chol-popc}} \rangle - \langle u_{\text{chol-chol}} + u_{\text{popc-popc}} \rangle / 2 > 0$$

non-ideality

Application 1 : GPCR-Bound Cholesterol



One restraint correction calculation (TI/SOS) gradually decreasing restraint strength




$$\frac{Z_g^{\circ\Delta}}{Z_g^\circ} \sim 10^{-3}$$

Less than 1 hour of calculation for single restraint correction


Application 1 : GPCR-Bound Cholesterol



$\beta 2$ -Adrenergic
3D4S




5-HT_{2B}
4NC3




μ -Opioid
5C1M

Application 1 : GPCR-Bound Cholesterol




$10^{16} \times$




$\beta 2$ -Adrenergic
3D4S

10^9



5-HT_{2B}
4NC3

10^2




μ -Opioid
5C1M

1


Application 1 : GPCR-Bound Cholesterol

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




Application 1 : GPCR-Bound Cholesterol

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




Application 1 : GPCR-Bound Cholesterol

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




Application 1 : GPCR-Bound Cholesterol

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




Application 1 : GPCR-Bound Cholesterol

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




Application 1 : GPCR-Bound Cholesterol

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^=} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$



Application 1 : GPCR-Bound Cholesterol

$$\kappa_{\mathcal{L}} \mathcal{L} = \frac{p_{\text{occ}}}{p_{\text{unocc}}} = \frac{Z_r^{\bullet}}{Z_b^{\square}} = \left(\frac{Z_b^=}{Z_b^{\square}} \right) \left(\frac{Z_b^- Z_g^=} {Z_b^=} \right) \left(\frac{Z_g^{\circ}}{Z_g^=} \right) \left(\frac{Z_g^{\circ\Delta}}{Z_g^{\circ}} \right) \left(\frac{Z_r^{\Delta}}{Z_g^{\Delta} Z_b^-} \right)$$




Ratio	Method	$\beta 2$ -adrenergic	5-HT2B	μ -opioid
κ_x	$\frac{Z_r^{\bullet}}{Z_b^- Z_g^{\circ}} \times \frac{Z_b^- Z_g^{\circ}}{Z_b^{\square}} \frac{1}{x_{\text{CHOL}}}$	$e^{2.7(1-x_{\text{CHOL}})^2} \times$		
		10^9	10^3	10
x_{50}	x_{CHOL} for which $\kappa_x x_{\text{CHOL}} = 1$	10^{-9}	10^{-3}	–



Application 1: Predicted ligand titration

$$p_{occ} = \frac{1}{1 + \frac{1}{\kappa_x x_{CHOL}}}$$



Ratio	Method	β_2 -adrenergic	5-HT2B	μ -opioid
κ_x	$\frac{Z_r^*}{Z_b^- Z_g^o} \times \frac{Z_b^- Z_g^o}{Z_b^o} \frac{1}{x_{CHOL}}$	$e^{2.7(1-x_{CHOL})^2}$	10^9	10^3
x_{50}	x_{CHOL} for which $\kappa_x x_{CHOL} = 1$	10^{-9}	10^{-3}	-




Application 1: Predicted ligand titration



β_2 -Adrenergic
3D4S
 $x_{50} = 10^{-9}$



μ -Opioid
5C1M
no x_{50}



5-HT2B
4NC3
 $x_{50} = 10^{-3}$

Natural x_{50} is in non-ideal range, probability of binding reduced because bulk has become so favorable

Salari, Joseph, Lohia, Henin, Brannigan
JCTC 2018 submitted

<https://arxiv.org/abs/1801.04901>

ACRES : Endpoint Solvation


Simplest AFEP process: assumes decoupled/annihilated state is equivalent to apo state.

Obviously not true


ACRES : Endpoint Solvation

Simplest AFEP process: assumes decoupled/annihilated state is equivalent to apo state.


Obviously not true



bound



annihilated




apo


ACRES : Endpoint Solvation

Simplest AFEP process: assumes decoupled/annihilated state is equivalent to apo state.


Obviously not true



bound



annihilated




apo


A robust calculation requires considering this - different ways to do so.

ACRES : Endpoint Solvation


Typical problem with gradual AFEP:



bound




intermediate




end-point

Overestimates
affinity!




actual apo


Common remedy - reverse for recoupling, combine with overlap sampling



end-point




intermediate




bound

ACRES : Endpoint Solvation


Typical problem with gradual AFEP:



bound




intermediate




end-point

Overestimates
affinity!




actual apo


Common remedy - reverse for recoupling, combine with overlap sampling



end-point



intermediate




bound


They match! (Both are **wrong**...
but they'll match!)

ACRES : Endpoint Solvation


Typical problem with gradual AFEP:



bound




intermediate




end-point

Overestimates
affinity!




actual apo


Common remedy - reverse for recoupling, combine with overlap sampling



end-point



intermediate



bound


They match! (Both are **wrong**...
but they'll match!)

REST can help




ACRES : Endpoint Solvation

Problem is amplified in membranes due to heterogeneous solvation:

if proper membrane apo site looks like



many possible
incorrect endpoints!




REST disrupts phase equilibrium




ACRES : Endpoint Solvation

Problem is amplified in membranes due to heterogeneous solvation:

if proper membrane apo site looks like



many possible
incorrect endpoints!




REST disrupts phase equilibrium

need to track endpoint solvation


ACRES : Endpoint Solvation

Problem is amplified in membranes due to heterogeneous solvation:

if proper membrane apo site looks like




many possible
incorrect endpoints!



REST disrupts phase equilibrium


need to track endpoint solvation





ACRES : Endpoint Solvation

Problem is amplified in membranes due to heterogeneous solvation:

if proper membrane apo site looks like




many possible
incorrect endpoints!



REST disrupts phase equilibrium


need to track endpoint solvation





ACRES : Endpoint Solvation

Problem is amplified in membranes due to heterogeneous solvation:

if proper membrane apo site looks like




many possible
incorrect endpoints!






REST disrupts phase equilibrium

need to track endpoint solvation




Application 2 : pseudosymmetric sites for propofol on GABA(A) receptors






Woll, Murlidaran...Brannigan, Garcia,
Eckenhoff, 2016, J Biol Chem


Application 2 : pseudosymmetric sites for propofol on GABA(A) receptors






Woll, Murlidaran...Brannigan, Garcia,
Eckenhoff, 2016, J Biol Chem




Application 2 : pseudosymmetric sites for propofol on GABA(A) receptors






Woll, Murlidaran...Brannigan, Garcia,
Eckenhoff, 2016, J Biol Chem




Application 2 : pseudosymmetric sites for propofol on GABA(A) receptors








Woll, Murlidaran...Brannigan, Garcia,
Eckenhoff, 2016, J Biol Chem




Application 2 : pseudosymmetric sites for propofol on GABA(A) receptors



Woll, Murlidaran...Brannigan, Garcia,
Eckenhoff, 2016, J Biol Chem



Application 2 : pseudosymmetric sites for sevoflurane on GABA(A) receptors



Murlidaran and Brannigan, in preparation

Summary

Summary

1. With a little adjustment of formalism, AFEP is well-suited to studying non-ideality.

Summary

1. With a little adjustment of formalism, AFEP is well-suited to studying non-ideality.
2. Non-ideality can be crucial for low-affinity ligands - even for relative binding affinities.

Summary

1. With a little adjustment of formalism, AFEP is well-suited to studying non-ideality.
2. Non-ideality can be crucial for low-affinity ligands - even for relative binding affinities.
3. DBC restraints simplify process of calculating absolute binding affinities; could improve automation.

Summary

1. With a little adjustment of formalism, AFEP is well-suited to studying non-ideality.
2. Non-ideality can be crucial for low-affinity ligands - even for relative binding affinities.
3. DBC restraints simplify process of calculating absolute binding affinities; could improve automation.
4. Cholesterol has a positive enthalpy of mixing in POPC, up to about ~50% cholesterol.

Summary

1. With a little adjustment of formalism, AFEP is well-suited to studying non-ideality.
2. Non-ideality can be crucial for low-affinity ligands - even for relative binding affinities.
3. DBC restraints simplify process of calculating absolute binding affinities; could improve automation.
4. Cholesterol has a positive enthalpy of mixing in POPC, up to about ~50% cholesterol.
5. Cholesterol affinities : b₂-adrenergic > serotonin > mu-opioid

Summary

1. With a little adjustment of formalism, AFEP is well-suited to studying non-ideality.
2. Non-ideality can be crucial for low-affinity ligands - even for relative binding affinities.
3. DBC restraints simplify process of calculating absolute binding affinities; could improve automation.
4. Cholesterol has a positive enthalpy of mixing in POPC, up to about ~50% cholesterol.
5. Cholesterol affinities : b₂-adrenergic > serotonin > mu-opioid
6. Heterogeneous lipid “solvation” increases likelihood of selectivity due to differential solvation.

Acknowledgments

Group Members

pLGICs -

Dr. Reza Salari

Dr. Sruthi Murlidaran

Liam Sharp

Kristen Woods

GPCRs

Dr. Reza Salari

Dr. Thomas Joseph

Neurotrophic factors

Ruchi Lohia

AFEP

Dr. Reza Salari

Dr. Thomas Joseph

Dr. Sruthi Murlidaran

Ruchi Lohia

VDAC

Dr. Reza Salari

Shashank Chavali

Collaborators

Dr. Jérôme Hénin, (CNRS-IPBC France)

Dr. Roderic Eckenhoff (University of Pennsylvania)

Dr. John Baenziger (University of Ottawa)

Dr. Joseph Martin (Rutgers)

Dr. Sebastian Hiller (University of Basel)

Dr. Keith Miller (Harvard)

Funding



XSEDE

Extreme Science and Engineering
Discovery Environment

