



# Things left to prove

## Characters and Blocks of Finite Groups

### Gabriel's conference 2025

Benjamin Sambale

31th January 2025

Gabriel likes (to propose) problems

# Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522

# Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281

# Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281
- 2004: *The McKay conjecture and Galois automorphisms*, Ann. of Math. 160, 1129–1140 (71 citations)

# Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281
- 2004: *The McKay conjecture and Galois automorphisms*, Ann. of Math. 160, 1129–1140 (71 citations)
- 2010: *Problems in character theory*, in: Character theory of finite groups, 97–125

# Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281
- 2004: *The McKay conjecture and Galois automorphisms*, Ann. of Math. 160, 1129–1140 (71 citations)
- 2010: *Problems in character theory*, in: Character theory of finite groups, 97–125
- 2023: *Problems on characters: solvable groups*, Publ. Mat. 67, 173–198

# Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281
- 2004: *The McKay conjecture and Galois automorphisms*, Ann. of Math. 160, 1129–1140 (71 citations)
- 2010: *Problems in character theory*, in: Character theory of finite groups, 97–125
- 2023: *Problems on characters: solvable groups*, Publ. Mat. 67, 173–198
- 2024: “All I would like to prove has been proved!”, e-mail correspondence



# Automorphisms

Let  $G$  be a finite group.

# Automorphisms

Let  $G$  be a finite group.

## Definition (Automorphism tower)

Let  $\text{Aut}^1(G) := \text{Aut}(G)$ ,  $\text{Aut}^2(G) := \text{Aut}(\text{Aut}(G)), \dots$

# Automorphisms

Let  $G$  be a finite group.

## Definition (Automorphism tower)

Let  $\text{Aut}^1(G) := \text{Aut}(G)$ ,  $\text{Aut}^2(G) := \text{Aut}(\text{Aut}(G)), \dots$

If  $Z(G) = 1$ , then  $G \cong \text{Inn}(G) \leq \text{Aut}^1(G) \leq \text{Aut}^2(G) \leq \dots$

# Automorphisms

Let  $G$  be a finite group.

## Definition (Automorphism tower)

Let  $\text{Aut}^1(G) := \text{Aut}(G)$ ,  $\text{Aut}^2(G) := \text{Aut}(\text{Aut}(G)), \dots$

If  $Z(G) = 1$ , then  $G \cong \text{Inn}(G) \leq \text{Aut}^1(G) \leq \text{Aut}^2(G) \leq \dots$

## Theorem (WIELANDT)

If  $Z(G) = 1$ , then  $\text{Aut}^n(G) \cong \text{Aut}^{n+1}(G)$  for some  $n$ .

# Automorphisms

## Theorem (HAMKINS)

*The transfinite automorphism tower of any group is bounded by some cardinal number.*

# Automorphisms

## Theorem (HAMKINS)

*The transfinite automorphism tower of any group is bounded by some cardinal number.*

## Problem (Kourovka notebook 11.123)

- Is there a constant  $c$  such that  $|\text{Aut}^n(G)| < c$  for all  $n$ ?
- Is  $\text{Aut}^n(G) \cong \text{Aut}^{n+1}(G)$  for some  $n$ ?
- Can  $\text{Aut}^n(G) \cong \text{Aut}^m(G) \not\cong \text{Aut}^{n+1}(G)$  with  $n < m$  happen?

# Automorphisms

## Example

- If  $G$  is non-abelian simple, then  $\text{Aut}^2(G) \cong \text{Aut}(G)$  (BURNSIDE).

# Automorphisms

## Example

- If  $G$  is non-abelian simple, then  $\text{Aut}^2(G) \cong \text{Aut}(G)$  (BURNSIDE).
- $\text{Aut}(C_2^n) \cong \text{GL}(n, 2)$  is simple (for  $n \geq 3$ ), so  $\text{Aut}^3(C_2^n) \cong \text{Aut}^2(C_2^n)$ .



# Automorphisms

## Example

- If  $G$  is non-abelian simple, then  $\text{Aut}^2(G) \cong \text{Aut}(G)$  (BURNSIDE).
- $\text{Aut}(C_2^n) \cong \text{GL}(n, 2)$  is simple (for  $n \geq 3$ ), so  $\text{Aut}^3(C_2^n) \cong \text{Aut}^2(C_2^n)$ .
- Center can stay non-trivial:  $\text{Aut}(D_8) \cong D_8$ .

# Automorphisms

## Example

- If  $G$  is non-abelian simple, then  $\text{Aut}^2(G) \cong \text{Aut}(G)$  (BURNSIDE).
- $\text{Aut}(C_2^n) \cong \text{GL}(n, 2)$  is simple (for  $n \geq 3$ ), so  $\text{Aut}^3(C_2^n) \cong \text{Aut}^2(C_2^n)$ .
- Center can stay non-trivial:  $\text{Aut}(D_8) \cong D_8$ .
- The sequence can decrease arbitrarily long:  $\text{Aut}(C_{2 \cdot 3^n}) \cong C_{2 \cdot 3^{n-1}}$

# Automorphisms

## Example

- If  $G$  is non-abelian simple, then  $\text{Aut}^2(G) \cong \text{Aut}(G)$  (BURNSIDE).
- $\text{Aut}(C_2^n) \cong \text{GL}(n, 2)$  is simple (for  $n \geq 3$ ), so  $\text{Aut}^3(C_2^n) \cong \text{Aut}^2(C_2^n)$ .
- Center can stay non-trivial:  $\text{Aut}(D_8) \cong D_8$ .
- The sequence can decrease arbitrarily long:  $\text{Aut}(C_{2 \cdot 3^n}) \cong C_{2 \cdot 3^{n-1}}$
- For  $G = \text{SmallGroup}(32, 13)$  we have

$n$	1	2	3	4	5
$ \text{Aut}^n(G) $	$2^7$	$2^{13}$	$2^{28}$	$2^{83}$	?

# Homomorphisms

## Theorem (FROBENIUS)

*For every  $n$ , the number of  $x \in G$  such that  $x^n = 1$  is divisible by  $\gcd(n, |G|)$ .*

# Homomorphisms

## Theorem (FROBENIUS)

*For every  $n$ , the number of  $x \in G$  such that  $x^n = 1$  is divisible by  $\gcd(n, |G|)$ .*

Equivalently,

$$|\mathrm{Hom}(C_n, G)| \equiv 0 \pmod{\gcd(n, |G|)}.$$

# Homomorphisms

## Theorem (FROBENIUS)

*For every  $n$ , the number of  $x \in G$  such that  $x^n = 1$  is divisible by  $\gcd(n, |G|)$ .*

Equivalently,

$$|\mathrm{Hom}(C_n, G)| \equiv 0 \pmod{\gcd(n, |G|)}.$$

## Theorem (YOSHIDA)

*For every finite abelian group  $A$ ,*

$$|\mathrm{Hom}(A, G)| \equiv 0 \pmod{\gcd(|A|, |G|)}.$$

# Homomorphisms

## Conjecture (ASAI–YOSHIDA)

For every finite group  $H$ ,

$$|\mathrm{Hom}(H, G)| \equiv 0 \pmod{\gcd(|H/H'|, |G|)}.$$

# Conjugacy classes

- Let  $k(G)$  be the number of conjugacy classes of  $G$ .



# Conjugacy classes

- Let  $k(G)$  be the number of conjugacy classes of  $G$ .
- LANDAU:  $k(G) > f(|G|)$  for some increasing function  $f$ .

# Conjugacy classes

- Let  $k(G)$  be the number of conjugacy classes of  $G$ .
- LANDAU:  $k(G) > f(|G|)$  for some increasing function  $f$ .

## Theorem (BAUMEISTER–MARÓTI–TONG–VIET)

*For every  $\epsilon > 0$  there exists  $\delta > 0$  such that*

$$k(G) > \frac{\delta \log |G|}{(\log \log |G|)^{3+\epsilon}}.$$

# Conjugacy classes

- Let  $k(G)$  be the number of conjugacy classes of  $G$ .
- LANDAU:  $k(G) > f(|G|)$  for some increasing function  $f$ .

## Theorem (BAUMEISTER–MARÓTI–TONG–VIET)

*For every  $\epsilon > 0$  there exists  $\delta > 0$  such that*

$$k(G) > \frac{\delta \log |G|}{(\log \log |G|)^{3+\epsilon}}.$$

## Conjecture (Folklore?)

Is there a constant  $c$  such that  $k(G) > c \log |G|$  for all  $G$ ?

# Constituents

Let  $\text{Irr}(G)$  be the set of irreducible complex characters of  $G$ .

# Constituents

Let  $\text{Irr}(G)$  be the set of irreducible complex characters of  $G$ .

## Conjecture (HÉTHELYI–KÜLSHAMMER)

Let  $P$  be a  $p$ -group and  $\chi \in \text{Irr}(P)$ . Then the number of irreducible constituents of  $\chi\bar{\chi}$  is  $1 \bmod p - 1$ .

# Constituents

Let  $\text{Irr}(G)$  be the set of irreducible complex characters of  $G$ .

## Conjecture (HÉTHELYI–KÜLSHAMMER)

Let  $P$  be a  $p$ -group and  $\chi \in \text{Irr}(P)$ . Then the number of irreducible constituents of  $\chi\bar{\chi}$  is  $1 \bmod p - 1$ .

Conjecture holds for  $|P| \leq p^6$ .

# Constituents

Let  $\text{Irr}(G)$  be the set of irreducible complex characters of  $G$ .

## Conjecture (HÉTHELYI–KÜLSHAMMER)

Let  $P$  be a  $p$ -group and  $\chi \in \text{Irr}(P)$ . Then the number of irreducible constituents of  $\chi\bar{\chi}$  is  $1 \bmod p-1$ .

Conjecture holds for  $|P| \leq p^6$ .

## Problem (KNUTSON–MURRAY)

Let  $P$  be a  $p$ -group and  $\chi \in \text{Irr}(P)$ . Is there a generalized character  $\psi$  such that  $\chi\psi$  is the regular character of  $P$ ?

# Constituents

Call  $\theta \in \text{Irr}(Z(G))$  **fully ramified** if  $\theta^G$  is a multiple of an irreducible character.



# Constituents

Call  $\theta \in \text{Irr}(Z(G))$  **fully ramified** if  $\theta^G$  is a multiple of an irreducible character.

## Theorem (HOWLETT–ISAACS)

*If  $\theta \in \text{Irr}(Z(G))$  is fully ramified, then  $G$  is solvable.*

# Constituents

Call  $\theta \in \text{Irr}(Z(G))$  **fully ramified** if  $\theta^G$  is a multiple of an irreducible character.

## Theorem (HOWLETT–ISAACS)

*If  $\theta \in \text{Irr}(Z(G))$  is fully ramified, then  $G$  is solvable.*

## Conjecture (HUMPHREYS, NAVARRO)

Let  $\theta \in \text{Irr}(Z(G))$  such that all constituents of  $\theta^G$  have the same degree. Then  $G$  is solvable.

# Constituents

Call  $\theta \in \text{Irr}(Z(G))$  **fully ramified** if  $\theta^G$  is a multiple of an irreducible character.

## Theorem (HOWLETT–ISAACS)

*If  $\theta \in \text{Irr}(Z(G))$  is fully ramified, then  $G$  is solvable.*

## Conjecture (HUMPHREYS, NAVARRO)

Let  $\theta \in \text{Irr}(Z(G))$  such that all constituents of  $\theta^G$  have the same degree. Then  $G$  is solvable.

Conjecture holds whenever  $\theta^G$  has at most two constituents (HIGGS).

# Field of values

For  $\chi \in \text{Irr}(G)$  define the abelian number fields

$$\begin{aligned}\mathbb{Q}(\chi) &:= \mathbb{Q}(\chi(g) : g \in G) \subseteq \mathbb{Q}_{|G|} \subseteq \mathbb{C}, \\ \mathbb{Q}(G) &:= \mathbb{Q}(\chi(g) : \chi \in \text{Irr}(G), g \in G) \subseteq \mathbb{Q}_{|G|}.\end{aligned}$$

# Field of values

For  $\chi \in \text{Irr}(G)$  define the abelian number fields

$$\begin{aligned}\mathbb{Q}(\chi) &:= \mathbb{Q}(\chi(g) : g \in G) \subseteq \mathbb{Q}_{|G|} \subseteq \mathbb{C}, \\ \mathbb{Q}(G) &:= \mathbb{Q}(\chi(g) : \chi \in \text{Irr}(G), g \in G) \subseteq \mathbb{Q}_{|G|}.\end{aligned}$$

By the KRONECKER–WEBER theorem, every abelian number field lies in some  $\mathbb{Q}_n$ .

# Field of values

For  $\chi \in \text{Irr}(G)$  define the abelian number fields

$$\begin{aligned}\mathbb{Q}(\chi) &:= \mathbb{Q}(\chi(g) : g \in G) \subseteq \mathbb{Q}_{|G|} \subseteq \mathbb{C}, \\ \mathbb{Q}(G) &:= \mathbb{Q}(\chi(g) : \chi \in \text{Irr}(G), g \in G) \subseteq \mathbb{Q}_{|G|}.\end{aligned}$$

By the KRONECKER–WEBER theorem, every abelian number field lies in some  $\mathbb{Q}_n$ .

## Theorem (FEIN–GORDON)

*For every abelian number field  $F$  there exist a group  $G$  and  $\chi \in \text{Irr}(G)$  such that  $\mathbb{Q}(\chi) = F$ .*

# Field of values

## Conjecture

- Not every abelian number field has the form  $\mathbb{Q}(G)$ .

# Field of values

## Conjecture

- Not every abelian number field has the form  $\mathbb{Q}(G)$ .
- For every  $d$ , there are only finitely many number fields  $\mathbb{Q}(G)$  of degree  $d$ .



# Field of values

## Conjecture

- Not every abelian number field has the form  $\mathbb{Q}(G)$ .
- For every  $d$ , there are only finitely many number fields  $\mathbb{Q}(G)$  of degree  $d$ .

There are only finitely many fields  $\mathbb{Q}(G)$  of degree  $d$  where  $G$  is solvable or simple (FARIAS E SOARES, FEIT–SEITZ).

# Field of values

## Conjecture

- Not every abelian number field has the form  $\mathbb{Q}(G)$ .
- For every  $d$ , there are only finitely many number fields  $\mathbb{Q}(G)$  of degree  $d$ .

There are only finitely many fields  $\mathbb{Q}(G)$  of degree  $d$  where  $G$  is solvable or simple (FARIAS E SOARES, FEIT–SEITZ).

## Example (quadratic fields)

Let  $d \mid 210$  or  $d \in \{-231, -11, 13, 17\}$ . Then there exists  $G$  with  $\mathbb{Q}(G) \cong \mathbb{Q}(\sqrt{d})$ .

# Field of values

## Theorem (ROBINSON–THOMPSON)

*For  $n > 24$  we have*

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

*In particular,  $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$  is an elementary abelian 2-group.*

# Field of values

## Theorem (ROBINSON–THOMPSON)

*For  $n > 24$  we have*

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

*In particular,  $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$  is an elementary abelian 2-group.*

## Problem

# Field of values

## Theorem (ROBINSON–THOMPSON)

*For  $n > 24$  we have*

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

*In particular,  $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$  is an elementary abelian 2-group.*

## Problem

- Is every abelian group the Galois group of some  $\mathbb{Q}(G)$ ?

# Field of values

## Theorem (ROBINSON–THOMPSON)

*For  $n > 24$  we have*

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt[(-1)^{\frac{p-1}{2}}]{p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

*In particular,  $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$  is an elementary abelian 2-group.*

## Problem

- Is every abelian group the Galois group of some  $\mathbb{Q}(G)$ ?
- Is there a group  $G$  such that  $|\mathbb{Q}(G) : \mathbb{Q}| = 7$ ?

# Field of values

## Theorem (ROBINSON–THOMPSON)

For  $n > 24$  we have

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

In particular,  $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$  is an elementary abelian 2-group.

## Problem

- Is every abelian group the Galois group of some  $\mathbb{Q}(G)$ ?
- Is there a group  $G$  such that  $|\mathbb{Q}(G) : \mathbb{Q}| = 7$ ?
- Is there a group  $G$  such that  $\mathbb{Q}(G) = \mathbb{Q}(\sqrt{11})$ ?

# Field of values

## Theorem (ROBINSON–THOMPSON)

*For  $n > 24$  we have*

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

*In particular,  $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$  is an elementary abelian 2-group.*

## Problem

- Is every abelian group the Galois group of some  $\mathbb{Q}(G)$ ?
- Is there a group  $G$  such that  $|\mathbb{Q}(G) : \mathbb{Q}| = 7$ ?
- Is there a group  $G$  such that  $\mathbb{Q}(G) = \mathbb{Q}(\sqrt{11})$ ?
- Is there a solvable group  $G$  such that  $\mathbb{Q}(G) = \mathbb{Q}(\sqrt{-5})$ ?



# Field of values

For  $\chi \in \text{Irr}(G)$  let  $f(\chi)$  be the smallest integer  $n$  such that  $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_n$ .

# Field of values

For  $\chi \in \text{Irr}(G)$  let  $f(\chi)$  be the smallest integer  $n$  such that  $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_n$ .

## Conjecture (HUNG–TIEP)

We have  $|\mathbb{Q}_{f(\chi)} : \mathbb{Q}(\chi)| \leq \chi(1)$  for all  $\chi \in \text{Irr}(G)$ .

# Field of values

For  $\chi \in \text{Irr}(G)$  let  $f(\chi)$  be the smallest integer  $n$  such that  $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_n$ .

## Conjecture (HUNG–TIEP)

We have  $|\mathbb{Q}_{f(\chi)} : \mathbb{Q}(\chi)| \leq \chi(1)$  for all  $\chi \in \text{Irr}(G)$ .

Conjecture holds whenever  $\chi(1)$  is a prime (HUNG–TIEP–ZALESSKI).

# Field of values

For  $\chi \in \text{Irr}(G)$  let  $f(\chi)$  be the smallest integer  $n$  such that  $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_n$ .

## Conjecture (HUNG–TIEP)

We have  $|\mathbb{Q}_{f(\chi)} : \mathbb{Q}(\chi)| \leq \chi(1)$  for all  $\chi \in \text{Irr}(G)$ .

Conjecture holds whenever  $\chi(1)$  is a prime (HUNG–TIEP–ZALESSKI).

## Problem (HUNG–TIEP)

Let  $z = \zeta_1 + \dots + \zeta_n \in \mathbb{C}$  be a sum of roots of unity. Let  $c(z)$  be the smallest  $m$  such that  $z \in \mathbb{Q}_m$ . Is  $|\mathbb{Q}_{c(z)} : \mathbb{Q}(z)| \leq n$ ?

# Algebraic integers

- Let  $\mathbb{Z}_G$  be the ring of integers of  $\mathbb{Q}(G)$ .

# Algebraic integers

- Let  $\mathbb{Z}_G$  be the ring of integers of  $\mathbb{Q}(G)$ .
- Let  $\mathbb{Z}[G] := \mathbb{Z}[\chi(g) : \chi \in \text{Irr}(G), g \in G] \subseteq \mathbb{Z}_G$ .

# Algebraic integers

- Let  $\mathbb{Z}_G$  be the ring of integers of  $\mathbb{Q}(G)$ .
- Let  $\mathbb{Z}[G] := \mathbb{Z}[\chi(g) : \chi \in \text{Irr}(G), g \in G] \subseteq \mathbb{Z}_G$ .
- Then  $\mathbb{Z}_G/\mathbb{Z}[G]$  is a finite abelian group.

# Algebraic integers

- Let  $\mathbb{Z}_G$  be the ring of integers of  $\mathbb{Q}(G)$ .
- Let  $\mathbb{Z}[G] := \mathbb{Z}[\chi(g) : \chi \in \text{Irr}(G), g \in G] \subseteq \mathbb{Z}_G$ .
- Then  $\mathbb{Z}_G/\mathbb{Z}[G]$  is a finite abelian group.

## Conjecture (BÄCHLE–S.)

The exponent of  $\mathbb{Z}_G/\mathbb{Z}[G]$  divides  $|G|$ .



# Algebraic integers

- Let  $\mathbb{Z}_G$  be the ring of integers of  $\mathbb{Q}(G)$ .
- Let  $\mathbb{Z}[G] := \mathbb{Z}[\chi(g) : \chi \in \text{Irr}(G), g \in G] \subseteq \mathbb{Z}_G$ .
- Then  $\mathbb{Z}_G/\mathbb{Z}[G]$  is a finite abelian group.

## Conjecture (BÄCHLE–S.)

The exponent of  $\mathbb{Z}_G/\mathbb{Z}[G]$  divides  $|G|$ .

## Example

Conjecture holds for nilpotent groups. For  $G = C_{15} \rtimes D_{16}$ ,

$$\mathbb{Z}_G/\mathbb{Z}[G] \cong C_{120}^2 \times C_{60}^2 \times C_{12}^4 \times C_4^4 \times C_2^{14}.$$

# Weights

- Let  $\pi$  be a set of primes.

# Weights

- Let  $\pi$  be a set of primes.
- Let  $l(G)$  be the number of conjugacy classes of  $\pi'$ -elements of  $G$ .

# Weights

- Let  $\pi$  be a set of primes.
- Let  $l(G)$  be the number of conjugacy classes of  $\pi'$ -elements of  $G$ .
- Let  $\chi \in \text{Irr}(G)$  of  $\pi$ -**defect** 0 if  $|G|/\chi(1)$  is a  $\pi'$ -number.

# Weights

- Let  $\pi$  be a set of primes.
- Let  $l(G)$  be the number of conjugacy classes of  $\pi'$ -elements of  $G$ .
- Let  $\chi \in \text{Irr}(G)$  of  $\pi$ -**defect** 0 if  $|G|/\chi(1)$  is a  $\pi'$ -number.
- Let  $k^0(G)$  be the number of  $\pi$ -defect 0 characters of  $G$ .

# Weights

- Let  $\pi$  be a set of primes.
- Let  $l(G)$  be the number of conjugacy classes of  $\pi'$ -elements of  $G$ .
- Let  $\chi \in \text{Irr}(G)$  of  $\pi$ -**defect** 0 if  $|G|/\chi(1)$  is a  $\pi'$ -number.
- Let  $k^0(G)$  be the number of  $\pi$ -defect 0 characters of  $G$ .

## Theorem (NAVARRO–S.)

*If  $G$  is  $\pi$ -separable with solvable Hall  $\pi$ -subgroups, then*

$$l(G) = \sum_P k^0(N_G(P)/P)$$

*where  $P$  runs through the nilpotent  $\pi$ -subgroups of  $G$  up to conjugation.*

# Weights

- Let  $\pi$  be a set of primes.
- Let  $l(G)$  be the number of conjugacy classes of  $\pi'$ -elements of  $G$ .
- Let  $\chi \in \text{Irr}(G)$  of  $\pi$ -defect 0 if  $|G|/\chi(1)$  is a  $\pi'$ -number.
- Let  $k^0(G)$  be the number of  $\pi$ -defect 0 characters of  $G$ .

## Theorem (NAVARRO–S.)

*If  $G$  is  $\pi$ -separable with solvable Hall  $\pi$ -subgroups, then*

$$l(G) = \sum_P k^0(N_G(P)/P)$$

*where  $P$  runs through the nilpotent  $\pi$ -subgroups of  $G$  up to conjugation.*

For  $\pi = \{p\}$ , this is Alperin's weight conjecture for  $p$ -solvable groups.

# Weights

## Definition

Define the **weight** of a group  $P$  by

$$\mu(P) = 1 - \sum_{\substack{Q < P \\ N_P(Q) = Q}} \mu(Q) \quad (Q \text{ up to conjugation}).$$



# Weights

## Definition

Define the **weight** of a group  $P$  by

$$\mu(P) = 1 - \sum_{\substack{Q < P \\ N_P(Q) = Q}} \mu(Q) \quad (Q \text{ up to conjugation}).$$

## Conjecture (NAVARRO–S.)

For every  $\pi$ -separable group  $G$ ,

$$l(G) = \sum_P \mu(P) k^0(N_G(P)/P)$$

where  $P$  runs through the  $\pi$ -subgroups of  $G$  up to conjugation.

# Weights

## Example

- If  $P$  is nilpotent, then  $\mu(P) = 1$ .

# Weights

## Example

- If  $P$  is nilpotent, then  $\mu(P) = 1$ .
- If  $P$  is solvable, but not nilpotent, then  $\mu(P) = 0$  (CARTER subgroups).

# Weights

## Example

- If  $P$  is nilpotent, then  $\mu(P) = 1$ .
- If  $P$  is solvable, but not nilpotent, then  $\mu(P) = 0$  (CARTER subgroups).
- On the other hand,

$P$	$A_5$	$A_6$	$A_7$	$A_8$	$\text{PSL}(3, 4)$	$A_5^2$	$S_n (3 \leq n \leq 11)$
$\mu(P)$	1	-2	1	-1	6	-1	0

# Characterization of blocks

Let  $G^0$  be the set of  $p$ -regular elements of  $G$ .

# Characterization of blocks

Let  $G^0$  be the set of  $p$ -regular elements of  $G$ .

## Theorem (OSIMA)

*A subset  $J \subseteq \text{Irr}(G)$  is a union of  $p$ -blocks if and only if*

$$\sum_{\chi \in J} \chi(g)\chi(h) = 0 \quad (\forall g \in G^0, h \in G \setminus G^0).$$

# Characterization of blocks

Let  $G^0$  be the set of  $p$ -regular elements of  $G$ .

## Theorem (OSIMA)

*A subset  $J \subseteq \text{Irr}(G)$  is a union of  $p$ -blocks if and only if*

$$\sum_{\chi \in J} \chi(g)\chi(h) = 0 \quad (\forall g \in G^0, h \in G \setminus G^0).$$

## Conjecture (HARADA)

It is enough to fix  $g = 1$  in Osima's theorem.

# Basic sets

- Let  $B$  be a  $p$ -block of  $G$ .



# Basic sets

- Let  $B$  be a  $p$ -block of  $G$ .
- For  $\chi \in \text{Irr}(G)$  let  $\chi^0$  be the restriction to  $G^0$ .

# Basic sets

- Let  $B$  be a  $p$ -block of  $G$ .
- For  $\chi \in \text{Irr}(G)$  let  $\chi^0$  be the restriction to  $G^0$ .
- The Brauer characters in  $p$ -solvable groups have the form  $\chi^0$  (FONG–SWAN).

# Basic sets

- Let  $B$  be a  $p$ -block of  $G$ .
- For  $\chi \in \text{Irr}(G)$  let  $\chi^0$  be the restriction to  $G^0$ .
- The Brauer characters in  $p$ -solvable groups have the form  $\chi^0$  (FONG–SWAN).

## Conjecture (ordinary basic set, GECK)

There exist  $\chi_1, \dots, \chi_l \in \text{Irr}(B)$  such that  $\chi_1^0, \dots, \chi_l^0$  is a  $\mathbb{Z}$ -basis for the ring of generalized Brauer characters of  $B$ .

# Basic sets

- Let  $B$  be a  $p$ -block of  $G$ .
- For  $\chi \in \text{Irr}(G)$  let  $\chi^0$  be the restriction to  $G^0$ .
- The Brauer characters in  $p$ -solvable groups have the form  $\chi^0$  (FONG–SWAN).

## Conjecture (ordinary basic set, GECK)

There exist  $\chi_1, \dots, \chi_l \in \text{Irr}(B)$  such that  $\chi_1^0, \dots, \chi_l^0$  is a  $\mathbb{Z}$ -basis for the ring of generalized Brauer characters of  $B$ .

This holds in a strong sense for “most” groups of Lie type (BRUNAT–DUDAS–TAYLOR).

# Number of characters

- For  $\chi, \psi \in \text{Irr}(G)$  let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

# Number of characters

- For  $\chi, \psi \in \text{Irr}(G)$  let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

- Let  $d$  be the defect of  $B$ . Then  $p^d[\chi, \chi]^0 \in \mathbb{N}$  for all  $\chi \in \text{Irr}(B)$ .

# Number of characters

- For  $\chi, \psi \in \text{Irr}(G)$  let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

- Let  $d$  be the defect of  $B$ . Then  $p^d[\chi, \chi]^0 \in \mathbb{N}$  for all  $\chi \in \text{Irr}(B)$ .

## Conjecture (NAVARRO-S.)

We have  $p^d[\chi, \chi]^0 \geq l(B)$  for all  $\chi \in \text{Irr}(B)$ .

# Number of characters

- For  $\chi, \psi \in \text{Irr}(G)$  let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

- Let  $d$  be the defect of  $B$ . Then  $p^d[\chi, \chi]^0 \in \mathbb{N}$  for all  $\chi \in \text{Irr}(B)$ .

## Conjecture (NAVARRO-S.)

We have  $p^d[\chi, \chi]^0 \geq l(B)$  for all  $\chi \in \text{Irr}(B)$ .

- This implies Brauer's conjecture  $k(B) \leq p^d$  with equality only if  $B$  has abelian defect groups.



# Number of characters

- For  $\chi, \psi \in \text{Irr}(G)$  let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

- Let  $d$  be the defect of  $B$ . Then  $p^d[\chi, \chi]^0 \in \mathbb{N}$  for all  $\chi \in \text{Irr}(B)$ .

## Conjecture (NAVARRO-S.)

We have  $p^d[\chi, \chi]^0 \geq l(B)$  for all  $\chi \in \text{Irr}(B)$ .

- This implies Brauer's conjecture  $k(B) \leq p^d$  with equality only if  $B$  has abelian defect groups.
- The case  $\chi = 1_G$  was conjectured by MURAI and implies FROBENIUS conjecture:  $G$  is  $p$ -nilpotent iff  $|G^0| = |G|_{p'}$  (known via CFSG).

# Fusion numbers

- Let

$$X_B := (\chi(g_i) : \chi \in \text{Irr}(B), i = 1, \dots, k(G))$$

be a submatrix of the character table.

# Fusion numbers

- Let

$$X_B := (\chi(g_i) : \chi \in \text{Irr}(B), i = 1, \dots, k(G))$$

be a submatrix of the character table.

- The non-zero elementary divisors of  $X_B^t \overline{X_B}$  over a complete discrete valuation ring are  $p$ -powers  $e_1, \dots, e_t$ .

# Fusion numbers

- Let

$$X_B := (\chi(g_i) : \chi \in \text{Irr}(B), i = 1, \dots, k(G))$$

be a submatrix of the character table.

- The non-zero elementary divisors of  $X_B^t \overline{X_B}$  over a complete discrete valuation ring are  $p$ -powers  $e_1, \dots, e_t$ .

## Conjecture (S.)

We have  $\gamma(B) := \frac{1}{e_1} + \dots + \frac{1}{e_t} \geq 1$  with equality if and only if  $B$  is nilpotent.

# Fusion numbers

- Let

$$X_B := (\chi(g_i) : \chi \in \text{Irr}(B), i = 1, \dots, k(G))$$

be a submatrix of the character table.

- The non-zero elementary divisors of  $X_B^t \overline{X_B}$  over a complete discrete valuation ring are  $p$ -powers  $e_1, \dots, e_t$ .

## Conjecture (S.)

We have  $\gamma(B) := \frac{1}{e_1} + \dots + \frac{1}{e_t} \geq 1$  with equality if and only if  $B$  is nilpotent.

## Example

Conjecture holds for symmetric groups and “ATLAS groups”. For the principal 2-block  $B$  of the Monster,  $\gamma(B) \approx 39.5$ .

## Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*

## Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands.* [finishing his second book]

## Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*



## Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*
- *Silly? I am in the middle of a restaurant.*

## Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*
- *Silly? I am in the middle of a restaurant.*
- *Seriously, I have thought on this for 3 minutes.*

## Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands.* [finishing his second book]
- *If these two guys are not Morita equivalent then I won't say the word Morita again!* [they weren't]
- *Silly? I am in the middle of a restaurant.*
- *Seriously, I have thought on this for 3 minutes.*
- *Zero.* [answer to: How many examples did you check?]

## Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*
- *Silly? I am in the middle of a restaurant.*
- *Seriously, I have thought on this for 3 minutes.*
- *Zero. [answer to: How many examples did you check?]*
- *I am also amazed by my intuition, sorry to say!*

## Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*
- *Silly? I am in the middle of a restaurant.*
- *Seriously, I have thought on this for 3 minutes.*
- *Zero. [answer to: How many examples did you check?]*
- *I am also amazed by my intuition, sorry to say!*
- *25th anniversary of my wedding. I have promised my wife not to think on mathematics for a couple of days. Now you are making my promise easy to break. . .*