



Things left to prove

Characters and Blocks of Finite Groups

Gabriel's conference 2025

Benjamin Sambale

31th January 2025

Gabriel likes (to propose) problems

Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522

Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281

Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281
- 2004: *The McKay conjecture and Galois automorphisms*, Ann. of Math. 160, 1129–1140 (71 citations)

Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281
- 2004: *The McKay conjecture and Galois automorphisms*, Ann. of Math. 160, 1129–1140 (71 citations)
- 2010: *Problems in character theory*, in: Character theory of finite groups, 97–125

Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281
- 2004: *The McKay conjecture and Galois automorphisms*, Ann. of Math. 160, 1129–1140 (71 citations)
- 2010: *Problems in character theory*, in: Character theory of finite groups, 97–125
- 2023: *Problems on characters: solvable groups*, Publ. Mat. 67, 173–198

Gabriel likes (to propose) problems

- 1994: *Some Open Problems on Coprime Action and Character Correspondences*, Bull. London Math. Soc. 26, 513–522
- 2004: *Problems on characters and Sylow subgroups*, in: Finite groups, 275–281
- 2004: *The McKay conjecture and Galois automorphisms*, Ann. of Math. 160, 1129–1140 (71 citations)
- 2010: *Problems in character theory*, in: Character theory of finite groups, 97–125
- 2023: *Problems on characters: solvable groups*, Publ. Mat. 67, 173–198
- 2024: “All I would like to prove has been proved!”, e-mail correspondence

Automorphisms

Let G be a finite group.

Automorphisms

Let G be a finite group.

Definition (Automorphism tower)

Let $\text{Aut}^0(G) := G$ and $\text{Aut}^{n+1}(G) := \text{Aut}(\text{Aut}^n(G))$ for $n \geq 0$.

Automorphisms

Let G be a finite group.

Definition (Automorphism tower)

Let $\text{Aut}^0(G) := G$ and $\text{Aut}^{n+1}(G) := \text{Aut}(\text{Aut}^n(G))$ for $n \geq 0$.

If $Z(G) = 1$, then $G \cong \text{Inn}(G) \leq \text{Aut}^1(G) \leq \text{Aut}^2(G) \leq \dots$

Automorphisms

Let G be a finite group.

Definition (Automorphism tower)

Let $\text{Aut}^0(G) := G$ and $\text{Aut}^{n+1}(G) := \text{Aut}(\text{Aut}^n(G))$ for $n \geq 0$.

If $Z(G) = 1$, then $G \cong \text{Inn}(G) \leq \text{Aut}^1(G) \leq \text{Aut}^2(G) \leq \dots$

Theorem (WIELANDT)

If $Z(G) = 1$, then $\text{Aut}^n(G) \cong \text{Aut}^{n+1}(G)$ for some n .

Automorphisms

Theorem (HAMKINS)

The transfinite automorphism tower of any group is bounded by some cardinal number.

Automorphisms

Theorem (HAMKINS)

The transfinite automorphism tower of any group is bounded by some cardinal number.

Problem (Kourovka notebook 11.123)

- Is there a constant c such that $|\text{Aut}^n(G)| < c$ for all n ?
- Is $\text{Aut}^n(G) \cong \text{Aut}^{n+1}(G)$ for some n ?
- Can $\text{Aut}^n(G) \cong \text{Aut}^m(G) \not\cong \text{Aut}^{n+1}(G)$ happen for some $n < m$?

Automorphisms

Example

- If G is non-abelian simple, then $\text{Aut}^2(G) \cong \text{Aut}(G)$ (BURNSIDE).

Automorphisms

Example

- If G is non-abelian simple, then $\text{Aut}^2(G) \cong \text{Aut}(G)$ (BURNSIDE).
- $\text{Aut}(C_2^n) \cong \text{GL}(n, 2)$ is simple (for $n \geq 3$), so $\text{Aut}^3(C_2^n) \cong \text{Aut}^2(C_2^n)$.

Automorphisms

Example

- If G is non-abelian simple, then $\text{Aut}^2(G) \cong \text{Aut}(G)$ (BURNSIDE).
- $\text{Aut}(C_2^n) \cong \text{GL}(n, 2)$ is simple (for $n \geq 3$), so $\text{Aut}^3(C_2^n) \cong \text{Aut}^2(C_2^n)$.
- Center can stay non-trivial: $\text{Aut}(D_8) \cong D_8$.

Automorphisms

Example

- If G is non-abelian simple, then $\text{Aut}^2(G) \cong \text{Aut}(G)$ (BURNSIDE).
- $\text{Aut}(C_2^n) \cong \text{GL}(n, 2)$ is simple (for $n \geq 3$), so $\text{Aut}^3(C_2^n) \cong \text{Aut}^2(C_2^n)$.
- Center can stay non-trivial: $\text{Aut}(D_8) \cong D_8$.
- The sequence can decrease arbitrarily long: $\text{Aut}(C_{2 \cdot 3^n}) \cong C_{2 \cdot 3^{n-1}}$

Automorphisms

Example

- If G is non-abelian simple, then $\text{Aut}^2(G) \cong \text{Aut}(G)$ (BURNSIDE).
- $\text{Aut}(C_2^n) \cong \text{GL}(n, 2)$ is simple (for $n \geq 3$), so $\text{Aut}^3(C_2^n) \cong \text{Aut}^2(C_2^n)$.
- Center can stay non-trivial: $\text{Aut}(D_8) \cong D_8$.
- The sequence can decrease arbitrarily long: $\text{Aut}(C_{2 \cdot 3^n}) \cong C_{2 \cdot 3^{n-1}}$
- For $G = \text{SmallGroup}(32, 13)$ we have

n	0	1	2	3	4	5
$ \text{Aut}^n(G) $	2^5	2^7	2^{13}	2^{28}	2^{83}	?

Homomorphisms

Theorem (FROBENIUS)

For every n , the number of $x \in G$ such that $x^n = 1$ is divisible by $\gcd(n, |G|)$.

Homomorphisms

Theorem (FROBENIUS)

For every n , the number of $x \in G$ such that $x^n = 1$ is divisible by $\gcd(n, |G|)$.

Equivalently,

$$|\mathrm{Hom}(C_n, G)| \equiv 0 \pmod{\gcd(n, |G|)}.$$

Homomorphisms

Theorem (FROBENIUS)

For every n , the number of $x \in G$ such that $x^n = 1$ is divisible by $\gcd(n, |G|)$.

Equivalently,

$$|\mathrm{Hom}(C_n, G)| \equiv 0 \pmod{\gcd(n, |G|)}.$$

Theorem (YOSHIDA)

For every finite abelian group A ,

$$|\mathrm{Hom}(A, G)| \equiv 0 \pmod{\gcd(|A|, |G|)}.$$

Homomorphisms

Conjecture (ASAI–YOSHIDA)

For every finite group H ,

$$|\mathrm{Hom}(H, G)| \equiv 0 \pmod{\gcd(|H/H'|, |G|)}.$$

Conjugacy classes

- Let $k(G)$ be the number of conjugacy classes of G .

Conjugacy classes

- Let $k(G)$ be the number of conjugacy classes of G .
- LANDAU: $k(G) > f(|G|)$ for some increasing function f .

Conjugacy classes

- Let $k(G)$ be the number of conjugacy classes of G .
- LANDAU: $k(G) > f(|G|)$ for some increasing function f .

Theorem (BAUMEISTER–MARÓTI–TONG–VIET)

For every $\epsilon > 0$ there exists $\delta > 0$ such that

$$k(G) > \frac{\delta \log |G|}{(\log \log |G|)^{3+\epsilon}}.$$

Conjugacy classes

- Let $k(G)$ be the number of conjugacy classes of G .
- LANDAU: $k(G) > f(|G|)$ for some increasing function f .

Theorem (BAUMEISTER–MARÓTI–TONG–VIET)

For every $\epsilon > 0$ there exists $\delta > 0$ such that

$$k(G) > \frac{\delta \log |G|}{(\log \log |G|)^{3+\epsilon}}.$$

Conjecture (Folklore?)

Is there a constant c such that $k(G) > c \log |G|$ for all G ?

Constituents

Let $\text{Irr}(G)$ be the set of irreducible complex characters of G .

Constituents

Let $\text{Irr}(G)$ be the set of irreducible complex characters of G .

Conjecture (HÉTHELYI–KÜLSHAMMER)

Let P be a p -group and $\chi \in \text{Irr}(P)$. Then the number of irreducible constituents of $\chi\bar{\chi}$ is $1 \bmod p - 1$.

Constituents

Let $\text{Irr}(G)$ be the set of irreducible complex characters of G .

Conjecture (HÉTHELYI–KÜLSHAMMER)

Let P be a p -group and $\chi \in \text{Irr}(P)$. Then the number of irreducible constituents of $\chi\bar{\chi}$ is $1 \bmod p-1$.

Conjecture holds for $|P| \leq p^6$.

Constituents

Let $\text{Irr}(G)$ be the set of irreducible complex characters of G .

Conjecture (HÉTHELYI–KÜLSHAMMER)

Let P be a p -group and $\chi \in \text{Irr}(P)$. Then the number of irreducible constituents of $\chi\bar{\chi}$ is $1 \bmod p-1$.

Conjecture holds for $|P| \leq p^6$.

Problem (KNUTSON–MURRAY)

Let P be a p -group and $\chi \in \text{Irr}(P)$. Is there a generalized character ψ such that $\chi\psi$ is the regular character of P ?

Constituents

Call $\theta \in \text{Irr}(Z(G))$ **fully ramified** if θ^G is a multiple of an irreducible character.

Constituents

Call $\theta \in \text{Irr}(Z(G))$ **fully ramified** if θ^G is a multiple of an irreducible character.

Theorem (HOWLETT–ISAACS)

If $\theta \in \text{Irr}(Z(G))$ is fully ramified, then G is solvable.

Constituents

Call $\theta \in \text{Irr}(Z(G))$ **fully ramified** if θ^G is a multiple of an irreducible character.

Theorem (HOWLETT–ISAACS)

If $\theta \in \text{Irr}(Z(G))$ is fully ramified, then G is solvable.

Conjecture (HUMPHREYS, NAVARRO)

Let $\theta \in \text{Irr}(Z(G))$ such that all constituents of θ^G have the same degree. Then G is solvable.

Constituents

Call $\theta \in \text{Irr}(Z(G))$ **fully ramified** if θ^G is a multiple of an irreducible character.

Theorem (HOWLETT–ISAACS)

If $\theta \in \text{Irr}(Z(G))$ is fully ramified, then G is solvable.

Conjecture (HUMPHREYS, NAVARRO)

Let $\theta \in \text{Irr}(Z(G))$ such that all constituents of θ^G have the same degree. Then G is solvable.

Conjecture holds whenever θ^G has at most two constituents (HIGGS).

Field of values

For $\chi \in \text{Irr}(G)$ define the abelian number fields

$$\mathbb{Q}(\chi) := \mathbb{Q}(\chi(g) : g \in G) \subseteq \mathbb{Q}_{|G|} \subseteq \mathbb{C},$$

$$\mathbb{Q}(G) := \mathbb{Q}(\chi(g) : \chi \in \text{Irr}(G), g \in G) \subseteq \mathbb{Q}_{|G|}.$$

Field of values

For $\chi \in \text{Irr}(G)$ define the abelian number fields

$$\begin{aligned}\mathbb{Q}(\chi) &:= \mathbb{Q}(\chi(g) : g \in G) \subseteq \mathbb{Q}_{|G|} \subseteq \mathbb{C}, \\ \mathbb{Q}(G) &:= \mathbb{Q}(\chi(g) : \chi \in \text{Irr}(G), g \in G) \subseteq \mathbb{Q}_{|G|}.\end{aligned}$$

By the KRONECKER–WEBER theorem, every abelian number field lies in some \mathbb{Q}_n .

Field of values

For $\chi \in \text{Irr}(G)$ define the abelian number fields

$$\begin{aligned}\mathbb{Q}(\chi) &:= \mathbb{Q}(\chi(g) : g \in G) \subseteq \mathbb{Q}_{|G|} \subseteq \mathbb{C}, \\ \mathbb{Q}(G) &:= \mathbb{Q}(\chi(g) : \chi \in \text{Irr}(G), g \in G) \subseteq \mathbb{Q}_{|G|}.\end{aligned}$$

By the KRONECKER–WEBER theorem, every abelian number field lies in some \mathbb{Q}_n .

Theorem (FEIN–GORDON)

For every abelian number field F there exist a group G and $\chi \in \text{Irr}(G)$ such that $\mathbb{Q}(\chi) = F$.

Field of values

Conjecture

- Not every abelian number field has the form $\mathbb{Q}(G)$.

Field of values

Conjecture

- Not every abelian number field has the form $\mathbb{Q}(G)$.
- For every d , there are only finitely many number fields $\mathbb{Q}(G)$ of degree d .

Field of values

Conjecture

- Not every abelian number field has the form $\mathbb{Q}(G)$.
- For every d , there are only finitely many number fields $\mathbb{Q}(G)$ of degree d .

There are only finitely many fields $\mathbb{Q}(G)$ of degree d where G is solvable or simple (FARIAS E SOARES, FEIT–SEITZ).

Field of values

Conjecture

- Not every abelian number field has the form $\mathbb{Q}(G)$.
- For every d , there are only finitely many number fields $\mathbb{Q}(G)$ of degree d .

There are only finitely many fields $\mathbb{Q}(G)$ of degree d where G is solvable or simple (FARIAS E SOARES, FEIT–SEITZ).

Example (quadratic fields)

Let $d \mid 210$ or $d \in \{-231, -11, 13, 17\}$. Then there exists G with $\mathbb{Q}(G) \cong \mathbb{Q}(\sqrt{d})$.

Field of values

Theorem (ROBINSON–THOMPSON)

For $n > 24$ we have

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

In particular, $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$ is an elementary abelian 2-group.

Field of values

Theorem (ROBINSON–THOMPSON)

For $n > 24$ we have

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

In particular, $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$ is an elementary abelian 2-group.

Problem

Field of values

Theorem (ROBINSON–THOMPSON)

For $n > 24$ we have

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

In particular, $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$ is an elementary abelian 2-group.

Problem

- Is every abelian group the Galois group of some $\mathbb{Q}(G)$?

Field of values

Theorem (ROBINSON–THOMPSON)

For $n > 24$ we have

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt[(-1)^{\frac{p-1}{2}}]{p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

In particular, $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$ is an elementary abelian 2-group.

Problem

- Is every abelian group the Galois group of some $\mathbb{Q}(G)$?
- Is there a group G such that $|\mathbb{Q}(G) : \mathbb{Q}| = 7$?

Field of values

Theorem (ROBINSON–THOMPSON)

For $n > 24$ we have

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

In particular, $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$ is an elementary abelian 2-group.

Problem

- Is every abelian group the Galois group of some $\mathbb{Q}(G)$?
- Is there a group G such that $|\mathbb{Q}(G) : \mathbb{Q}| = 7$?
- Is there a group G such that $\mathbb{Q}(G) = \mathbb{Q}(\sqrt{11})$?

Field of values

Theorem (ROBINSON–THOMPSON)

For $n > 24$ we have

$$\mathbb{Q}(A_n) = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p} : p \text{ odd prime } \leq n, p \neq n-2\right).$$

In particular, $\text{Gal}(\mathbb{Q}(A_n)|\mathbb{Q})$ is an elementary abelian 2-group.

Problem

- Is every abelian group the Galois group of some $\mathbb{Q}(G)$?
- Is there a group G such that $|\mathbb{Q}(G) : \mathbb{Q}| = 7$?
- Is there a group G such that $\mathbb{Q}(G) = \mathbb{Q}(\sqrt{11})$?
- Is there a solvable group G such that $\mathbb{Q}(G) = \mathbb{Q}(\sqrt{-5})$?

Field of values

For $\chi \in \text{Irr}(G)$ let $f(\chi)$ be the smallest integer n such that $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_n$.

Field of values

For $\chi \in \text{Irr}(G)$ let $f(\chi)$ be the smallest integer n such that $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_n$.

Conjecture (HUNG–TIEP)

We have $|\mathbb{Q}_{f(\chi)} : \mathbb{Q}(\chi)| \leq \chi(1)$ for all $\chi \in \text{Irr}(G)$.

Field of values

For $\chi \in \text{Irr}(G)$ let $f(\chi)$ be the smallest integer n such that $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_n$.

Conjecture (HUNG–TIEP)

We have $|\mathbb{Q}_{f(\chi)} : \mathbb{Q}(\chi)| \leq \chi(1)$ for all $\chi \in \text{Irr}(G)$.

Conjecture holds whenever $\chi(1)$ is a prime.

Field of values

For $\chi \in \text{Irr}(G)$ let $f(\chi)$ be the smallest integer n such that $\mathbb{Q}(\chi) \subseteq \mathbb{Q}_n$.

Conjecture (HUNG–TIEP)

We have $|\mathbb{Q}_{f(\chi)} : \mathbb{Q}(\chi)| \leq \chi(1)$ for all $\chi \in \text{Irr}(G)$.

Conjecture holds whenever $\chi(1)$ is a prime.

Problem (HUNG–TIEP)

Let $z = \zeta_1 + \dots + \zeta_n \in \mathbb{C}$ be a sum of roots of unity. Let $c(z)$ be the smallest n such that $z \in \mathbb{Q}_n$. Is $|\mathbb{Q}_{c(z)} : \mathbb{Q}(z)| \leq n$?

Algebraic integers

- Let \mathbb{Z}_G be the ring of integers of $\mathbb{Q}(G)$.

Algebraic integers

- Let \mathbb{Z}_G be the ring of integers of $\mathbb{Q}(G)$.
- Let $\mathbb{Z}[G] := \mathbb{Z}[\chi(g) : \chi \in \text{Irr}(G), g \in G] \subseteq \mathbb{Z}_G$.

Algebraic integers

- Let \mathbb{Z}_G be the ring of integers of $\mathbb{Q}(G)$.
- Let $\mathbb{Z}[G] := \mathbb{Z}[\chi(g) : \chi \in \text{Irr}(G), g \in G] \subseteq \mathbb{Z}_G$.
- Then $\mathbb{Z}_G/\mathbb{Z}[G]$ is a finite abelian group.

Algebraic integers

- Let \mathbb{Z}_G be the ring of integers of $\mathbb{Q}(G)$.
- Let $\mathbb{Z}[G] := \mathbb{Z}[\chi(g) : \chi \in \text{Irr}(G), g \in G] \subseteq \mathbb{Z}_G$.
- Then $\mathbb{Z}_G/\mathbb{Z}[G]$ is a finite abelian group.

Conjecture (BÄCHLE–S.)

The exponent of $\mathbb{Z}_G/\mathbb{Z}[G]$ divides $|G|$.

Algebraic integers

- Let \mathbb{Z}_G be the ring of integers of $\mathbb{Q}(G)$.
- Let $\mathbb{Z}[G] := \mathbb{Z}[\chi(g) : \chi \in \text{Irr}(G), g \in G] \subseteq \mathbb{Z}_G$.
- Then $\mathbb{Z}_G/\mathbb{Z}[G]$ is a finite abelian group.

Conjecture (BÄCHLE–S.)

The exponent of $\mathbb{Z}_G/\mathbb{Z}[G]$ divides $|G|$.

Example

Conjecture hold for nilpotent groups. For $G = C_{15} \rtimes D_{16}$,

$$\mathbb{Z}_G/\mathbb{Z}[G] \cong C_{120}^2 \times C_{60}^2 \times C_{12}^4 \times C_4^4 \times C_2^{14}.$$

Weights

- Let π be a set of primes.

Weights

- Let π be a set of primes.
- Let $l(G)$ be the number of conjugacy classes of π' -elements of G .

Weights

- Let π be a set of primes.
- Let $l(G)$ be the number of conjugacy classes of π' -elements of G .
- Let $\chi \in \text{Irr}(G)$ of π -**defect** 0 if $|G|/\chi(1)$ is a π' -number.

Weights

- Let π be a set of primes.
- Let $l(G)$ be the number of conjugacy classes of π' -elements of G .
- Let $\chi \in \text{Irr}(G)$ of π -**defect** 0 if $|G|/\chi(1)$ is a π' -number.
- Let $k^0(G)$ be the number of π -defect 0 characters of G .

Weights

- Let π be a set of primes.
- Let $l(G)$ be the number of conjugacy classes of π' -elements of G .
- Let $\chi \in \text{Irr}(G)$ of π -**defect** 0 if $|G|/\chi(1)$ is a π' -number.
- Let $k^0(G)$ be the number of π -defect 0 characters of G .

Theorem (NAVARRO–S.)

If G is π -separable with solvable Hall π -subgroups, then

$$l(G) = \sum_P k^0(N_G(P)/P)$$

where P runs through the nilpotent π -subgroups of G up to conjugation.

Weights

- Let π be a set of primes.
- Let $l(G)$ be the number of conjugacy classes of π' -elements of G .
- Let $\chi \in \text{Irr}(G)$ of π -**defect** 0 if $|G|/\chi(1)$ is a π' -number.
- Let $k^0(G)$ be the number of π -defect 0 characters of G .

Theorem (NAVARRO–S.)

If G is π -separable with solvable Hall π -subgroups, then

$$l(G) = \sum_P k^0(N_G(P)/P)$$

where P runs through the nilpotent π -subgroups of G up to conjugation.

For $\pi = \{p\}$, this is Alperin's weight conjecture for p -solvable groups.

Weights

Definition

Define the **weight** of a group P by

$$w(P) = 1 - \sum_{\substack{Q < P \\ N_P(Q) = Q}} w(Q) \quad (Q \text{ up to conjugation}).$$

Weights

Definition

Define the **weight** of a group P by

$$w(P) = 1 - \sum_{\substack{Q < P \\ N_P(Q) = Q}} w(Q) \quad (Q \text{ up to conjugation}).$$

Conjecture (NAVARRO–S.)

For every π -separable group G ,

$$l(G) = \sum_P w(P) k^0(N_G(P)/P)$$

where P runs through the π -subgroups of G up to conjugation.

Weights

Example

- If P is nilpotent, then $w(P) = 1$.

Weights

Example

- If P is nilpotent, then $w(P) = 1$.
- If P is solvable, but not nilpotent, then $w(P) = 0$ (CARTER subgroups).

Weights

Example

- If P is nilpotent, then $w(P) = 1$.
- If P is solvable, but not nilpotent, then $w(P) = 0$ (CARTER subgroups).
- On the other hand,

P	A_5	A_6	A_7	A_8	$\text{PSL}(3, 4)$	A_5^2	$S_n (3 \leq n \leq 11)$
$w(P)$	1	-2	1	-1	6	-1	0

Characterization of blocks

Let G^0 be the set of p -regular elements of G .

Characterization of blocks

Let G^0 be the set of p -regular elements of G .

Theorem (OSIMA)

A subset $J \subseteq \text{Irr}(G)$ is a union of p -blocks if and only if

$$\sum_{\chi \in J} \chi(g)\chi(h) = 0 \quad (\forall g \in G^0, h \in G \setminus G^0).$$

Characterization of blocks

Let G^0 be the set of p -regular elements of G .

Theorem (OSIMA)

A subset $J \subseteq \text{Irr}(G)$ is a union of p -blocks if and only if

$$\sum_{\chi \in J} \chi(g)\chi(h) = 0 \quad (\forall g \in G^0, h \in G \setminus G^0).$$

Conjecture (HARADA)

It is enough to fix $g = 1$ in Osima's theorem.

Basic sets

- Let B be a p -block of G .

Basic sets

- Let B be a p -block of G .
- For $\chi \in \text{Irr}(G)$ let χ^0 be the restriction to G^0 .

Basic sets

- Let B be a p -block of G .
- For $\chi \in \text{Irr}(G)$ let χ^0 be the restriction to G^0 .
- The Brauer characters in p -solvable groups have the form χ^0 (FONG–SWAN).

Basic sets

- Let B be a p -block of G .
- For $\chi \in \text{Irr}(G)$ let χ^0 be the restriction to G^0 .
- The Brauer characters in p -solvable groups have the form χ^0 (FONG–SWAN).

Conjecture (ordinary basic set, GECK)

There exist $\chi_1, \dots, \chi_l \in \text{Irr}(B)$ such that $\chi_1^0, \dots, \chi_l^0$ is a \mathbb{Z} -basis for the ring of generalized Brauer characters of B .

Basic sets

- Let B be a p -block of G .
- For $\chi \in \text{Irr}(G)$ let χ^0 be the restriction to G^0 .
- The Brauer characters in p -solvable groups have the form χ^0 (FONG–SWAN).

Conjecture (ordinary basic set, GECK)

There exist $\chi_1, \dots, \chi_l \in \text{Irr}(B)$ such that $\chi_1^0, \dots, \chi_l^0$ is a \mathbb{Z} -basis for the ring of generalized Brauer characters of B .

This holds in a strong sense for “most” groups of Lie type (BRUNAT–DUDAS–TAYLOR).

Number of characters

- For $\chi, \psi \in \text{Irr}(G)$ let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

Number of characters

- For $\chi, \psi \in \text{Irr}(G)$ let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

- Let d be the defect of B . Then $p^d[\chi, \chi]^0 \in \mathbb{N}$ for all $\chi \in \text{Irr}(B)$.

Number of characters

- For $\chi, \psi \in \text{Irr}(G)$ let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

- Let d be the defect of B . Then $p^d[\chi, \chi]^0 \in \mathbb{N}$ for all $\chi \in \text{Irr}(B)$.

Conjecture (NAVARRO-S.)

We have $p^d[\chi, \chi]^0 \geq l(B)$ for all $\chi \in \text{Irr}(B)$.

Number of characters

- For $\chi, \psi \in \text{Irr}(G)$ let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

- Let d be the defect of B . Then $p^d[\chi, \chi]^0 \in \mathbb{N}$ for all $\chi \in \text{Irr}(B)$.

Conjecture (NAVARRO-S.)

We have $p^d[\chi, \chi]^0 \geq l(B)$ for all $\chi \in \text{Irr}(B)$.

- This implies Brauer's conjecture $k(B) \leq p^d$ with equality only if B has abelian defect groups.

Number of characters

- For $\chi, \psi \in \text{Irr}(G)$ let

$$[\chi, \psi]^0 = \frac{1}{|G|} \sum_{g \in G^0} \chi(g) \overline{\psi(g)}.$$

- Let d be the defect of B . Then $p^d[\chi, \chi]^0 \in \mathbb{N}$ for all $\chi \in \text{Irr}(B)$.

Conjecture (NAVARRO-S.)

We have $p^d[\chi, \chi]^0 \geq l(B)$ for all $\chi \in \text{Irr}(B)$.

- This implies Brauer's conjecture $k(B) \leq p^d$ with equality only if B has abelian defect groups.
- The case $\chi = 1_G$ was conjectured by MURAI and implies FROBENIUS conjecture: G is p -nilpotent iff $|G^0| = |G|_{p'}$ (known via CFSG).

Fusion numbers

- Let

$$X_B := (\chi(g_i) : \chi \in \text{Irr}(B), i = 1, \dots, k(G))$$

be a submatrix of the character table.

Fusion numbers

- Let

$$X_B := (\chi(g_i) : \chi \in \text{Irr}(B), i = 1, \dots, k(G))$$

be a submatrix of the character table.

- The non-zero elementary divisors of $X_B^t \overline{X_B}$ over a complete discrete valuation ring are p -powers e_1, \dots, e_t .

Fusion numbers

- Let

$$X_B := (\chi(g_i) : \chi \in \text{Irr}(B), i = 1, \dots, k(G))$$

be a submatrix of the character table.

- The non-zero elementary divisors of $X_B^t \overline{X_B}$ over a complete discrete valuation ring are p -powers e_1, \dots, e_t .

Conjecture (S.)

We have $\gamma(B) := \frac{1}{e_1} + \dots + \frac{1}{e_t} \geq 1$ with equality if and only if B is nilpotent.

Fusion numbers

- Let

$$X_B := (\chi(g_i) : \chi \in \text{Irr}(B), i = 1, \dots, k(G))$$

be a submatrix of the character table.

- The non-zero elementary divisors of $X_B^t \overline{X_B}$ over a complete discrete valuation ring are p -powers e_1, \dots, e_t .

Conjecture (S.)

We have $\gamma(B) := \frac{1}{e_1} + \dots + \frac{1}{e_t} \geq 1$ with equality if and only if B is nilpotent.

Example

Conjecture holds for symmetric groups and “ATLAS groups”. For the principal 2-block B of the Monster, $\gamma(B) \approx 39.5$.

Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*

Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands.* [finishing his second book]

Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*

Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*
- *Silly? I am in the middle of a restaurant.*

Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*
- *Silly? I am in the middle of a restaurant.*
- *Seriously, I have thought on this for 3 minutes.*

Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands.* [finishing his second book]
- *If these two guys are not Morita equivalent then I won't say the word Morita again!* [they weren't]
- *Silly? I am in the middle of a restaurant.*
- *Seriously, I have thought on this for 3 minutes.*
- *Zero.* [answer to: How many examples did you check?]

Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*
- *Silly? I am in the middle of a restaurant.*
- *Seriously, I have thought on this for 3 minutes.*
- *Zero. [answer to: How many examples did you check?]*
- *I am also amazed by my intuition, sorry to say!*

Some memories

- *I read your question just before going to bed and you have ruined my night. Good question. I dreamt that I had an example.*
- *I am two days, 18 hours each, trying to clean up the inductive McKay, no waving hands. [finishing his second book]*
- *If these two guys are not Morita equivalent then I won't say the word Morita again! [they weren't]*
- *Silly? I am in the middle of a restaurant.*
- *Seriously, I have thought on this for 3 minutes.*
- *Zero. [answer to: How many examples did you check?]*
- *I am also amazed by my intuition, sorry to say!*
- *25th anniversary of my wedding. I have promised my wife not to think on mathematics for a couple of days. Now you are making my promise easy to break. . .*