


Lighting


Light/Material Interaction




Color Multiplication – “*” Operator

- Color channels are independent
- S ...light color
- M ...material color


$$\begin{pmatrix} S_R \\ S_G \\ S_B \end{pmatrix} * \begin{pmatrix} M_R \\ M_G \\ M_B \end{pmatrix} := \begin{pmatrix} S_R M_R \\ S_G M_G \\ S_B M_B \end{pmatrix}$$




Color Multiplication – “*” Operator




Light Sources



Reflection Models




Shading Models




Light Sources

- Where is the light coming from?




Directional Light

- Light source is infinitely far away
- Light rays are parallel, like sun
- l ... direction to the light = $-dir$




Point Light

- Has a certain position in space $l = \text{normalize}(\text{light}_{pos} - pos)$



Spot Light


- Has a certain position and cone in space
- Cone can be specified by opening angle and central direction




Spot Light

- Point is in cone iff


$$\cos^{-1} \text{dot}(l, -\text{light}_{dir}) < \theta$$




Light Sources



Reflection Models



Shading Models




Reflection/Illumination/Lighting model






- How is light reflected by a surface?
- What is the resulting color?
- What properties can we simulate with a given model?
- What lighting effects can we create?




Glossy → Matte




Surface lighting effects



Lighting Models



Phong Illumination Model



component ... lighting model component

Ambient Light Reflection




- Background light
 - No direction because scattered so often
 - “Color inside shadow”
 - Approximation of global diffuse lighting effects
- $S_{ambient}$...background light color
- M ...material color

$$ambient = M * S_{ambient}$$

Diffuse Light Reflection




- The flatter light falls on a surface, the darker it will appear
- Ideal diffuse reflectors (Lambertian reflectors)
- Brightness depends on orientation of surface



Diffuse Light Reflection





- Diffuse brightness is dependent on angle between
 n ... surface normal and
 l ... direction to the light



Diffuse Light Reflection




- Diffuse brightness is dependent on angle between
 n ... surface normal and
 l ... direction to the light
- $\cos \theta = \text{dot}(n, l)$



Diffuse Light Reflection




- S ...light color
- M ...material color
- $diffuse = M * S \cdot \text{dot}(n, l)$



Light from Behind





- Should be ignored
- $diffuse = M * S \cdot \max(0, \text{dot}(\mathbf{n}, \mathbf{l}))$



Lambertian (Diffuse) Reflection




Specular Highlights




Light source reflection
directly into the viewer's eye

Specular Reflection Model


- Reflection of incident light around specular-reflection angle




- Empirical Phong model
- $$\text{specular} = M * S \cdot \cos^p \sigma$$
$$= M * S \cdot \text{dot}(\mathbf{r}, \mathbf{v})^p$$




Specular Reflection Coefficient p




Specular Reflection Coefficient p




Specular Reflection Coefficient p



Specular Reflection Coefficient p





Specular Reflection Coefficient p




Specular Reflection Coefficient


- Empirical Phong model specular = $M * S \cdot \cos^p \sigma$
 - p large \Rightarrow shiny surface
 - p small \Rightarrow dull surface



Specular reflection



$$\text{specular} = M * S \cdot \text{dot}(r, v)^p$$




$$r + l = 2 \cdot \text{dot}(n, l) \cdot n$$

$$r = 2 \cdot \text{dot}(n, l) \cdot n - l$$

Blinn-Phong



- Halfway vector **h**

$$L_{\text{spec}} = M * S \cdot (\mathbf{v} \cdot \mathbf{r})^p \rightarrow L_{\text{spec}} = M * S \cdot (\mathbf{n} \cdot \mathbf{h})^p$$




$$\mathbf{h} = \frac{\mathbf{l} + \mathbf{v}}{\|\mathbf{l} + \mathbf{v}\|}$$

Phong vs Blinn-Phong




Lighting Models




Gooch

- Blend between a cool and a warm color




Toon shading

- Discrete color steps for diffuse




Cel shading


- Detect edges and color them




Light Sources



Reflection Models



Shading Models



Shading Models




Shading model

- Shading \neq shadows (shadowing)
- Coloring / shading the model
- When to evaluate lighting model





Flat-shading

- 1 color for the mesh (polygon)
- Really fast
- Really ugly
- *If an object really is faceted, is this accurate?*
- No:
 - Point light sources
 - Direction to light varies across the facet
 - Specular reflectance
 - Direction to eye varies across the facet





Gouraud shading

1. Calculate the normal vector for each vertex
2. Calculate the intensity for each vertex




Vertex Normals

- Vertex normals may be
 - Provided with the model
 - Artist
 - 3d program
 - Computed from first principles
 - Mathematic description of model




Sharp Edges and Normals



Vertex Normals


- Approximated by averaging the normals of the facets that share the vertex



$$\mathbf{n}_v = \frac{\sum_{k=1}^N \mathbf{n}_k}{\left\| \sum_{k=1}^N \mathbf{n}_k \right\|}$$



Gouraud shading

1. Calculate the normal vector for each vertex
2. Calculate the intensity for each vertex
3. Color interpolation along edges





Gouraud shading

1. Calculate the normal vector for each vertex
2. Calculate the intensity for each vertex
3. Color interpolation along edges
4. Color interpolation along scanline




Phong Shading Model


1. Normal Vector for each vertex
2. Normal vectors are interpolated along the edge
3. Normal vectors are interpolated along the scanline
4. Calculate the intensity using the normal vectors



Normal Interpolation



Flat / Gouraud / Phong Comparison



Transforming Normals

- Differential scaling changes shape and normals
- If \mathbf{M} transforms points, then $(\mathbf{M}^T)^{-1}$ transforms normals

