

When the membership and homogeneous-existence loci coincide, and a proof that they do for ansatz 5 on hydrogen

July 23, 2026

A companion note to *A New Method of Solving PDEs*, and a sequel to *Membership versus variety, and the partial-solution strata* and *Recovering the partial-solution strata by membership alone: a GCRD closure construction*. The core algorithm computes the **membership locus** $V_V(A) = \{c : P \in [A(c)]\}$; the existential computation on $A \cup \{P\}$ computes the **homogeneous existence locus** $V_{\exists \setminus \Psi}(A) = \{c : 1 \notin [A(c), P] : \Psi^\infty\}$. This note (i) records the exact condition under which the two coincide, (ii) proves coincidence unconditionally for one-dimensional ansätze, and (iii) proves by an explicit greatest-common-right-divisor (GCRD) computation that *ansatz 5 on the hydrogen Schrödinger equation has no genuine partial strata*, so that the two loci coincide there off the bad locus, and the whole gap between them is confined to the ODE-degeneration loci that the differential Thomas decomposition already resolves cell by cell.

1 The exact gap, and its GCRD reading

The relationship between the two loci was established in the two companion notes. Writing the existential projection as membership plus a remainder,

$$V_{\exists \setminus \Psi}(A) = V_V(A) \sqcup \{\text{partial-solution strata}\}, \quad (1)$$

where a *partial-solution stratum* (companion note I, Definition 1) is a c at which a proper nontrivial subfamily of $\text{Sol}(A(c))$ satisfies P while the family as a whole does not. Hence

$$\boxed{V_V(A) = V_{\exists \setminus \Psi}(A) \iff A \text{ has no partial-solution strata.}} \quad (2)$$

For an ansatz whose top element is a *linear* ODE L of order n in the inner variable v , companion note II sharpens this into an order condition. Ritt-reducing P modulo A and splitting the remainder over base power products yields operators $R_1, \dots, R_s \in K(v)[\partial]$ of order $\leq n - 1$; with

$$G(c^*) = \text{GCRD}(L(c^*), R_1(c^*), \dots, R_s(c^*)), \quad k = \text{ord } G(c^*),$$

the trichotomy (note II, Theorem 1) reads: $k = n \iff c^* \in V_V$ (the whole family solves P); $k = 0 \iff$ no common solution; and $0 < k < n \iff c^*$ is a partial stratum. Therefore

$$V_V(A) = V_{\exists \setminus \Psi}(A) \quad \text{off the bad locus} \iff \text{ord } G(c^*) \in \{0, n\} \quad \text{for all such } c^*. \quad (3)$$

Coincidence is exactly the statement that the GCRD never has intermediate order.

4 Summary

- $V_{\forall} = V_{\exists \setminus \Psi} \iff$ no partial strata $\iff$ (linear-ODE ansätze) $\text{ord GCRD}(L, \{R_{\beta}\}) \in \{0, n\}$ everywhere ((2), (3)).
- One-dimensional ansätze coincide unconditionally (Lemma 1); this covers the first-order and pure-exponential library families.
- Ansatz 5 on hydrogen has no genuine ($h \neq 0$) partial strata (Theorem 1); its $V_{\forall}$ -vs- $V_{\exists \setminus \Psi}$ gap is entirely on the ODE-degeneration loci resolved by the Thomas decomposition, so per-cell membership recovers $V_{\exists \setminus \Psi}$ with no existential computation.
- Open: the log ansätze dimension check (2, 17, 17.1, 18, 19); the quadratic-coefficient and coefficient-ring second-order families; the nested (12) and product (10) ansätze; and the helium Hamiltonian.