

MAT 472 - Intermediate Real Analysis I

Instructor: Dr. Steven Kaliszewski

Notes written by Brett Hansen

Contents

1	Week of August 21st, 2016	2
1.1	The Completeness Property of Real Numbers	2
1.1.1	Axiom of Completeness	2
2	Week of August 28th, 2016	3
3	Week of September 4th, 2016	3
4	Week of September 11th, 2016	3
5	Week of September 18th, 2016	3
6	Week of September 25th, 2016	3
7	Week of October 2nd, 2016	3
8	Week of October 9th, 2016	3
9	Week of October 16th, 2016	3
10	Week of October 23rd, 2016	3
11	Week of October 30th, 2016	3
12	Week of November 6th, 2016	3
13	Week of November 13th, 2016	3
14	Week of November 20th, 2016	3
15	Week of November 27th, 2016	3

1 Week of August 21st, 2016

1.1 The Completeness Property of Real Numbers

\mathbb{R} and \mathbb{Q} are ordered fields.

- $|\mathbb{Q}| < |\mathbb{R}|$
- \mathbb{Q} is countable, \mathbb{R} is uncountable (cardinality)
- $x^2 - 2 = 0$ has solutions in \mathbb{R} but none in \mathbb{Q}
- \mathbb{R} has no gaps (order)

1.1.1 Axiom of Completeness

Every non-empty subset of \mathbb{R} that has an upper bound has a least upper bound. Along with the greatest lower bound, these bounds can be classified using the following definitions:

Upper Bound Let the set S be a non-empty subset of \mathbb{R} and suppose $b \in \mathbb{R}$. b is an upper bound of the set S if $\forall x \in S, x \leq b$.

Lower Bound Let the set S be a non-empty subset of \mathbb{R} and suppose $b \in \mathbb{R}$. b is a lower bound of the set S if $\forall x \in S, x \geq b$.

Supremum Let B be the set of upper bounds of the non-empty subset, S , of \mathbb{R} and suppose $b \in B$. If $\forall x \in B, b \leq x$, then b is the least upper bound of S , or *supremum* of S . This can be denoted as $\sup S = b$.

Infinum Let B be the set of lower bounds of the non-empty subset, S , of \mathbb{R} and suppose $b \in B$. If $\forall x \in B, b \geq x$, then b is the greatest lower bound of S , or *infinum* of S . This can be denoted as $\inf S = b$.

Example $A = [0, 2] \longrightarrow \sup A = 2$
 $A = [0, 2) \longrightarrow \sup A = 2$

Maximum If the supremum of a set is also a member of the set, then the supremum is also the maximum.

Example Let $A = \{1/n \mid n \in \mathbb{N}\}$, then $\max A = \sup A = 1$ and $\inf A = 0$, however there is no minimum.

There are non-empty bounded subsets of \mathbb{Q} with no supremum (or infimum) in \mathbb{Q} .

Example Let $S = \{r \in \mathbb{Q} \mid r^2 < 2\}$. There does not exist any $b \in \mathbb{Q}$ such that $b \leq c$ where $c \in \text{UB}\{S\}$ and $c \in \mathbb{Q}$.

Proof S is non-empty and bounded above by 2. Suppose $x \in \mathbb{Q}$ and $x > 2$. Then $0 \leq 2 < x$ so $0 \leq 2 \cdot 2 < 2x$ and $0 \leq 2x < x^2$. Then $2 < 4 < x^2$ and thus $x \notin S$.

QED

- 2 Week of August 28th, 2016
- 3 Week of September 4th, 2016
- 4 Week of September 11th, 2016
- 5 Week of September 18th, 2016
- 6 Week of September 25th, 2016
- 7 Week of October 2nd, 2016
- 8 Week of October 9th, 2016
- 9 Week of October 16th, 2016
- 10 Week of October 23rd, 2016
- 11 Week of October 30th, 2016
- 12 Week of November 6th, 2016
- 13 Week of November 13th, 2016
- 14 Week of November 20th, 2016
- 15 Week of November 27th, 2016