

INTRO to DATA SCIENCE

LECTURE 6: REGRESSION & REGULARIZATION

Francesco Mosconi
DAT16 SF // August 17, 2015

INTRO TO DATA SCIENCE, REGRESSION & REGULARIZATION

DATA SCIENCE IN THE NEWS

DATA SCIENCE IN THE NEWS

WHY SECURITY DATA SCIENCE MATTERS AND HOW ITS
DIFFERENT: PITFALLS AND PROMISES OF DATA
SCIENCE BASED BREACH DETECTION AND THREAT
INTELLIGENCE



FROM FALSE POSITIVES TO ACTIONABLE ANALYSIS:
BEHAVIORAL INTRUSION DETECTION MACHINE
LEARNING AND THE SOC

THE APPLICATIONS OF DEEP LEARNING ON TRAFFIC
IDENTIFICATION

Why security data science matters and how it's different: pitfalls and promises of data science based breach detection and threat intelligence

Joshua Saxe, Invincea Labs

Work presented in this talk contains significant contributions from Alex Long, David Slater, Giacomo Bergamo, Konstantin Berlin, and Robert Gove

DATA SCIENCE IN THE NEWS



CLASSIFICATION OF ENCRYPTED WEB TRAFFIC USING MACHINE LEARNING ALGORITHMS

THESIS

William Charles Barto

AFIT-ENG-13-J-11

Cleaning MoMA's Artwork Collection Using Python

09 AUG 2015 on python, art, and cleaning

Use Python to Clean the Museum of Modern Art's Collection

Art is a messy business. Over centuries, artists have created everything from simple paintings to complex sculptures, and art historians have been cataloging everything they can along the way. The Museum of Modern Art, or MoMA for short, is considered one of the most influential museums in the world and recently released a dataset of all the artworks they've cataloged in their collection. This dataset contains basic information on metadata for each artwork and is part of MoMA's push to make art more accessible to everyone.

<https://www.dataquest.io/blog/use-python-to-clean-moma-collection/>

LAST TIME:

I. CROSS VALIDATION



II. INTRO TO PROBABILITY

III. NAÏVE BAYESIAN CLASSIFICATION

EXERCISES:

IV. NAÏVE BAYES CLASSIFICATION IN PYTHON

QUESTIONS?



INTRO TO DATA SCIENCE

QUESTIONS?

WHAT WAS THE MOST INTERESTING THING YOU LEARNT?

WHAT WAS THE HARDEST TO GRASP?

REVIEW GAME



AGENDA

I. LINEAR REGRESSION (INCL. MULTIPLE REGRESSION)

II. POLYNOMIAL REGRESSION

III. REGULARIZATION

LAB:

**IV. IMPLEMENTING MULTIPLE REGRESSION & POLYNOMIAL
REGRESSION IN PYTHON**

KEY OBJECTIVES

- **WHAT ARE LINEAR AND POLYNOMIAL REGRESSION**
- **WHICH PROBLEMS CAN BE TACKLED WITH REGRESSION TECHNIQUES**
- **HOW TO IMPLEMENT LINEAR AND POLYNOMIAL REGRESSION IN PYTHON**
- **WHAT IS REGULARIZATION**
- **HOW REGULARIZATION CAN HELP WHEN DATA IS NOISY**
- **HOW TO IMPLEMENT REGULARIZATION IN PYTHON**

I. LINEAR REGRESSION

	<i>Continuous</i>	<i>Categorical</i>
<i>Supervised</i>	???	???
<i>Unsupervised</i>	???	???

	<i>Continuous</i>	<i>Categorical</i>
<i>Supervised</i>	<i>regression</i>	<i>classification</i>
<i>Unsupervised</i>	<i>dimension reduction</i>	<i>clustering</i>

Q: What is a regression model?

Q: What is a regression model?

A: A functional relationship between input & response variables.

Q: What is a regression model?

A: A functional relationship between input & response variables

The simple linear regression model captures a linear relationship between a single input variable x and a response variable y :

Q: What is a regression model?

A: A functional relationship between input & response variables

The simple linear regression model captures a linear relationship between a single input variable x and a response variable y :

$$y = \alpha + \beta x + \varepsilon$$

Q: What do the terms in this model mean?

$$y = \alpha + \beta x + \varepsilon$$

Q: What do the terms in this model mean?

$$y = \alpha + \beta X + \varepsilon$$

A: y = response variable (the one we want to predict)

Q: What do the terms in this model mean?

$$y = \alpha + \beta x + \varepsilon$$

A: y = response variable (the one we want to predict)

x = input variable (the one we use to train the model)

Q: What do the terms in this model mean?

$$y = \alpha + \beta x + \varepsilon$$

A: y = response variable (the one we want to predict)

x = input variable (the one we use to train the model)

α = intercept (where the line crosses the y-axis)

Q: What do the terms in this model mean?

$$y = \alpha + \beta x + \varepsilon$$

A: y = response variable (the one we want to predict)

x = input variable (the one we use to train the model)

α = intercept (where the line crosses the y-axis)

β = regression coefficient (the model “parameter”)

Q: What do the terms in this model mean?

$$y = \alpha + \beta x + \varepsilon$$

A: y = response variable (the one we want to predict)

x = input variable (the one we use to train the model)

α = intercept (where the line crosses the y -axis)

β = regression coefficient (the model “parameter”)

ε = residual (the prediction error)

We can extend this model to several input variables, giving us the multiple linear regression model:

We can extend this model to several input variables, giving us the multiple linear regression model:

$$y = \alpha + \beta_1 x_1 + \dots + \beta_n x_n + \varepsilon$$



We can extend this model to several input variables, giving us the multiple linear regression model:

$$y = \alpha + \beta_1 x_1 + \dots + \beta_n x_n + \varepsilon$$

$$y = \alpha + \begin{bmatrix} \beta_1 & \beta_2 & \dots & \beta_n \end{bmatrix} * \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix} + \varepsilon$$

Linear regression involves several technical assumptions and is often presented with lots of mathematical formality.

The math is not very important for our purposes, but you should check it out if you get serious about solving regression problems.



Q: How do we fit a regression model to a dataset?

Q: How do we fit a regression model to a dataset?

A: In theory, minimize the sum of the squared residuals (OLS).

Q: How do we fit a regression model to a dataset?

A: In theory, minimize the sum of the squared residuals (OLS).

In practice, any respectable piece of software will do this for you.

Q: How do we fit a regression model to a dataset?

A: In theory, minimize the sum of the squared residuals (OLS).

In practice, any respectable piece of software will do this for you.

But again, if you get serious about regression, you should learn how this works!


Q: How do we fit a regression model to a dataset?

A: In theory, minimize the sum of the squared residuals (OLS).

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i.$$


$$\hat{u}_i = y_i - \hat{y}_i = y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i.$$

$$\sum_{i=1}^n \hat{u}_i^2 = \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2,$$



Q: How do we fit a regression model to a dataset?

A: In theory, minimize the sum of the squared residuals (OLS).



II: POLYNOMIAL REGRESSION

Consider the following polynomial regression model:

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \varepsilon$$

Consider the following polynomial regression model:

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \varepsilon$$

Q: This represents a nonlinear relationship. Is it still a linear model?

Consider the following polynomial regression model:

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \varepsilon$$

Q: This represents a nonlinear relationship. Is it still a linear model?

A: Yes, because it's linear in the β 's!


Consider the following polynomial regression model:

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \varepsilon$$

Q: This represents a nonlinear relationship. Is it still a linear model?

A: Yes, because it's linear in the β 's!

“Although polynomial regression fits a nonlinear model to the data, as a statistical estimation problem it is linear, in the sense that the regression function $E(y|x)$ is linear in the unknown parameters that are estimated from the data. For this reason, polynomial regression is considered to be a special case of multiple linear regression.” -- Wikipedia



Polynomial regression allows us to fit very complex curves to data.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

Polynomial regression allows us to fit very complex curves to data.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

But there is one problem with the model we've written down so far.

Polynomial regression allows us to fit very complex curves to data.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

But there is one problem with the model we've written down so far.

Q: Does anyone know what it is?

Polynomial regression allows us to fit very complex curves to data.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

But there is one problem with the model we've written down so far.

Q: Does anyone know what it is?

A: This model violates one of the assumptions of linear regression!



This model displays multicollinearity, which means the predictor variables are highly correlated with each other.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

Multicollinearity causes the linear regression model to break down, because it can't tell the predictor variables apart.

Q: What can we do about this?

Q: What can we do about this?

A: Replace the correlated predictors with uncorrelated predictors.

Q: What can we do about this?

A: Replace the correlated predictors with uncorrelated predictors.

$$y = \alpha + \beta_1 f_1(x) + \beta_2 f_2(x^2) + \dots + \beta_n f_n(x^n) + \varepsilon$$

Q: What can we do about this?

A: Replace the correlated predictors with uncorrelated predictors.

$$y = \alpha + \beta_1 f_1(x) + \beta_2 f_2(x^2) + \dots + \beta_n f_n(x^n) + \varepsilon$$

OPTIONAL NOTE

These polynomial functions form an *orthogonal basis* of the function space.

So far, we've seen how polynomial regression allows us to fit complex nonlinear relationships, and even to avoid multicollinearity (by using basis functions).

So far, we've seen how polynomial regression allows us to fit complex nonlinear relationships, and even to avoid multicollinearity (by using basis functions).

Q: Can a regression model be too complex?

III: REGULARIZATION

Recall our earlier discussion of overfitting.


Recall our earlier discussion of overfitting.

When we talked about this in the context of classification, we said that it was a result of matching the training set too closely.

Recall our earlier discussion of overfitting.

When we talked about this in the context of classification, we said that it was a result of matching the training set too closely.


In other words, an overfit model matches the noise in the dataset instead of the signal.



The same thing can happen in regression.

It's possible to design a regression model that matches the noise in the data instead of the signal.

This happens when our model becomes too complex for the data to support.



Q: How do we define the complexity of a regression model?

Q: How do we define the complexity of a regression model?

A: One method is to define complexity as a function of the size of the coefficients.

Q: How do we define the complexity of a regression model?

A: One method is to define complexity as a function of the size of the coefficients.

Ex 1: $\sum |\beta_i|$

Ex 2: $\sum \beta_i^2$

Q: How do we define the complexity of a regression model?

A: One method is to define complexity as a function of the size of the coefficients.

Ex 1: $\sum |\beta_i|$ this is called the L1-norm

Ex 2: $\sum \beta_i^2$ this is called the L2-norm

These measures of complexity lead to the following regularization techniques:

These measures of complexity lead to the following regularization techniques:

L1 regularization: $y = \sum \beta_i x_i + \varepsilon \quad \text{st.} \quad \sum |\beta_i| < s$

These measures of complexity lead to the following regularization techniques:

L1 regularization: $y = \sum \beta_i x_i + \varepsilon \quad \text{st.} \quad \sum |\beta_i| < s$

L2 regularization: $y = \sum \beta_i x_i + \varepsilon \quad \text{st.} \quad \sum \beta_i^2 < s$

These measures of complexity lead to the following regularization techniques:

L1 regularization: $y = \sum \beta_i x_i + \varepsilon \quad \text{st.} \quad \sum |\beta_i| < s$

L2 regularization: $y = \sum \beta_i x_i + \varepsilon \quad \text{st.} \quad \sum \beta_i^2 < s$

Regularization refers to the method of preventing overfitting by explicitly controlling model complexity.

These regularization problems can also be expressed as:

L1 regularization: $\min(\|y - x\beta\|^2 + \lambda\|\beta\|)$

L2 regularization: $\min(\|y - x\beta\|^2 + \lambda\|\beta\|^2)$

These regularization problems can also be expressed as:

L1 regularization (Lasso): $\min(\|y - x\beta\|^2 + \lambda\|x\|)$

L2 regularization (Ridge): $\min(\|y - x\beta\|^2 + \lambda\|x\|^2)$

This (Lagrangian) formulation reflects the fact that there is a cost associated with regularization.

Q: What are bias and variance?


Q: What are bias and variance?

A: Bias refers to predictions that are systematically inaccurate.

Q: What are bias and variance?

A: Bias refers to predictions that are systematically inaccurate.

Variance refers to predictions that are generally inaccurate.




Q: What are bias and variance?

A: Bias refers to predictions that are systematically inaccurate.


Variance refers to predictions that are generally inaccurate.

It turns out (after some math) that the generalization error in our model can be decomposed into a bias component and variance component.

This is another example of the bias-variance tradeoff.



This is another example of the bias-variance tradeoff.

**NOTE**

The “meta-parameter” here is the λ we saw above.

A more typical term is “hyperparameter”.


This tradeoff is regulated by a hyperparameter λ , which we've already seen:

L1 regularization: $y = \sum \beta_i x_i + \varepsilon \quad \text{st.} \quad \sum |\beta_i| < \lambda$


L2 regularization: $y = \sum \beta_i x_i + \varepsilon \quad \text{st.} \quad \sum \beta_i^2 < \lambda$

So regularization represents a method to trade away some variance for a little bias in our model, thus achieving a better overall fit.


L2 regularization: $\min(\|y - x\beta\|^2 + \lambda\|x\|^2)$



L2 regularization: $\min(\|y - \mathbf{x}\beta\|^2 + \lambda\|\mathbf{x}\|^2)$



L2 regularization: $\min(\|y - \mathbf{x}\beta\|^2 + \lambda\|\mathbf{x}\|^2)$



- Linear regression
- Multiple regression
- Polynomial regression
- The concept of minimizing some error or “cost” function
- Regularization

LAB: POLYNOMIAL REGRESSION & REGULARIZATION