

Data Handling and Data Mining

FIXING THE SPARROW DATA SET

-

BASIC STATISTICS FOR BIOLOGISTS

Erik Kusch

PhD Student

Aarhus University

Department of Bioscience

Section for Ecoinformatics & Biodiversity

Center for Biodiversity and Dynamics in a Changing World (BIOCHANGE)

Ny Munkegade 116, Building 1540

8000 Aarhus

Denmark

email: erik@i-solution.de

Summary:

These are the solutions to the exercises contained within the handout to A Primer For Statistical Tests which walks you through the basics of variables, their scales and distributions. Keep in mind that there is probably a myriad of other ways to reach the same conclusions as presented in these solutions.

Contents

1	Loading the R Environment Object	2
2	Variables	2
2.1	Finding Variables	2
2.2	Colour	2
2.3	Depth	3
2.4	IndividualsPassingBy	4
2.5	Length	5
2.6	Reproducing	6
2.7	Sex	7
2.8	Size	8
2.9	Temperature	9
3	Distributions	10
3.1	Length	10
3.2	Reproducing	11
3.3	IndividualsPassingBy	12
3.4	Depth	13

1. Loading the R Environment Object

```
load("Data/Primer.RData") # load data file from Data folder
```

2. Variables

2.1 Finding Variables

```
ls() # list all elements in working environment
```


```
## [1] "Colour"          "Depth"           "IndividualsPassingBy"  
## [4] "Length"          "Reproducing"     "Sex"  
## [7] "Size"            "Temperature"
```

2.2 Colour

```
class(Colour) # mode
```

```
## [1] "character"
```

```
barplot(table(Colour)) # fitting?
```




Question	Answer
Mode?	character
Which scale?	Nominal
What's implied?	Categorical data that can't be ordered
Does data fit scale?	Yes

2.3 Depth

```
class(Depth) # mode

## [1] "numeric"

barplot(Depth) # fitting?
```




Question	Answer
Mode?	numeric
Which scale?	Interval/Discrete
What's implied?	Continuous data with a non-absence point of origin
Does data fit scale?	Debatable (is 0 depth absence of depth?)

2.4 IndividualsPassingBy

```
class(IndividualsPassingBy) # mode

## [1] "integer"

barplot(IndividualsPassingBy) # fitting?
```



Question	Answer
Mode?	integer
Which scale?	Integer
What's implied?	Only integer numbers with an absence point of origin
Does data fit scale?	Yes

2.5 Length

```
class(Length) # mode
```

```
## [1] "numeric"
```

```
barplot(Length) # fitting?
```




Question	Answer
Mode?	numeric
Which scale?	Relation/Ratio
What's implied?	Continuous data with an absence point of origin
Does data fit scale?	Yes

2.6 Reproducing

```
class(Reproducing) # mode
```

```
## [1] "integer"
```

```
barplot(Reproducing) # fitting?
```




Question	Answer
Mode?	integer
Which scale?	Integer
What's implied?	Only integer numbers with an absence point of origin
Does data fit scale?	Yes

2.7 Sex

```
class(Sex) # mode
```

```
## [1] "factor"
```

```
barplot(table(Sex)) # fitting?
```




Question	Answer
Mode?	factor
Which scale?	Binary
What's implied?	Only two possible outcomes
Does data fit scale?	Yes

2.8 Size

```
class(Size) # mode
```

```
## [1] "character"
```

```
barplot(table(Size)) # fitting?
```




Question	Answer
Mode?	character
Which scale?	Ordinal
What's implied?	Categorical data that can be ordered
Does data fit scale?	Yes

2.9 Temperature

```
class(Temperature) # mode

## [1] "numeric"

barplot(Temperature) # fitting?
```




Question	Answer
Mode?	numeric
Which scale?	Interval/Discrete
What's implied?	Continuous data with a non-absence point of origin
Does data fit scale?	Yes (the data is clearly recorded in degree Celsius)

3. Distributions

3.1 Length

```
plot(density(Length)) # distribution plot
```




```
shapiro.test(Length) # normality check
```

```
##  
## Shapiro-Wilk normality test  
##  
## data: Length  
## W = 1, p-value = 0.4
```

The data is **normal distributed**.

3.2 Reproducing

```
plot(density(Reproducing)) # distribution
```




```
shapiro.test(Reproducing) # normality check
```

```
##  
## Shapiro-Wilk normality test  
##  
## data: Reproducing  
## W = 1, p-value = 0.3
```

The data is **binomial distributed** (i.e. “How many individuals manage to reproduce”) but looks **normal distributed**. The normal distribution doesn’t make sense here because it implies continuity whilst the data only comes in integers.

3.3 IndividualsPassingBy

```
plot(density(IndividualsPassingBy)) # distribution
```




```
shapiro.test(IndividualsPassingBy) # normality check
```

```
##  
## Shapiro-Wilk normality test  
##  
## data:  IndividualsPassingBy  
## W = 1, p-value = 0.02
```

The data is **poisson distributed** (i.e. “How many individuals pass by an observer in a given time frame?”).

3.4 Depth

```
plot(density(Depth)) # distribution
```



The data is **uniform distributed**. You don't know this distribution class from the lectures and I only wanted to confuse you with this to show you that there's much more out there than I can show in our lectures.