F-Mail

Article Alert

Share/Bookmark

SUBSCRIBE OR RENEW

Includes NEJM iPad Edition, 20 FREE Online CME Exams and more >>



HOME

ARTICI ES & MUI TIMEDIA

ISSUES

SPECIAL TIES & TOPICS

FOR AUTHORS

CMF

Keyword, Title, Author, or Citation

Download Citation

Supplementary

TOOLS

PDF

Print



ORIGINAL ARTICLE

A Randomized, Controlled Trial of an Aerosolized Vaccine against Measles

Nicola Low, M.D., Ashish Bavdekar, D.N.B., Lakshmanan Jeyaseelan, Ph.D., Siddhivinayak Hirve, Ph.D., Kavitha Ramanathan, M.Sc., Nicholas J. Andrews, Ph.D., Naseem Shaikh, Ph.D., Ramesh S. Jadi, Ph.D., Arunachalam Rajagopal, M.C.A., Kevin E. Brown, M.D., David Brown, F.R.C.Path., James B. Fink, Ph.D., Oommen John, M.D., Pippa Scott, Ph.D., A. Ximena Riveros-Balta, B.Sc., Michel Greco, M.B.A., Rajeev Dhere, Ph.D., Prasad S. Kulkarni, M.D., and Ana Maria Henao Restrepo, M.D.

N Engl J Med 2015; 372:1519-1529 | April 16, 2015 | DOI: 10.1056/NEJMoa1407417

Share:

Abstract

Article

References

Citing Articles (4)

BACKGROUND

Aerosolized vaccine can be used as a needle-free method of immunization against measles, a disease that remains a major cause of illness and death. Data on the immunogenicity of aerosolized vaccine against measles in children are inconsistent.

Full Text of Background...

METHODS

We conducted an open-label noninferiority trial involving children 9.0 to 11.9 months of age in India who were eligible to receive a first dose of measles vaccine. Children were randomly assigned to receive a single dose of vaccine by means of either aerosol inhalation or a subcutaneous injection. The primary end points were seropositivity for antibodies against measles and adverse events 91 days after vaccination. The noninferiority margin was 5 percentage points.

Full Text of Methods...

RESULTS

A total of 1001 children were assigned to receive aerosolized vaccine, and 1003 children were assigned to receive subcutaneous vaccine; 1956 of all the children (97.6%) were followed to day 91, but outcome data were missing for 331 children because of thawed specimens. In the per-protocol population, data on 1560 of 2004 children (77.8%) could be evaluated. At day 91, a total of 662 of 775 children (85.4%; 95% confidence interval [CI], 82.5 to 88.0) in the aerosol group, as compared with 743 of 785 children (94.6%; 95% CI, 92.7 to 96.1) in the subcutaneous group, were seropositive, a difference of -9.2 percentage points (95% CI, -12.2 to -6.3). Findings were similar in the full-analysis set (673 of 788 children in the aerosol group [85.4%] and 754 of 796 children in the subcutaneous group [94.7%] were seropositive at day 91, a difference of -9.3 percentage points [95% CI, -12.3 to -6.4]) and after multiple imputation of missing results. No serious adverse events were attributable to measles vaccination. Adverse-event profiles were similar in the two groups.

Full Text of Results

MEDIA IN THIS ARTICLE

FIGURE 1



Eligibility, Randomization, and Follow-up of Children in the Per-Protocol

FIGURE 2



Immunogenicity Outcomes at Day 91 in the Aerosolized and Vaccine Groups

ARTICLE ACTIVITY

4 articles have cited this article

TOPICS

Vaccines **Viral Infections Global Health** Immunization Childhood Diseases

Research April 16, 2015

MORE IN

TRENDS

Most Viewed (Last Week)

Data Sharing [56,955 views] January 21, 2016 | D.L. Longo and J.M. Drazen

IMAGES IN CLINICAL MEDICINE

Taenia saginata Infestation [29,200 views] January 21, 2016 | J. Li and E. Guo

ORIGINAL ARTICLE

CDX2 as a Prognostic Biomarker in Stage II and Stage III Colon Cancer [27,821 views] January 21, 2016 | P. Dalerba and Others

More Trends

(NEIM CareerCenter

PHYSICIAN JOBS

January 29, 2016

Nephrology

NEPHROLOGISTS ARIZONA

Internal Medicine

Boston: Primary Care Provider MASSACHUSETTS

Obstetrics & Gynecology

Established OBGYN Practice - No State Income Tax TENNESSEE

Neurology

NEURORADIOI OGIST PENNSYI VANIA

Endocrinology

Endocrinologist NEW JERSEY

Surgery, General

CONCLUSIONS Aerosolized vaccine against measles was immunogenic, but, at the prespecified margin, the aerosolized vaccine was inferior to the subcutaneous vaccine with respect to the rate of seropositivity. (Funded by the Bill and Melinda Gates Foundation; Measles Aerosol Vaccine Project Clinical Trials Registry-India number, CTRI/2009/091/000673.)

Sign up for the FREE >> Weekly Table of Contents

Orthopedic Surgeon Jobs in Spokane, WA

WASHINGTON

Full Text of Discussion...

Read the Full Article...

CONTENT: Home | Current Issue | Articles | Issue Index | Specialties & Topics | Multimedia & Images | Archive 1812-1989

INFORMATION FOR: Authors | Reviewers | Subscribers | Institutions | Media | Advertisers

SERVICES: Subscribe | Renew | Pay Bill | Activate Subscription | Create or Manage Account | Alerts | RSS & Podcasts | Submit a Manuscript | Mobile

RESOURCES: Physician Jobs | Reprints | Permissions | Medical Meetings | Conventions | FAQs | NEJM Knowledge+ | NEJM Journal Watch | NEJM Catalyst | Help | Contact Us

NEJM: About | Product Information | Editors & Publishers | 200th Anniversary | Terms of Use | Privacy Policy | Copyright | Advertising Policies | NEJM Group

CME: Weekly CME Program | Browse Weekly Exams | Your CME Activity | Purchase Exams | Review CME Program









nejmcareercenter.org