

STAT W4201 001, Homework 3

Brian Weinstein (bmw2148)

Feb 17, 2016

Code is attached here and also posted at <https://github.com/BrianWeinstein/advanced-data-analysis>. Where relevant, code snippets and output are included in-line.

Problem 1: Ramsey 4.30

Problem 2: Ramsey 4.32

Problem 3: Ramsey 5.19

(a) The pooled estimate of the variance s_p^2 is given by

$$\begin{aligned} s_p^2 &= \frac{\sum_{i=1}^I (n_i - 1) s_i^2}{\sum_{i=1}^I (n_i - 1)} \\ &= \frac{(127 - 1)(0.4979)^2 + (44 - 1)(0.4235)^2 + (24 - 1)(0.3955)^2 + \dots}{(127 - 1) + (44 - 1) + (24 - 1) + \dots} \\ &\quad \frac{\dots + (41 - 1)(0.3183)^2 + (18 - 1)(0.3111)^2 + (16 - 1)(0.4649)^2 + \dots}{\dots + (41 - 1) + (18 - 1) + (16 - 1) + \dots} \\ &\quad \frac{\dots + (11 - 1)(0.2963)^2 + (7 - 1)(0.3242)^2 + (6 - 1)(0.5842)^2}{\dots + (11 - 1) + (7 - 1) + (6 - 1)} \\ &= 0.1919. \end{aligned}$$

(b) To construct an ANOVA table to test for species differences, we first compute SS_W , the sum of squared residuals of the “within” group.

$$\begin{aligned} SS_W &= \sum_{i=1}^9 \left[\sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_i)^2 \right] = \sum_{i=1}^9 [(n_i - 1) \text{SE}(Y_i)] \\ &= (127 - 1)(0.4979)^2 + (44 - 1)(0.4235)^2 + \dots + (6 - 1)(0.5842)^2 \\ &= 122.8658. \end{aligned}$$

Given the sample standard deviation of all 294 observations as one group is $\text{SD}_{Total} = 0.4962$, the total sum of squared residuals SS_{Total} is

Should
SS total
be -9 (-1)
instead
of -1 ?

$$\begin{aligned}
SS_{Total} &= \sum_{i=1}^9 \left[\sum_{j=1}^{n_i} (Y_{ij} - \bar{Y})^2 \right] = (n_1 + n_2 + \cdots + n_9 - 1)(SD_{Total}) \\
&= (127 + 44 + \cdots + 6 - 1)(0.4962) = (293)(0.4962) \\
&= 145.3866.
\end{aligned}$$

Therefore, the sum of squared residuals of the “between” group, SS_B is:

$$SS_B = SS_{Total} - SS_W = 145.3866 - 122.8658 = 22.5208.$$

The Total, Within, and Between degrees of freedom are $df_{Total} = (n - 1) = 293$, $df_W = (n - I) = (294 - 9) = 285$, and $df_B = df_{Total} - df_W = 8$, respectively.

The Mean Square for the Within and Between groups is the [(Sum of Squares) / df], and the F-statistic is defined as the [(Between Mean Square) / (Within Mean Square)].

Thus the ANOVA table to test for species differences is:

Source of Variation	Sum of Squares	d.f.	Mean Square	F-Statistic	p-Value
Between Groups	22.5208	8	2.8151	6.5300	8.0551×10^{-8}
Within Groups	122.8658	285	0.4311		
Total	145.3866	293			

Where the p-value for the F-Statistic is given by the CDF of an F distribution with 8 and 285 degrees of freedom.

```
> # p-value of F-statistic
> pf(q=6.5300, df1=8, df2=285, lower.tail=FALSE)
[1] 8.055137e-08
```

(c) The ANOVA method for calculating SS_B yields the same answer as the formula:

$$SS_B = \sum_{i=1}^I (n_i \bar{Y}_i^2) - n \bar{Y}^2 = \sum_{i=1}^9 (n_i \bar{Y}_i^2) - n \bar{Y}^2.$$

The overall mean $\bar{Y} = 7.4746$ is computed as a weighted average of the group means and $n = 294$, thus

$$\begin{aligned}
SS_B &= \sum_{i=1}^9 (n_i \bar{Y}_i^2) - n \bar{Y}^2 \\
&= (127)(7.347)^2 + (44)(7.368)^2 + (24)(7.418)^2 + (41)(7.487)^2 + (18)(7.563)^2 + \cdots \\
&\quad \cdots + (16)(7.568)^2 + (11)(8.214)^2 + (7)(8.272)^2 + (6)(8.297)^2 - (294)(7.4746)^2 \\
&= 17.45402
\end{aligned}$$

17.45402
!=
22.5208
: maybe
my SS
total calc
is incor-
rect?

(d) If the first 6 species have one common mean and the last 3 have another common mean, we find that

finish 3d

Problem 4:

Consider the Bumpuss data in Chapter 2, compute the power of the two-sided two sample t -test of size 0.05 (i.e., reject the null hypothesis if the absolute value the t -statistic is greater than or equal to 2), under the alternative that $\mu_x - \mu_y = \bar{x} - \bar{y} = 0.01$ and $\sigma = s_p = 0.0214$.

Problem 5:

Show that the two-sided two sample t -test is equivalent to the anova F -test, if the number of groups is two.

For $I = 2$ groups, the F -statistic is given by

$$F\text{-statistic} = \frac{SS_B / [(n-1) - (n-I)]}{SS_W / (n-I)},$$

where n_1 and n_2 are the sizes of samples 1 and 2, respectively, $n = n_1 + n_2$ is the total sample size, SS_B is the “between groups” sum of squared residuals, and SS_W is the “within groups” sum of squared residuals.

Simplifying, we find

$$F\text{-statistic} = \frac{SS_B / (I-1)}{SS_W / (n-I)} = \frac{SS_B / (2-1)}{SS_W / (n-2)} = \frac{SS_B / 1}{SS_W / (n-2)}.$$

If the observations from group 1 are $\sim N(\mu_1, \sigma^2)$ and the observations from group 2 are $\sim N(\mu_2, \sigma^2)$, we know that

$$F\text{-statistic} \sim F_{1, n-2}, \text{ which is equivalent to } t_{n-2}^2.$$

i.e., an F distribution with a numerator degrees of freedom of 1 and a denominator degrees of freedom of $n-2$ is equivalent to the square of a t distribution with $n-2$ degrees of freedom.

Finish
problem
5

Problem 6:

Consider X_1, \dots, X_{10} are i.i.d. $N(0, \sigma^2)$, Y_1, \dots, Y_{10} are i.i.d. $N(\mu, \sigma^2)$ and hypothesis testing:

$$H_0 : \mu = 0$$

$$H_A : \mu \neq 0.$$

Compute the power of a two sided two sample t -test of size 0.05 when $\sigma^2 = 1$ and $\mu = 0.1, 0.5, 1, \text{ and } 2$. Plot the power as a function of μ . Then, increase the sample size in each group to 20 and draw the power function in the same plot as that of the sample size 10.

Problem 7:

Under the setting of the previous problem, show that, under the null hypothesis, the p -value follows the uniform distribution on the interval $[0, 1]$ and perform simulations to confirm it.

Todo list

Should SS total be -9 (-I) instead of -1 ?	1
17.45402 != 22.5208 : maybe my SS total calc is incorrect?	2
finish 3d	2
Finish problem 5	3