

Statistics

Introduction to R for Public Health Researchers

Statistics

Now we are going to cover how to perform a variety of basic statistical tests in R.

- Correlation
- T-tests/Rank-sum tests
- Linear Regression
- Logistic Regression
- Proportion tests
- Chi-squared
- Fisher's Exact Test

Note: We will be glossing over the statistical theory and “formulas” for these tests. There are plenty of resources online for learning more about these tests, as well as dedicated Biostatistics series at the School of Public Health

Correlation

`cor()` performs correlation in R

```
cor(x, y = NULL, use = "everything",  
    method = c("pearson", "kendall", "spearman"))
```

Like other functions, if there are NAs, you get NA as the result. But if you specify use only the complete observations, then it will give you correlation on the non-missing data.

```
library(readr)  
circ = read_csv("http://johnmuschelli.com/intro_to_r/data/Charm_City_Circulator_Rider.  
cor(circ$orangeAverage, circ$purpleAverage, use="complete.obs")
```

```
[1] 0.9195356
```

Correlation with **corrr**

The **corrr** package allows you to do correlations easily:

```
library(corrr); library(dplyr)
circ %>%
  select(orangeAverage, purpleAverage) %>%
  correlate()
```

Correlation method: 'pearson'
Missing treated using: 'pairwise.complete.obs'

```
# A tibble: 2 x 3
  rowname      orangeAverage purpleAverage
  <chr>          <dbl>          <dbl>
1 orangeAverage    NA              0.920
2 purpleAverage  0.920           NA
```

Correlation


`correlate` is usually better with more than 2 columns:

```
avgs = circ %>% select(ends_with("Average"))
cobj = avgs %>% correlate(use = "complete.obs", diagonal = 1)
cobj %>% fashion(decimals = 3)
```

	rowname	orangeAverage	purpleAverage	greenAverage	bannerAverage
1	orangeAverage	1.000	.908	.840	.545
2	purpleAverage	.908	1.000	.867	.521
3	greenAverage	.840	.867	1.000	.453
4	bannerAverage	.545	.521	.453	1.000

Correlation

```
cobj %>% rplot()
```



Correlation

You can also use `cor.test()` to test for whether correlation is significant (ie non-zero). Note that linear regression may be better, especially if you want to regress out other confounders.

```
ct = cor.test(circ$orangeAverage, circ$purpleAverage,  
              use = "complete.obs")  
ct
```

Pearson's product-moment correlation

```
data: circ$orangeAverage and circ$purpleAverage  
t = 73.656, df = 991, p-value < 2.2e-16  
alternative hypothesis: true correlation is not equal to 0  
95 percent confidence interval:  
 0.9093438 0.9286245  
sample estimates:  
      cor  
0.9195356
```

Correlation

For many of these testing result objects, you can extract specific slots/results as numbers, as the `ct` object is just a list.

```
# str(ct)
names(ct)
```

```
[1] "statistic"  "parameter"  "p.value"    "estimate"   "null.value"
[6] "alternative" "method"     "data.name"  "conf.int"
```

```
ct$statistic
```

```
      t
73.65553
```

```
ct$p.value
```

```
[1] 0
```


Broom package

The **broom** package has a **tidy** function that puts most objects into `data.frames` so that they are easily manipulated:

```
library(broom)
tidy_ct = tidy(ct)
tidy_ct
```


```
# A tibble: 1 x 8
```

	estimate	statistic	p.value	parameter	conf.low	conf.high	method	alternative
	<dbl>	<dbl>	<dbl>	<int>	<dbl>	<dbl>	<chr>	<chr>
1	0.920	73.7	0	991	0.909	0.929	Pearson's...	two.sided

Correlation

Note that you can add the correlation to a plot, via the `annotate`

```
library(ggplot2)
txt = paste0("r=", signif(ct$estimate,3))
q = qplot(data = circ, x = orangeAverage, y = purpleAverage)
q + annotate("text", x = 4000, y = 8000, label = txt, size = 5)
```



T-tests

The T-test is performed using the `t.test()` function, which essentially tests for the difference in means of a variable between two groups.

In this syntax, x and y are the column of data for each group.

```
tt = t.test(circ$orangeAverage, circ$purpleAverage)
tt
```

Welch Two Sample t-test

```
data: circ$orangeAverage and circ$purpleAverage
t = -17.076, df = 1984, p-value < 2.2e-16
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 -1096.7602 -870.7867
sample estimates:
mean of x mean of y
 3033.161  4016.935
```

T-tests

Using `t.test` treats the data as independent. Realistically, this data should be treated as a paired t-test. The `paired = TRUE` argument to do a paired test

```
t.test(circ$orangeAverage, circ$purpleAverage, paired = TRUE)
```

Paired t-test

data: circ\$orangeAverage and circ\$purpleAverage

t = -42.075, df = 992, p-value < 2.2e-16

alternative hypothesis: true difference in means is not equal to 0

95 percent confidence interval:

-799.783 -728.505

sample estimates:

mean of the differences

-764.144

T-tests

`t.test` saves a lot of information: the difference in means `estimate`, confidence interval for the difference `conf.int`, the p-value `p.value`, etc.

```
names(tt)
```

```
[1] "statistic"  "parameter"  "p.value"    "conf.int"   "estimate"  
[6] "null.value" "stderr"     "alternative" "method"     "data.name"
```

T-tests

```
tidy(tt)
```

```
# A tibble: 1 x 10
```

	estimate	estimate1	estimate2	statistic	p.value	parameter	conf.low	conf.high
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	-984.	3033.	4017.	-17.1	4.20e-61	1984.	-1097.	-871.

```
# ... with 2 more variables: method <chr>, alternative <chr>
```

T-tests

You can also use the 'formula' notation. In this syntax, it is $y \sim x$, where x is a factor with 2 levels or a binary variable and y is a vector of the same length.

```
library(tidyr)
long = circ %>%
  select(date, orangeAverage, purpleAverage) %>%
  gather(key = line, value = avg, -date)
tt = t.test(avg ~ line, data = long)
tidy(tt)
```

A tibble: 1 x 10

	estimate	estimate1	estimate2	statistic	p.value	parameter	conf.low	conf.high
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	-984.	3033.	4017.	-17.1	4.20e-61	1984.	-1097.	-871.

... with 2 more variables: method <chr>, alternative <chr>

Wilcoxon Rank-Sum Tests

Nonparametric analog to t-test (testing medians):

```
tidy(wilcox.test(avg ~ line, data = long))
```

```
# A tibble: 1 x 4
```

	statistic	p.value	method	alternative
	<dbl>	<dbl>	<chr>	<chr>
1	336714.	4.56e-58	Wilcoxon rank sum test with continuity correct...	two.sided

Lab Part 1

[Website](#)

Analysis of Variance

The `aov` function exists for ANOVA, but we'd recommend `lm` (Linear models):

```
long3 = circ %>%  
  select(date, orangeAverage, purpleAverage, bannerAverage) %>%  
  gather(key = line, value = avg, -date)  
anova_res = aov(avg ~ line, data = long3)  
anova_res
```

Call:

```
aov(formula = avg ~ line, data = long3)
```

Terms:

	line	Residuals
Sum of Squares	2214242251	3724379811
Deg. of Freedom	2	2396

Residual standard error: 1246.762

Estimated effects may be unbalanced

1039 observations deleted due to missingness

Linear Regression

Now we will briefly cover linear regression. I will use a little notation here so some of the commands are easier to put in the proper context.

$$y_i = \alpha + \beta x_i + \varepsilon_i$$

where:

- y_i is the outcome for person i
- α is the intercept
- β is the slope
- x_i is the predictor for person i
- ε_i is the residual variation for person i

Linear Regression

The R version of the regression model is:

$y \sim x$

where:

- y is your outcome
- x is/are your predictor(s)

Linear Regression

For a linear regression, when the predictor is binary this is the same as a t-test:

```
fit = lm(avg ~ line, data = long)
fit
```

```
Call:
lm(formula = avg ~ line, data = long)
```

```
Coefficients:
      (Intercept)  linepurpleAverage
          3033.2             983.8
```

'(Intercept)' is α

'linepurpleAverage' is β

Linear Regression

```
fit = lm(avg ~ line, data = long3)
fit
```

```
Call:
lm(formula = avg ~ line, data = long3)
```

```
Coefficients:
      (Intercept) lineorangeAverage linepurpleAverage
            827.3             2205.9             3189.7
```

Linear Regression

The `summary` command gets all the additional information (p-values, t-statistics, r-square) that you usually want from a regression.

```
sfit = summary(fit)
print(sfit)
```

```
Call:
lm(formula = avg ~ line, data = long3)
```

Residuals:

Min	1Q	Median	3Q	Max
-4016.9	-1008.4	5.6	973.1	4072.6

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	827.27	75.88	10.90	<2e-16	***
lineorangeAverage	2205.89	84.41	26.13	<2e-16	***
linepurpleAverage	3189.67	85.57	37.27	<2e-16	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1247 on 2396 degrees of freedom
(1039 observations deleted due to missingness)

Multiple R-squared: 0.3729, Adjusted R-squared: 0.3723

F-statistic: 712.2 on 2 and 2396 DF, p-value: < 2.2e-16

Linear Regression

We can **tidy** linear models as well and it gives us all of this in a tibble:

```
tidy(fit)
```

```
# A tibble: 3 x 5
```

	term <chr>	estimate <dbl>	std.error <dbl>	statistic <dbl>	p.value <dbl>
1	(Intercept)	827.	75.9	10.9	4.76e- 27
2	lineorangeAverage	2206.	84.4	26.1	1.16e-132
3	linepurpleAverage	3190.	85.6	37.3	2.98e-240

Linear Regression

The `confint` argument allows for confidence intervals

```
tidy(fit, confint = TRUE)
```

```
# A tibble: 3 x 5
```

	term <chr>	estimate <dbl>	std.error <dbl>	statistic <dbl>	p.value <dbl>
1	(Intercept)	827.	75.9	10.9	4.76e- 27
2	lineorangeAverage	2206.	84.4	26.1	1.16e-132
3	linepurpleAverage	3190.	85.6	37.3	2.98e-240

Using Cars Data

```
http_data_dir = "http://johnmuschelli.com/intro_to_r/data/"
cars = read_csv(
  paste0(http_data_dir, "kaggleCarAuction.csv"),
  col_types = cols(VehBCost = col_double()))
head(cars)
```

```
# A tibble: 6 x 34
```

```
  RefId IsBadBuy PurchDate Auction VehYear VehicleAge Make Model Trim SubModel
  <dbl>   <dbl>   <chr>    <chr>    <dbl>    <dbl> <chr> <chr> <chr> <chr>
1     1     0 12/7/2009 ADESA      2006         3 MAZDA MAZD... i    4D SEDA...
2     2     0 12/7/2009 ADESA      2004         5 DODGE 1500... ST   QUAD CA...
3     3     0 12/7/2009 ADESA      2005         4 DODGE STRA... SXT   4D SEDA...
4     4     0 12/7/2009 ADESA      2004         5 DODGE NEON   SXT   4D SEDAN
5     5     0 12/7/2009 ADESA      2005         4 FORD  FOCUS  ZX3   2D COUP...
6     6     0 12/7/2009 ADESA      2004         5 MITS... GALA... ES    4D SEDA...
# ... with 24 more variables: Color <chr>, Transmission <chr>, WheelTypeID <chr>,
# WheelType <chr>, VehOdo <dbl>, Nationality <chr>, Size <chr>,
# TopThreeAmericanName <chr>, MMRAcquisitionAuctionAveragePrice <chr>,
# MMRAcquisitionAuctionCleanPrice <chr>,
# MMRAcquisitionRetailAveragePrice <chr>,
# MMRAcquisitonRetailCleanPrice <chr>, MMRCurrentAuctionAveragePrice <chr>,
# MMRCurrentAuctionCleanPrice <chr>, MMRCurrentRetailAveragePrice <chr>,
# MMRCurrentRetailCleanPrice <chr>, PRIMEUNIT <chr>, AUCGUART <chr>,
# BYRNO <dbl>, VNZIP1 <dbl>, VNST <chr>, VehBCost <dbl>, IsOnlineSale <dbl>,
# WarrantyCost <dbl>
```

Linear Regression

We'll look at vehicle odometer value by vehicle age:

```
fit = lm(VehOdo~VehicleAge, data = cars)
print(fit)
```

Call:

```
lm(formula = VehOdo ~ VehicleAge, data = cars)
```

Coefficients:

(Intercept)	VehicleAge
60127	2723

Linear Regression

Note that you can have more than 1 predictor in regression models. The interpretation for each slope is change in the predictor corresponding to a one-unit change in the outcome, holding all other predictors constant.

```
fit2 = lm(VehOdo ~ IsBadBuy + VehicleAge, data = cars)
summary(fit2)
```

Call:

```
lm(formula = VehOdo ~ IsBadBuy + VehicleAge, data = cars)
```

Residuals:

Min	1Q	Median	3Q	Max
-70856	-9490	1390	10311	41193

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	60141.77	134.75	446.33	<2e-16	***
IsBadBuy	1329.00	157.84	8.42	<2e-16	***
VehicleAge	2680.33	30.27	88.53	<2e-16	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 13810 on 72980 degrees of freedom

Multiple R-squared: 0.1031, Adjusted R-squared: 0.1031

F-statistic: 4196 on 2 and 72980 DF, p-value: < 2.2e-16

Linear Regression: Interactions

The * does interactions:

```
fit3 = lm(VehOdo ~ IsBadBuy * VehicleAge, data = cars)
summary(fit3)
```

Call:

```
lm(formula = VehOdo ~ IsBadBuy * VehicleAge, data = cars)
```

Residuals:

Min	1Q	Median	3Q	Max
-70857	-9490	1391	10311	41193

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	60139.70	143.23	419.872	< 2e-16	***
IsBadBuy	1347.28	456.23	2.953	0.00315	**
VehicleAge	2680.84	32.54	82.380	< 2e-16	***
IsBadBuy:VehicleAge	-3.79	88.74	-0.043	0.96594	

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 13810 on 72979 degrees of freedom

Multiple R-squared: 0.1031, Adjusted R-squared: 0.1031

F-statistic: 2798 on 3 and 72979 DF, p-value: < 2.2e-16

Linear Regression: Interactions

You can take out main effects with minus

```
fit4 = lm(VehOdo ~ IsBadBuy * VehicleAge -IsBadBuy , data = cars)
summary(fit4)
```

Call:

```
lm(formula = VehOdo ~ IsBadBuy * VehicleAge - IsBadBuy, data = cars)
```

Residuals:

Min	1Q	Median	3Q	Max
-70822	-9493	1389	10311	41172

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	60272.49	136.00	443.184	< 2e-16	***
VehicleAge	2652.94	31.14	85.186	< 2e-16	***
IsBadBuy:VehicleAge	242.08	30.70	7.885	3.19e-15	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 13810 on 72980 degrees of freedom

Multiple R-squared: 0.103, Adjusted R-squared: 0.103

F-statistic: 4192 on 2 and 72980 DF, p-value: < 2.2e-16

Linear Regression

Factors get special treatment in regression models - lowest level of the factor is the comparison group, and all other factors are relative to its values.

```
fit3 = lm(VehOdo ~ factor(TopThreeAmericanName), data = cars)
summary(fit3)
```

Call:

```
lm(formula = VehOdo ~ factor(TopThreeAmericanName), data = cars)
```

Residuals:

Min	1Q	Median	3Q	Max
-71947	-9634	1532	10472	45936

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	68248.48	92.98	733.984	< 2e-16	***
factor(TopThreeAmericanName)FORD	8523.49	158.35	53.828	< 2e-16	***
factor(TopThreeAmericanName)GM	4952.18	128.99	38.393	< 2e-16	***
factor(TopThreeAmericanName)NULL	-2004.68	6361.60	-0.315	0.752670	
factor(TopThreeAmericanName)OTHER	584.87	159.92	3.657	0.000255	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 14220 on 72978 degrees of freedom

Multiple R-squared: 0.04822, Adjusted R-squared: 0.04817

F-statistic: 924.3 on 4 and 72978 DF, p-value: < 2.2e-16

Logistic Regression and GLMs

Generalized Linear Models (GLMs) allow for fitting regressions for non-continuous/normal outcomes. The `glm` has similar syntax to the `lm` command. Logistic regression is one example.

```
glmfit = glm(IsBadBuy ~ VehOdo + VehicleAge, data=cars, family = binomial())
summary(glmfit)
```

Call:

```
glm(formula = IsBadBuy ~ VehOdo + VehicleAge, family = binomial(),
     data = cars)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-0.9943	-0.5481	-0.4534	-0.3783	2.6318

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	-3.778e+00	6.381e-02	-59.211	<2e-16	***
VehOdo	8.341e-06	8.526e-07	9.783	<2e-16	***
VehicleAge	2.681e-01	6.772e-03	39.589	<2e-16	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 54421 on 72982 degrees of freedom
Residual deviance: 52346 on 72980 degrees of freedom
AIC: 52352

Number of Fisher Scoring iterations: 5

Tidying GLMs

```
tidy(glmfit, conf.int = TRUE)
```

```
# A tibble: 3 x 7
```

	term <chr>	estimate <dbl>	std.error <dbl>	statistic <dbl>	p.value <dbl>	conf.low <dbl>	conf.high <dbl>
1	(Intercept)	-3.78	0.0638	-59.2	0.	-3.90	-3.65
2	VehOdo	0.00000834	0.000000853	9.78	1.33e-22	0.00000667	0.0000100
3	VehicleAge	0.268	0.00677	39.6	0.	0.255	0.281

Tidying GLMs

```
tidy(glmfit, conf.int = TRUE, exponentiate = TRUE)
```

```
# A tibble: 3 x 7
```

	term <chr>	estimate <dbl>	std.error <dbl>	statistic <dbl>	p.value <dbl>	conf.low <dbl>	conf.high <dbl>
1	(Intercept)	0.0229	0.0638	-59.2	0.	0.0202	0.0259
2	VehOdo	1.00	0.000000853	9.78	1.33e-22	1.00	1.00
3	VehicleAge	1.31	0.00677	39.6	0.	1.29	1.32

Logistic Regression

Note the coefficients are on the original scale, we must exponentiate them for odds ratios:

```
exp(coef(glmfit))
```

(Intercept)	VehOdo	VehicleAge
0.02286316	1.00000834	1.30748911

Chi-squared tests

`chisq.test()` performs chi-squared contingency table tests and goodness-of-fit tests.

```
chisq.test(x, y = NULL, correct = TRUE,  
           p = rep(1/length(x), length(x)), rescale.p = FALSE,  
           simulate.p.value = FALSE, B = 2000)
```

```
tab = table(cars$IsBadBuy, cars$IsOnlineSale)  
tab
```

	0	1
0	62375	1632
1	8763	213

Chi-squared tests

You can also pass in a table object (such as `tab` here)

```
cq = chisq.test(tab)
cq
```

Pearson's Chi-squared test with Yates' continuity correction

```
data:  tab
X-squared = 0.92735, df = 1, p-value = 0.3356
```

```
names(cq)
```

```
[1] "statistic" "parameter" "p.value"    "method"    "data.name" "observed"
[7] "expected"  "residuals" "stdres"
```

```
cq$p.value
```

```
[1] 0.3355516
```

Chi-squared tests

Note that does the same test as `prop.test`, for a 2x2 table (`prop.test` not relevant for greater than 2x2).

```
chisq.test(tab)
```

Pearson's Chi-squared test with Yates' continuity correction

```
data: tab  
X-squared = 0.92735, df = 1, p-value = 0.3356
```

```
prop.test(tab)
```

2-sample test for equality of proportions with continuity correction

```
data: tab  
X-squared = 0.92735, df = 1, p-value = 0.3356  
alternative hypothesis: two.sided  
95 percent confidence interval:  
 -0.005208049 0.001673519  
sample estimates:  
  prop 1    prop 2  
0.9745028 0.9762701
```

Fisher's Exact test

`fisher.test()` performs contingency table test using the hypergeometric distribution (used for small sample sizes).

```
fisher.test(x, y = NULL, workspace = 200000, hybrid = FALSE,  
            control = list(), or = 1, alternative = "two.sided",  
            conf.int = TRUE, conf.level = 0.95,  
            simulate.p.value = FALSE, B = 2000)
```

```
fisher.test(tab)
```


Fisher's Exact Test for Count Data

```
data:  tab  
p-value = 0.3324  
alternative hypothesis: true odds ratio is not equal to 1  
95 percent confidence interval:  
 0.8001727 1.0742114  
sample estimates:  
odds ratio  
 0.9289923
```

Sampling

Also, if you want to only plot a subset of the data (for speed/time or overplotting)

```
samp.cars = sample_n(cars, size = 10000)
samp.cars = sample_frac(cars, size = 0.2)
ggplot(aes(x = VehBCost, y = VehOdo),
       data = samp.cars) + geom_point()
```



Lab Part 2

[Website](#)