

# Matrix

see [math notation](#)

**definition** *formally in my [math notation](#)* a [matrix](#) in  $\mathbb{R}^{m,n}$  is a [set theoryetical function](#) with domain at least  $\langle x,y \rangle \rightarrow \mathbb{N}x \wedge \mathbb{N}y \wedge 0 \leq x < m \wedge 0 \leq y < n$  that takes an [ordered pair](#) as an index and returns the element at that index

**notation**

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

## Multiplication by a Scalar

see [matrix vector space](#), [vector space](#)

**definition**

$$(kA)^{i,j} = kA^{i,j} \dashv \mathbb{N}i \wedge \mathbb{N}j \wedge \mathbb{R}k \wedge \mathbb{M}A$$

**properties**

*commutativity with [scalars](#)*  $kA = Ak$

## Matrix Addition

see [matrix vector space](#), [vector space](#)

**definition**

$$(A : B)^{i,j} = A^{i,j} : B^{i,j} \dashv \mathbb{N}i \wedge \mathbb{N}j \wedge \mathbb{M}^{m,n}A \wedge \mathbb{M}^{m,n}B$$

## Matrix Multiplication

see [dot product](#), [vector in rn](#)

**definition**

$AB \neq \emptyset \equiv \mathbb{M}^{m,n}A \wedge \mathbb{M}^{n,p}B \wedge \mathbb{M}^{m,p}AB$  ( $AB$  is defined if the number of columns in  $A$  is equal to the number of rows in  $B$ . their product will be an  $m$  by  $p$  [matrix](#))

$(AB)^{i,j} = A^{i, \cdot} \cdot B^{\cdot, j} \dashv \mathbb{N}i \wedge \mathbb{N}j$ , see dot product

intuitively, matrix multiplication is the dot product of **every row** of the first matrix by **every column** of the second matrix

## notation

$$AA = A^2 = [A]^2 \dashv \mathbb{M}A$$

$$\text{and therefore } AA \dots A = [A]^n \wedge \mathbb{N}n \dashv \mathbb{M}A$$

## properties

*not commutative*  $AB = BA \not\wedge \mathbb{M}A \wedge \mathbb{M}B$  or  $AB \neq BA \wedge \mathbb{M}A \wedge \mathbb{M}B$

$AB = 0 \not\Leftarrow A = 0 \vee B = 0$  (it can happen that  $AB = 0$ , but  $A \neq 0$  and  $B \neq 0$ ) ( $AB$  being equal to 0 does not imply that  $A = 0$  or that  $B = 0$ )

$AC = BC \wedge C \neq 0 \not\Leftarrow A = B$  ( $AC = BC$  and  $C \neq 0$  does not imply that  $A = B$ )

*associative*  $(AB)C = A(BC)$

*distributive*  $A(B : C) = AB : AC$

*distributive*  $(B : C)A = BA : CA$

*associative with scalars*  $k(AB) = (kA)B = A(kB)$

## applications

matrix multiplication can be used to represent a linear system of linear equations:

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

matrix multiplication can be used to represent any linear transformation

# Identity Matrix

## definition

$$(I^{a,b} = 1 \wedge a = b) \vee (I^{a,b} = 0 \wedge a \neq b) \dashv \mathbb{N}a \wedge \mathbb{N}b \wedge \mathbb{M}^{n,n}I$$

## examples

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

properties

$$AI = A \wedge IA = A \dashv \mathbb{M}A$$

## Zero Matrix

see [matrix vector space](#), [vector space](#)

definition

$$O^{a,b} = 0 \dashv \mathbb{N}a \wedge \mathbb{N}b \wedge \mathbb{M}^{n,m}O$$

examples

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

properties

$$A_{m,n}O_{n,p} = O_{m,p} \dashv \mathbb{M}^{n,p}O_{n,p} \wedge \mathbb{M}^{m,p}O_{m,p} \wedge \mathbb{M}^{m,n}A_{m,n}$$

$$O_{q,m}A_{m,n} = O_{q,n} \dashv \mathbb{M}^{q,m}O_{q,m} \wedge \mathbb{M}^{q,n}O_{q,n} \wedge \mathbb{M}^{m,n}A_{m,n}$$

## Rank of a Matrix

*the number of pivots in any REF of the matrix*

notation

*rank*  $A$ , where

- $A$  is the matrix to find the rank of

## Matrix Vector Spaces

*Null Space (Nullspace, Kernel), Column Space, Row Space*

## notations

*kernel*  $\text{Ker } A \equiv \text{Null } A$

*column space*  $\text{Col } A$

*row space*  $\text{Row } A$

## definitions

*kernel*  $(\text{Ker } A) \ x \equiv (\text{Null } A) \ x \equiv Ax = O \wedge \mathbb{M}^{n,n} A \wedge \mathbb{M}^{n,1} x$

*column space*  $\text{Col } A = \text{span } A^n \dashv \mathbb{N}n$

*row space*  $\text{Row } A = \text{span } A^n, \dashv \mathbb{N}n$

**procedure** *computing the kernel of a matrix use row reduction*

## theorems

the Null Space, Row Space and Column Space of a matrix are always vector spaces

number of free variables in  $A$  : number of pivots in  $A$  = number of columns in  $A$

$\dim \text{Null } A$  = number of free variables in  $A$

$\text{rank } A$  = number of pivots in  $A$

the nonzero rows in any REF of a matrix  $A$  forms a basis for  $\text{Row } A$ . therefore,  
 $\dim \text{Row } A = \text{rank } A$  (see rank of a matrix)

if  $A$  and  $B$  are row-equivalent, then  $\text{Row } A = \text{Row } B$ , see linear system

the spanning set of  $\text{Null } A$  obtained from applying row reduction on the system  $Ax = O$  is a basis for  $\text{Null } A$

$\text{Row } A$  does not change when applying elementary operations on the rows of  $A$ , see linear system

## properties

$\text{Col } A = \text{Row } A^T \wedge \text{Row } A = \text{Col } A^T \dashv \mathbb{M}A$ , see transpose matrix

## applications

row spaces can be used to find a basis for a spanning set of vectors through row reduction

the basis for the row space of a matrix can be found by applying row reduction and spanning the **row-reduced columns** in the REF form of the matrix

the basis for the column space of a matrix can be found by applying row reduction and spanning the **original columns** that became pivots in the REF form of the matrix

the same can be said for *Col A*

**example** *transforming a vector space into the null space of a certain matrix*

let  $W = \text{span} \langle (1, 0, 0, 1), (1, 1, 1, 0), (2, 1, \cdot 1, 1) \rangle$

after solving the linear system, we get  $W(x, y, z, w) \equiv \cdot x : y : w = 0$ . therefore,  $W$  is the null space of  $A = \begin{bmatrix} \cdot & 1 & 0 & 1 \end{bmatrix}$

## Transpose Matrix

*the transpose of a matrix*

*flips a matrix around its diagonal*

**note** the *diagonal* of a square matrix goes from its top left element to its bottom right element (triplicate)

### definition

$$(A^\top)^{i,j} = (A)^{j,i} \dashv \mathbb{N}i \wedge \mathbb{N}j \wedge \mathbb{M}A$$

### properties

$$(A^\top)^\top = A \dashv \mathbb{M}A$$

$$(AB)^\top = B^\top A^\top \dashv \mathbb{M}A \wedge \mathbb{M}B$$

### representation

A

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$$

—

[https://upload.wikimedia.org/wikipedia/commons/thumb/e/e4/Matrix\\_transpose.gif/200px-Matrix\\_transpose.gif](https://upload.wikimedia.org/wikipedia/commons/thumb/e/e4/Matrix_transpose.gif/200px-Matrix_transpose.gif)

## Matrix Inverse

the inverse of a matrix

### definition

$AA^{-1} = A^{-1}A = I \vdash \mathbb{M}A$ , where

- $A$  is a square matrix
- $A^{-1}$  is the *inverse matrix* of  $A$

## Invertability

**definition** an *invertible matrix* has a corresponding inverse matrix

see theorems below for invertability criteria

### properties

let  $A$  and  $C$  be invertible matrixes, let  $\mathbb{Z}p$  and let  $\mathbb{R}k \wedge k \neq 0$ . then,

$$AA^{-1} = A^{-1}A = I$$

$$(A^{-1})^{-1} = A$$

$$(A^p)^{-1} = (A^{-1})^p$$

$$(kA)^{-1} = -k \mid A^{-1} \text{ (restriction might not be necessary, see improved expression evaluation)}$$

$$(AC)^{-} = C^{-}A^{-}$$

**note** in the equation above, the order of the matrixes has changed. this is significant as matrix multiplication is not commutative

if  $AC$  is invertible, then  $A$  is invertible and  $C$  is invertible

**procedure** *computing the inverse of a matrix*

let  $\mathbb{M}^{n,n}A$

solve the system  $AA^{-} = I$  by extending the matrix with the identity matrix and solve the linear system up to RREF using row reduction.  $[A \mid I] \sim \dots [I \mid A^{-}]$

**procedure** *computing the inverse of a 2 by 2 matrix*

see determinant

$$\text{let } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$A$  is invertible if and only if  $|A| \neq 0$

$$A^{-} = -|A| \mid \begin{bmatrix} d & \cdot b \\ \cdot c & a \end{bmatrix}$$

**applications** *using a matrix inverse to solve a linear system*

$$\text{let } A = \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix}$$

then, calculate  $B$  such that  $B \equiv A^{-}$

this can be used to solve a linear system such as:

$$Ax = \begin{bmatrix} \cdot 1 \\ 1 \end{bmatrix}$$

$$BAx = B \begin{bmatrix} \cdot 1 \\ 1 \end{bmatrix}$$

$$Ix = x = B \begin{bmatrix} \cdot 1 \\ 1 \end{bmatrix}$$

## Triangular Matrix

**definition** a matrix is said to be *triangular* if every entry below its diagonal **or** above its diagonal is 0

**note** the *diagonal* of a square matrix goes from its top left element to its bottom right element (triplicate)

## Diagonal Matrix

**definition** a matrix is said to be *diagonal* if every entry below its diagonal **and** above its diagonal is 0

**note** the *diagonal* of a square matrix goes from its top left element to its bottom right element (triplicate)

let  $D$  be a diagonal matrix

**applications**

$[D]x$  can be calculated by raising every entry of  $D$  to the power  $x$

## Diagonalizable Matrix

see eigenvector

**definition** an  $n$  by  $n$  matrix  $A$  is said to be *diagonalizable over the reals* if there exists a basis of  $\mathbb{R}^n$  consisting entirely of eigenvectors of  $A$

a matrix is *diagonalizable* if and only if the geometric multiplicity of an eigenvalue is equal to the algebraic multiplicity of said eigenvalue, for every eigenvalue of the matrix (see eigenvector and eigenvalue)

**note** a matrix may also be diagonalizable over other number fields such as the set of complex numbers  $\mathbb{C}$

**note** some matrixes do not have "enough" real eigenvalues or "enough" eigenvectors to be diagonalizable

**example** the matrix  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$  is diagonalizable over the reals as  $\langle\langle (1,1), (1,-1) \rangle\rangle$  is a basis of  $\mathbb{R}^2$  consisting entirely of eigenvectors of  $A$



**example** the matrix  $A = \begin{bmatrix} 1 & 1 \\ .1 & 1 \end{bmatrix}$  is not diagonalizable over the reals as it does not have any real eigenvalues

**example** the matrix  $A = \begin{bmatrix} 3 & 1 \\ 0 & 3 \end{bmatrix}$  is not diagonalizable over the reals as it only has one eigenvalue, and therefore only one set of linearly dependent eigenvectors (see eigenvector and eigenvalue)

**example** the matrix  $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  is diagonalizable over the reals as, even though  $A$  has a single eigenvalue  $\lambda = 1$ , its eigenspace spans  $\mathbb{R}^2$ . this is the case for both  $A = I \wedge \lambda = 1$  and  $A = O \wedge \lambda = 0$

**proof** let  $A = I \wedge \lambda = 1 \wedge E_1 = x$ . we then have  $O = A \cdot \lambda I \mid x = I \cdot 1I \mid E_1 = O \mid E_1$ . therefore,  $E_1 \equiv \mathbb{R}^2$ . see eigenvector and eigenvalue

**example** let  $\mathbb{M}^{n,n}A \wedge \mathbb{N}n$  and suppose  $A$  has  $n$  distinct eigenvalues. deduce that  $A$  is diagonalizable over the reals

**proof**  $A$  has at most  $n$  eigenvalues  $\rightarrow$  the algebraic multiplicity of every eigenvalue of  $A$  is 1 as they are all distinct and must be greater than 1  $\rightarrow$  the geometric multiplicity of every eigenvalue of  $A$  is 1 as it must be greater than 1 and less than its algebraic multiplicity  $\rightarrow$  all algebraic multiplicities and geometric multiplicities are equal  $\rightarrow A$  is diagonalizable. see eigenvector and eigenvalue

## eigenvector and eigenvalues

### theorems

see linear system

#### theorem

let  $\mathbb{M}^{m,n}A$  (see matrix). the following logic statements are equivalent:

- every variable is a leading variable
- there is a leading variable in every column of the RREF of  $A$
- the system  $Ax = O$  has a unique solution

- the columns of  $A$  are linearly independent
- $\text{Ker } A = \langle \langle 0 \rangle \rangle$
- $\dim \text{Ker } A = 0$
- $\text{rank } A = n$

see linear system theorem proof

### **theorem**

let  $\mathbb{M}^{n,n}A$  (see matrix). the following logic statements are equivalent:

**note** all logic statements below are valid for both  $A$  and  $A^T$ , see transpose matrix

- $\text{rank } A = n$
- every linear system of the form  $Ax = b$  has a unique solution
- the RREF of  $A$  is the identity matrix
- $\text{Ker } A = \langle \langle 0 \rangle \rangle$
- $\text{Col } A = \mathbb{R}^n$
- $\text{Row } A = \mathbb{R}^n$
- the columns of  $A$  are linearly independent
- the rows of  $A$  are linearly independent
- the columns of  $A$  form a basis for  $\mathbb{R}^n$
- the rows of  $A$  form a basis for  $\mathbb{R}^n$
- $A$  is an invertible matrix
- $\det A \neq 0$