





clams_analysis



Quynh Tran



November 23, 2015


This is an R Markdown document. Markdown is a simple formatting syntax for authoring HTML, PDF, and MS Word documents. For more details on using R Markdown see <http://rmarkdown.rstudio.com>.

Body Weight Analysis









	pval	padj
Weight	0.00000000	0.00000000
Fat	0.00000000	0.00000000
Lean	0.00000000	0.00000000
Percent.Fat	0.00000000	0.00000000

Table 1: Oneway ANOVA with Welch correction for effects of Fasting and HFD on Weight, Total Fat Mass, and Lean Mass; Kruskal Wallis test on Percent Fat Mass. P-values are adjusted by the method of Benjamini and Hochberg

	diff	lwr	upr	p adj
After_fasting-Pre_fasting	-1.18518519	-2.30496396	-0.06540641	0.03527128
After_HFD-Pre_fasting	15.97407407	14.85429530	17.09385285	0.00000000
After_HFD-After_fasting	17.15925926	16.03948049	18.27903803	0.00000000

Table 2: Tukey HSD for pairwise comparisons among Fasting stages on Body Weight

	diff	lwr	upr	p adj
After_fasting-Pre_fasting	-0.35055556	-1.44144958	0.74033847	0.72786160
After_HFD-Pre_fasting	12.27166667	11.18077264	13.36256069	0.00000000
After_HFD-After_fasting	12.62222222	11.53132820	13.71311625	0.00000000

Table 3: Tukey HSD for pairwise comparisons among Fasting stages on Fat Mass

	diff	lwr	upr	p adj
After_fasting-Pre_fasting	-0.98388889	-1.62497062	-0.34280716	0.00110244
After_HFD-Pre_fasting	3.67111111	3.03002938	4.31219284	0.00000000
After_HFD-After_fasting	4.65500000	4.01391827	5.29608173	0.00000000

Table 4: Tukey HSD for pairwise comparisons among Fasting stages on Lean Mass

	Pre_fasting	After_fasting
After_fasting	0.06877140	
After_HFD	0.00000000	0.00000000

Table 5: Pairwise comparisons using Nemenyi-test with Chi-squared approximation for independent samples