

Tissue Weights for Muscle Tsc1 Knockout Mice on HFD

Dave Bridges and Erin Stephenson

2019-03-01

Contents

1 Purpose	1
2 Experimental Details	1
3 Analysis	1
3.1 Fat Pad Weights	2
3.2 Muscle Weights	4
4 Session Information	4

1 Purpose

To determine tissue weights at sacrifice for fat pads and muscle tissues

2 Experimental Details

At sacrifice, after a 16h fast data were entered and collected in the raw data sheet

These data can be found in `/Users/davebrid/Documents/GitHub/TissueSpecificTscKnockouts/Mouse Data/Muscle Tsc1 Knockout/HFD/Sacrifice Analysis` in a file named **HFD Sacrifice Data.xlsx**.


This script was most recently updated on **Fri Mar 1 14:51:43 2019**.


3 Analysis

Sex	GonadalWAT_mean.na	InguinalWAT_mean.na	Quadriceps_mean.na	TricepsSurae_mean.na	Heart_mean.na
Male	0.792	0.574	0.224	0.172	0.101
Male	0.181	0.116	0.192	0.156	0.076
Female	0.492	0.316	0.166	0.132	0.081
Female	0.101	0.076	0.142	0.121	0.076

Sex	GonadalWAT_length	InguinalWAT_length	Quadriceps_length	TricepsSurae_length	Heart_length
Male	25	25	25	25	25
Male	7	7	7	7	7
Female	53	53	53	53	53
Female	8	8	8	8	8

3.1 Fat Pad Weights





For the male mice, the fat pads were reduced in weight:

Table 3: Changes in Gonadal Fat Pad Weights

Sex	Wild-Type	Knockout	Difference	Pct.Difference
Male	0.792	0.181	0.611	77.1
Female	0.492	0.101	0.391	79.5

Table 4: Changes in Inguinal Fat Pad Weights

Sex	Wild-Type	Knockout	Difference	Pct.Difference
Male	0.574	0.116	0.458	79.8
Female	0.316	0.076	0.240	75.9

Table 5: Shapiro-Wilk Tests for each group

Sex	Knockout	InguinalWAT_shapiro.p	GonadalWAT_shapiro.p
Male	FALSE	0.400	0.172
Male	TRUE	0.856	0.319
Female	FALSE	0.000	0.000
Female	TRUE	0.703	0.975

Table 6: Pairwise tests for effects of knockout on Inguinal WAT weights.

Sex	Levene	Mann.Whitney	Welch	Student
Female	0.018	0.001	0	0.007
Male	0.008	0.000	0	0.002

Table 7: Pairwise tests for effects of knockout on Gonadal WAT weights.

Sex	Levene	Mann.Whitney	Welch	Student
Female	0.016	0	0	0.007
Male	0.009	0	0	0.000

For the male mice, normality can be assumed, but not equal variance, so a Welch’s t-test is used, which had a p-value of 9.802×10^{-7} for inguinal WAT and 4.917×10^{-8} for gonadal WAT.

For the female mice, normality cannot be assumed so a Mann-Whitney test is used, which had a p-value of 0.001 for inguinal WAT and 0 for gonadal WAT.

3.2 Muscle Weights

4 Session Information

```
sessionInfo()
```

```
## R version 3.5.0 (2018-04-23)
## Platform: x86_64-apple-darwin15.6.0 (64-bit)
## Running under: macOS 10.14.2
##
## Matrix products: default
## BLAS: /Library/Frameworks/R.framework/Versions/3.5/Resources/lib/libRblas.0.dylib
## LAPACK: /Library/Frameworks/R.framework/Versions/3.5/Resources/lib/libRlapack.dylib
##
## locale:
## [1] en_US.UTF-8/en_US.UTF-8/en_US.UTF-8/C/en_US.UTF-8/en_US.UTF-8
##
## attached base packages:
## [1] stats      graphics  grDevices utils      datasets  methods   base
##
## other attached packages:
## [1] car_3.0-2      carData_3.0-2    forcats_0.3.0
## [4] gridExtra_2.3  ggplot2_3.1.0    RColorBrewer_1.1-2
## [7] bindrcpp_0.2.2 readxl_1.2.0     dplyr_0.7.8
## [10] tidyr_0.8.2    knitr_1.21
##
## loaded via a namespace (and not attached):
## [1] zip_1.0.0      Rcpp_1.0.0      pillar_1.3.1
## [4] compiler_3.5.0 cellranger_1.1.0 highr_0.7
```




Figure 1: Weights of Muscle Depots at Sacrifice

## [7]	plyr_1.8.4	bindr_0.1.1	tools_3.5.0
## [10]	digest_0.6.18	evaluate_0.12	tibble_2.0.0
## [13]	gtable_0.2.0	pkgconfig_2.0.2	rlang_0.3.1
## [16]	openxlsx_4.1.0	cli_1.0.1	curl_3.2
## [19]	yaml_2.2.0	haven_2.0.0	xfun_0.4
## [22]	rio_0.5.16	withr_2.1.2	stringr_1.3.1
## [25]	hms_0.4.2	grid_3.5.0	tidyselect_0.2.5
## [28]	data.table_1.11.8	glue_1.3.0	R6_2.3.0
## [31]	fansi_0.4.0	foreign_0.8-71	rmarkdown_1.11
## [34]	purrr_0.2.5	magrittr_1.5	scales_1.0.0
## [37]	htmltools_0.3.6	abind_1.4-5	assertthat_0.2.0
## [40]	colorspace_1.3-2	labeling_0.3	utf8_1.1.4
## [43]	stringi_1.2.4	lazyeval_0.2.1	munsell_0.5.0
## [46]	crayon_1.3.4		