

Statistical__Inference1

Laura Bennett

June 3, 2017

Statistical Inference for Data Science: Chapter 5 Exercises

1. If I have a random sample from a population, the sample variance is an estimate of what?
 - **The population variance**
2. The distribution of the sample variance of a random sample from a population is centered at what?
 - **The population variance**
3. I keep drawing samples of size n from a population with variance σ^2 and taking their average. I do this thousands of times. If I were to take the variance of the collection of averages, what would it be?
 - **In this case, you are seeking the standard error of the mean, a measure of the variability of the mean of a distribution. Thus, the variance of the collection of averages should be $\frac{\sigma^2}{n}$**
4. You get a random sample of n observations from a population and take their average. You would like to estimate the variability of averages of n observations from this population to better understand how precise of an estimate it is. Do you need to repeatedly collect averages to do this?
 - **No; you may instead divide the variance by n and obtain the standard error of the mean.**
5. A random variable takes the value -4 with probability 0.2 and 1 with probability 0.8. What is the variance of this random variable?
 - **The variance of a random variable may be found as $Var(X) = E(X^2) - E(X)^2$, where $E(X) = \sum x_i p(x)$. In this case, $E(X) = (-4)(0.2) + (1)(0.8) = 0$ and $E(X^2) = (-4)^2(0.2) + (1)^2(0.8) = 4.0$; therefore, $Var(X) = 4.0 - 0 = 4.0$**
6. If \bar{X} and \bar{Y} are comprised of n iid random variables arising from distributions having means μ_x and μ_y respectively and common variance σ^2 , what is the variance $\bar{X} - \bar{Y}$?
 - **According to the central limit theorem, $Var(X \pm Y) = Var(X) + Var(Y)$ IF X and Y are independent. Given that the sample variance from both X and Y is $\frac{\sigma^2}{n}$, $Var(\bar{X} - \bar{Y}) = \frac{2\sigma^2}{n}$**
7. Let X be a random variable have a standard deviation σ . What can be said about the variance of $\frac{X}{\sigma}$?
 - **The variance of a constant times a random variable is given by $Var(aX) = a^2 Var(X)$. Thus, $Var\left(\frac{X}{\sigma}\right) = \frac{\sigma^2}{\sigma^2} = 1$**
8. What would be the variance for a random variable with values [2:5] and accompanying probabilities (0.1, 0.2, 0.3, 0.4)?
 - ```
p<-c(.1,.2,.3,.4)
x<-2:5
variance <- sum(x^2*p)-(sum(x*p))^2
print(variance)
```
- ## [1] 1
9. If you roll ten standard dice, take their average, then repeat this process over and over and construct a histogram, what would be its variance expressed to 3 decimal places?

- The variance of a standard die may be calculated as  $Var(X) = \frac{1}{6} [1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2] - \frac{1}{6^2} [1 + 2 + 3 + 4 + 5 + 6]^2 = 2.92$ . Since we are essentially looking for the variance of the mean, we may calculate it as  $Var(\bar{X}) = \frac{\sigma^2}{n} = \frac{2.92}{10} = 0.292$