






# 13. Refinement of Extended Structures

John S.O. Evans  
BCA School, 2005



## Extended Structures

- "Non Molecular" species
- Inorganic Compounds
  - oxides, chalcogenides, alloys
- Minerals
- Coordination polymers



BCA 2005

john.evans@durham.ac.uk www.durham.ac.uk/jsoe



## Overview of Lecture

- What is an extended structure?
- Potential pitfalls during data collection
- Potential pitfalls during solution/refinement
- Disordered materials
- Limits of Bragg Diffraction
- How to know you've got it right/wrong
- Supporting techniques
- Case histories
- Thanks to Clare, Graham, Matt, Sarah, Martin, Ivana, Thomas Proffen for data/examples!

BCA 2005

john.evans@durham.ac.uk www.durham.ac.uk/jsoe



## Molecular Symmetry

BCA 2005

john.evans@durham.ac.uk www.durham.ac.uk/jsoe



## Extended Symmetry

BCA 2005

john.evans@durham.ac.uk www.durham.ac.uk/jsoe




## Pitfall Avoidance

BCA 2005

john.evans@durham.ac.uk www.durham.ac.uk/jsoe

## Pitfall Avoidance



© ICA 2005

## Data Collection – Crystal Quality, etc

- Coordination polymers and materials grown hydrothermally frequently very insoluble and form as polycrystalline powders – can't recrystallise
- Crystals frequently grown by slow cooling from melt – must be chipped away from other material
- Twinning – wait for later!
- May need to use synchrotron source or powder diffraction methods
- Good absorption corrections can become vital



© ICA 2005

## $\lambda/2$ Contamination

- Most diffractometers use graphite 001 monochromator to select a single wavelength

$$I = 2d_{001} \sin \theta \quad \frac{I}{2d_{001}} = \sin \theta \quad \frac{I}{2d_{001}} = \frac{(I/2)}{2(d_{001}/2)}$$

- "Monochromatic" beam will also contain  $\lambda/2$

© ICA 2005


## $\lambda/2$ Contamination/Solutions

$$F'_{hkl} = F_{hkl} + kF_{2h2k2l}$$

- Each reflection (2h,2k,2l) will contribute to intensity of (h,k,l) due to  $\lambda/2$
- Area detectors may see extra peaks
- All reflection intensities may be effected
- Important when  $F_{hkl}$  small and  $F_{2h2k2l}$  large
- For Mo run tube at 34.9 kV


$$I_{\text{tube}} = hc/eV = 1239 \text{ eV}$$

- Use a Si/Ge (111) monochromator –  $F_{222}$  small



© ICA 2005


## $\lambda/2$ Contamination – $\text{Bi}_{4.5}\text{CaVO}_{10.5}$



50 kV, 40 mA      35 kV, 50 mA

© ICA 2005

## $\lambda/2$ Contamination – $\text{Bi}_{4.5}\text{CaVO}_{10.5}$



50 kV, 40 mA      35 kV, 50 mA

- $F^2_{-6-3-3} = 498.45$ , 13<sup>th</sup> strongest reflection in data set

© ICA 2005

## $\lambda/2$ Contamination Solutions

$$F'_{hkl} = F_{hkl} + kF_{2h2k2l} \quad k \approx 0.001$$

“for routine data sets” ..... “the effect on final refinements was negligible. There was no significant change in the final agreement factors and the changes in the geometrical parameters were all smaller than 1 e.s.d.”

“.....it is quite easy to obtain the wrong unit cell for strongly scattering samples when pure  $\lambda/2$  reflections are included.....”


Kirschbaum et al, *J. Appl. Cryst.* (1997), **30**, 514-6

ICMA 2005     info.xray@diffrao.ac.uk     www.diffrao.ac.uk/info/





## Heavy Atoms

- Heavy metal atoms may dominate diffraction
- $\text{Bi}_2\text{O}_3$  contains Bi (83 electrons) and O (8 electrons)
- $\text{CH}_3$  contains C (6 electrons) and H (1 electron)

- O may have very small contribution to diffraction pattern
- Pseudo symmetry due to heavy atoms dominating




ICMA 2005     info.xray@diffrao.ac.uk     www.diffrao.ac.uk/info/




## Partial Occupancy and Disorder

- Often crucial to the properties of materials







YBa<sub>2</sub>Cu<sub>3</sub>O<sub>7-δ</sub>  
Superconductor for δ<0.6



Remove O  
Swap La/Sr


La<sub>1-x</sub>Sr<sub>x</sub>MnO<sub>3</sub>  
Antiferromagnetic Insulator to Ferromagnetic Metal

ICFA 2005    john.arnold@durham.ac.uk    www.durham.ac.uk/John.Arnold









## Caveats/Solutions

- Beware e.g. 2:1 disorder of H:D in neutron experiments (b=-0.373/+0.667)
- Don't try and work on MO<sub>x</sub>F<sub>y</sub> systems!
- La<sub>1-x</sub>Ca<sub>x</sub>MnO<sub>3</sub>: Beware e.g. La<sub>0.5</sub>Ca<sub>0.5</sub>MnO<sub>3</sub> where cation ordering can cause complications
- Beware incommensurate superstructures



- Combine X-ray and neutron data
- Consider anomalous scattering
- Consider other techniques – e.g. solid state NMR
- Think

ICFA 2005    john.arnold@durham.ac.uk    www.durham.ac.uk/John.Arnold



## Order and Disorder

cations anions

© ICA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/JohnEvans](http://www.durham.ac.uk/JohnEvans)

## Structure Validation

- Molecular structures relatively straightforward
- C-C 1.54 ?
- C=C 1.34 ?
- CSD for more complex examples
- Metal coordination less predictable, depends on oxidation state, Jahn Teller/Lone pair distortions, etc
- e.g. Bi(III) vs Bi(V)

© ICA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/JohnEvans](http://www.durham.ac.uk/JohnEvans)

## Bond Valence Sums

$$v_{ij} = \exp\left[\frac{(R_{ij} - d_{ij})}{0.37}\right]$$

Bond valence parameter From tables.  
e.g.  $R_{\text{Bi-O}} = 2.07$   
 $R_{\text{Re-O}} = 1.79$

$$V = \sum_j v_{ij}$$

Total Valence (V) for each cation/anion in structure should match expected values: Bi 3, Mg 2, V 5, O 2, etc

e.g. Bi(III)  
4\*O at 2.25 ? ?  $4*v_i = 0.649 = 2.592$   
3\*O at 2.80 ? ?  $3*v_i = 0.147 = 0.441$   
**3.033**

$v_{13} = \exp[(2.07 - 2.80)/0.37] = 0.147$   
 $d_{13} = 2.25 ? ; v_{13} = 0.649$

© ICA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/JohnEvans](http://www.durham.ac.uk/JohnEvans)

## Typical values

		O1	O1	O1	O1	O2	O3	O3	O4	Sum
Bi1	dÅ	2.199	2.199	2.236	2.236					2.87
	$v_i$	0.75	0.75	0.68	0.68					1.98
Mg1	dÅ			2.066	2.066	1.980	2.038	2.038		1.98
	$v_i$			0.36	0.36	0.46	0.39	0.39		1.95
Mg2	dÅ			2.066	2.066		2.042	2.042	1.995	
	$v_i$			0.36	0.36		0.39	0.39	0.44	
V1	dÅ					1.688	1.733	1.733	1.684	
	$v_i$					1.36	1.21	1.21	1.38	5.16
Sum				(2*Bi1)	2.16	1.82		1.99	1.82	

- After refinement calculate all bond valence parameters from distances
- e.g.  $\text{BiMg}_2\text{VO}_6$  (see later)

© ICA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/JohnEvans](http://www.durham.ac.uk/JohnEvans)

## Think Beyond Diffraction

(202) (022) (220)

- Best diffraction data tells you  $\text{ZrP}_2\text{O}_7$  is complicated – 50 atoms in the asymmetric unit, perhaps higher
- 2D MQ  $^{31}\text{P}$  solid state NMR **PROVES** it has 136 ( $\pm 0$ )

© ICA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/JohnEvans](http://www.durham.ac.uk/JohnEvans)

## Think Beyond Diffraction

SnP<sub>2</sub>O<sub>7</sub>

ZrW<sub>2</sub>O<sub>8</sub>

ZrV<sub>2</sub>O<sub>7</sub>


© ICA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/JohnEvans](http://www.durham.ac.uk/JohnEvans)

## Case History 1 – BiMg<sub>2</sub>VO<sub>6</sub>

- Simple metal oxide originally published in J. Solid State Chemistry in 1992
- Space group Cmc<sub>2</sub>m
- Straightforward material, simple structure, low R factor no particular reason to think anything might be wrong

CC-BY 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/ds/bio/evans](http://www.durham.ac.uk/ds/bio/evans)

## Case History 1 – BiMg<sub>2</sub>VO<sub>6</sub>




- Simple metal oxide
- Cmc<sub>2</sub>m
- R=2.10%
- Bond distances etc sensible

4+n coord Bi, 5 coordinate Mg, VO<sub>4</sub> tetrahedra

CC-BY 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/ds/bio/evans](http://www.durham.ac.uk/ds/bio/evans)

## Case History 1 – BiMg<sub>2</sub>VO<sub>6</sub>




- Simple metal oxide
- Cmc<sub>2</sub>m
- R=2.10%
- Bond distances etc sensible

4+n coord Bi, 5 coordinate Mg, VO<sub>4</sub> tetrahedra

CC-BY 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/ds/bio/evans](http://www.durham.ac.uk/ds/bio/evans)

## Case History 1 – BiMg<sub>2</sub>VO<sub>6</sub>



- Simple metal oxide
- Pnma
- R=2.13%
- Face index + sadabs
- Everything sensible

4+n coord Bi, 5 coordinate Mg, VO<sub>4</sub> tetrahedra

CC-BY 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/ds/bio/evans](http://www.durham.ac.uk/ds/bio/evans)

## Tedious Detail - 1







		O1	O1	O1	O1	O2	O3	O3	O4	Sum
Bi1	dA	2.199	2.199	2.236	2.236					
	W <sub>i</sub>	0.75	0.75	0.68	0.68					2.87
Mg1	dA			2.066	2.066	1.980	2.038	2.038		
	W <sub>i</sub>			0.36	0.36	0.46	0.39	0.39		1.98
Mg2	dA			2.066	2.066		2.042	2.042	1.995	
	W <sub>i</sub>			0.36	0.36		0.39	0.39	0.44	1.95
V1	dA					1.698	1.733	1.733	1.684	
	W <sub>i</sub>					1.36	1.21	1.21	1.38	5.16
Sum				(2*Bi1)	2.16	1.82		1.99	1.82	

CC-BY 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/ds/bio/evans](http://www.durham.ac.uk/ds/bio/evans)

## Tedious Detail - 2

	Untreated	Face Indexed	SADABS	FI + SADABS
Rint (%)	25.8	12.8	5.2	2.9
No. Reflections	8983	8983	7680	8955
R/wR (%), isotropic refinement	7.20 / 17.53	3.80 / 10.14	5.44 / 11.85	2.43 / 6.36
R/wR (%), anisotropic refinement	6.70 / 16.16	3.19 / 8.47	3.82 / 8.77*	2.21 / 6.03
R/wR (%), aniso ref + optimal weights	6.75 / 10.85	3.14 / 7.05	4.05 / 7.79**	2.13 / 5.27


CC-BY 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/ds/bio/evans](http://www.durham.ac.uk/ds/bio/evans)





## Mo<sub>2</sub>P<sub>4</sub>O<sub>15</sub> – Structure Solution

- Space group Pn
- 441 atoms in asymmetric unit
- How to solve?




ICMA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/john.evans](http://www.durham.ac.uk/john.evans)

## Mo<sub>2</sub>P<sub>4</sub>O<sub>15</sub>


incorrect subcell

$a = 8.31 \text{ \AA}$   
 $b = 6.52 \text{ \AA}$   
 $c = 10.71 \text{ \AA}$   
 $\beta = 107.0^\circ$   
 $V = 555 \text{ \AA}^3$



correct cell


$a = 24.11 \text{ \AA}$   
 $b = 19.53 \text{ \AA}$   
 $c = 25.09 \text{ \AA}$   
 $\beta = 100.0^\circ$   
 $V = 11635 \text{ \AA}^3$



ICMA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/john.evans](http://www.durham.ac.uk/john.evans)

## Mo<sub>2</sub>P<sub>4</sub>O<sub>15</sub> – Structure Solution


- Space group Pn
- 441 atoms in asymmetric unit
- How to solve?



ICMA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/john.evans](http://www.durham.ac.uk/john.evans)

## Distance Least Squares

- We know a lot about the structure!
- 42 MoO<sub>6</sub> octahedra
- 84 P<sub>2</sub>O<sub>7</sub> groups
- We know octahedra share corners with tetrahedra



ICMA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/john.evans](http://www.durham.ac.uk/john.evans)


## Distance Least Squares Refinement

- Structural model has a bond distance of  $d_{\text{model}}$  ?
- Ideal bond distance of  $d_{\text{ideal}}$  ?
- Calculate and minimise  $w(d_{\text{ideal}} - d_{\text{model}})^2$
- Do this for all the bonds/angles known in the structure




ICMA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/john.evans](http://www.durham.ac.uk/john.evans)

## Mo<sub>2</sub>P<sub>4</sub>O<sub>15</sub> – Structure Solution


- Space group Pn
- 441 atoms in asymmetric unit
- Solved by simulated annealing approach in Topas
- R=3.49% for 43738 reflections



ICMA 2005 [john.evans@durham.ac.uk](mailto:john.evans@durham.ac.uk) [www.durham.ac.uk/john.evans](http://www.durham.ac.uk/john.evans)



- ## The Problems
- Getting the right unit cell, pseudo symmetry
  - Getting the right space group
  - Transformation matrices
  - Spotting a chemically nonsensical result
  - Designing a good experiment
- ICMA 2005 [john.arnold@durham.ac.uk](mailto:john.arnold@durham.ac.uk) [www.durham.ac.uk/ajb/ajb.htm](http://www.durham.ac.uk/ajb/ajb.htm)



## Handy Hint

- Inverse of  $\begin{pmatrix} 2 & -1 & 0 \\ 2 & -1 & 1 \\ -2 & 0 & -2 \end{pmatrix}$
- is  $\begin{pmatrix} 1 & -1 & -0.5 \\ 1 & -2 & -1 \\ -1 & 1 & 0 \end{pmatrix}$

ICMA 2005 [john.arnold@durham.ac.uk](mailto:john.arnold@durham.ac.uk) [www.durham.ac.uk/ajb/ajb.htm](http://www.durham.ac.uk/ajb/ajb.htm)

