

	1990		2005		2010		p value
	n	Rate (95% CI)	n	Rate (95% CI)	n	Rate (95% CI)	
(Continued from previous page)							
Globally							
<75 years							
Incidence	6 353 868	159.22 (145.32–174.98)	9 288 048	167.45 (150.96–187.11)	10 469 624	168.75 (152.43–187.09)	0.208
Prevalence	13 234 062	324.26 (288.74–374.96)	20 187 246	358.58 (317.58–412.79)	23 052 804	366.93 (328.04–420.66)	0.086
MIR	..	0.359 (0.318–0.409)	..	0.293 (0.249–0.332)	..	0.254 (0.212–0.287)	<0.001
DALYs lost	63 991 864	1543.96 (1452.03–1728.25)	74 855 520	1326.17 (1172.08–1388.74)	73 293 552	1163.448 (1011.43–1232.19)	<0.001
Mortality	2 301 435	57.38 (54.12–64.27)	2 734 251	49.16 (43.60–51.55)	2 668 499	42.89 (37.65–45.81)	<0.001
≥75 years							
Incidence	3 725 067	3173.50 (2932.14–3422.23)	5 446 077	3082.97 (2819.52–3372.55)	6 424 911	3113.00 (2850.95–3403.57)	0.361
Prevalence	4 681 276	3974.37 (3609.66–4441.23)	8 308 337	4700.18 (4239.37–5256.84)	9 972 153	4835.38 (4382.63–5433.92)	0.005
MIR	..	0.634 (0.575–0.709)	..	0.543 (0.476–0.607)	..	0.500 (0.439–0.560)	<0.001
DALYs	22 018 520	18665.35 (17 464.55–20 408.51)	27 096 178	15 300.36 (13 987.78–16 317.62)	28 938 754	14 053.63 (12 761.98–15 088.12)	<0.001
Mortality	2 359 013	2033.21 (1888.78–2233.65)	2 950 719	1678.65 (1528.60–1807.22)	3 205 682	1545.29 (1412.76–1685.12)	<0.001
All ages							
Incidence	10 078 935	250.55 (229.70–273.25)	14 734 124	255.79 (232.10–283.88)	16 894 536	257.96 (234.40–284.11)	0.335
Prevalence	17 915 338	434.86 (389.45–496.84)	28 495 582	490.13 (436.60–557.52)	33 024 958	502.32 (451.26–572.18)	0.047
MIR	..	0.461 (0.415–0.518)	..	0.386 (0.336–0.432)	..	0.348 (0.299–0.390)	<0.001
DALYs lost	86 010 384	2062.74 (1949.53–2280.29)	101 951 696	1749.59 (1568.67–1830.82)	102 232 304	1554.02 (1373.94–1642.26)	<0.001
Mortality	4 660 449	117.25 (111.51–129.68)	5 684 970	98.53 (89.02–103.86)	5 874 182	88.41 (79.84–94.41)	<0.001

*p value for the difference in age-adjusted rates between 1990 and 2010 only.

Table 1: Age-adjusted annual incidence and mortality rates (per 100 000 person-years), disability-adjusted life-years (DALYs) lost, prevalence (per 100 000 people), and mortality-to-incidence ratio (MIR) by age groups in high-income and low-income and middle-income countries, and globally in 1990, 2005, and 2010